

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07H 21/04, C07K 14/00, 16/00, C12N 15/00, 15/63, 15/85, 15/86, C12Q 1/68, G01N 33/53</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/56755</b> <b>(43) International Publication Date:</b> 28 September 2000 (28.09.00)
<b>(21) International Application Number:</b> PCT/US00/06830 <b>(22) International Filing Date:</b> 16 March 2000 (16.03.00)  <b>(30) Priority Data:</b> 60/125,361 19 March 1999 (19.03.99) US 60/169,910 10 December 1999 (10.12.99) US  <b>(71) Applicant (for all designated States except US):</b> HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hill Drive, Laytonsville, MD 20882 (US). KOMAT-SOULIS, George [US/US]; 9518 Garwood Street, Silver Spring, MD 20901 (US).  <b>(74) Agents:</b> HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> 49 HUMAN SECRETED PROTEINS  <b>(57) Abstract</b> <p>The present invention relates to 49 novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## 49 Human Secreted Proteins

### *Field of the Invention*

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and their production.

### *Background of the Invention*

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include the commercially valuable human insulin, interferon, Factor VIII, human growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of

the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent medical disorders by using secreted proteins or the genes that encode them.

5

### *Summary of the Invention*

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant and synthetic methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders and conditions related to the polypeptides and polynucleotides, and therapeutic methods for treating such disorders and conditions. The invention further relates to screening methods for identifying binding partners of the polypeptides.

15

### *Detailed Description*

#### **Definitions**

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide/sequences of the present invention.

30



In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released  
5 into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000  
10 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding  
15 sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid  
20 sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants  
25 of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig  
30 analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the

ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

- 5 A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC  
10 (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

- Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions.  
15 Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl;  
20 0.2M NaH<sub>2</sub>PO<sub>4</sub>; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

- 25 Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking  
30 reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using oligo dT as a primer).

The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of

ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Many proteins (and translated DNA sequences) contain regions where the amino acid composition is highly biased toward a small subset of the available

residues. For example, membrane spanning domains and signal peptides (which are also membrane spanning) typically contain long stretches where Leucine (L), Valine (V), Alanine (A), and Isoleucine (I) predominate. Poly-Adenosine tracts (polyA) at the end of cDNAs appear in forward translations as poly-Lysine (poly-K) and poly-  
5 Phenylalanine (poly-F) when the reverse complement is translated. These regions are often referred to as "low complexity" regions.

Such regions can cause database similarity search programs such as BLAST to find high-scoring sequence matches that do not imply true homology. The problem is exacerbated by the fact that most weight matrices (used to score the alignments  
10 generated by BLAST) give a match between any of a group of hydrophobic amino acids (L,V and I) that are commonly found in certain low complexity regions almost as high a score as for exact matches.

In order to compensate for this, BLASTX.2 (version 2.0a5MP-WashU) employs two filters ("seg" and "xnu") which "mask" the low complexity regions in a  
15 particular sequence. These filters parse the sequence for such regions, and create a new sequence in which the amino acids in the low complexity region have been replaced with the character "X". This is then used as the input sequence (sometimes referred to herein as "Query" and/or "Q") to the BLASTX program. While this regime helps to ensure that high-scoring matches represent true homology, there is a  
20 negative consequence in that the BLASTX program uses the query sequence that has been masked by the filters to draw alignments.

Thus, a stretch of "X"s in an alignment shown in the following application does not necessarily indicate that either the underlying DNA sequence or the translated protein sequence is unknown or uncertain. Nor is the presence of such  
25 stretches meant to indicate that the sequence is identical or not identical to the sequence disclosed in the alignment of the present invention. Such stretches may simply indicate that the BLASTX program masked amino acids in that region due to the detection of a low complexity region, as defined above. In all cases, the reference sequence(s) (sometimes referred to herein as "Subject", "Sbjct", and/or "S") indicated  
30 in the specification, sequence table (Table 1), and/or the deposited clone is (are) the definitive embodiment(s) of the present invention, and should not be construed as

limiting the present invention to the partial sequence shown in an alignment, unless specifically noted otherwise herein.

### Polynucleotides and Polypeptides of the Invention

#### 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 1

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. bbsl127516 (all information available through the recited accession number is  
10 incorporated herein by reference) which is described therein as " type XVI collagen alpha 1 chain, alpha 1 (XVI) [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

15 >bbsl127516 type XVI collagen alpha 1 chain, alpha 1 (XVI) [human, placenta,  
Peptide Partial, 1186 aa] [Homo sapiens]  
Length = 1186

20 Plus Strand HSPs:

Score = 84 (29.6 bits), Expect = 0.057, Sum P(3) = 0.056  
Identities = 19/49 (38%), Positives = 26/49 (53%), Frame = +3

25 Q: 726 PSQPGWLPSPKRPGRAGRQGLQVLPALRPRPQISNNEQTLGPAWAGDLG 872  
P +PG SP PG+ GR+G Q LP +R P + +G AG+ G  
S: 1066 PGRPGAPGSPGLPGQIGREGRQGLPGVRGLPGTKGEKGDIGIGIAGENG 1114

Score = 62 (21.8 bits), Expect = 0.097, Sum P(4) = 0.092  
Identities = 30/100 (30%), Positives = 40/100 (40%), Frame = +3

30 Q: 573 PHLPRPHRAQNRRGLPQPAEHRVSGAPIHRCRRGY\*W\*PPCGFEGSSPWPHPSPQPGWLPS 752  
P P P Q G+ +P + G P +G PP G +G P P G +  
S: 323 PGQPGPKGEQGPEGVGRPGKPGQPGLP---GVQG---PP-GLKGVQGEPPGRG-VQG 373

35 Q: 753 PK-RPGRAGRQGLQVLPALRPRPQISNNEQTLG-PAWAGDLG 872  
P+ PG G G+Q LP R P + + G P G G  
S: 374 PQGEPGAPGLPGIQGLPGPRGPPGPTGEKGAQGSPPGVKGATG 415

The segments of bbsl127516 that are shown as "S" above are set out as  
40 sequences SEQ ID NO. 109 and/or SEQ ID NO. 111. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 110 and/or SEQ ID NO. 112 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

5 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen 1NFLS and to a lesser extent in Stratagene fetal spleen (#937205); H Macrophage (GM-CSF treated), re-excision; Soares infant brain 1NIB; Human Thymus Tumor; Fetal Liver, subtraction II; Soares adult brain N2b5HB55Y; Human adult testis, large inserts; Early Stage Human Brain;  
10 Human Fetal Heart; CD34 depleted Buffy Coat (Cord Blood), re-excision; Nine Week Old Early Stage Human.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 60 as residues: Gln-43 to Arg-49, Leu-77 to Asn-88.

Many polynucleotide sequences, such as EST sequences, are publicly  
15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention  
20 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1315 of SEQ ID NO:11, b is an integer of 15 to 1329, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

25

## **FEATURES OF PROTEIN ENCODED BY GENE NO: 2**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.  
30 gill150857 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " rev gene product

[Simian immunodeficiency virus]." A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >gi|1150857 rev gene product [Simian immunodeficiency virus]
      >sp|Q87626|Q87626
          GAG (FRAGMENT).
          Length = 65

10      Plus Strand HSPs:

      Score = 65 (22.9 bits), Expect = 2.6, P = 0.92
      Identities = 19/49 (38%), Positives = 23/49 (46%), Frame = +1

15      Q:   229 GTKWSLLPEQKLALIASQNPHPHPGVSAGCRPSK--WTPRDRSGDCLA 369
          GT   LL   K+   I   QNP+P P GS A  R   +   W   R   +   LA
      S:     9 GTDEQLLKAVKIIKILYQNPYPKPAGSRAARRNKRRRWKKRQDQVEGLA 57

```

The segment of gill150857 that is shown as "S" above is set out as sequence SEQ ID NO. 113.

20 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 114 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following  
25 tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Soares\_NhHMPu\_S1; Soares\_fetal\_lung\_NbHL19W; Stratagene placenta (#937225); Nine Week Old Early Stage Human; Human Prostate Cancer, Stage C fraction; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Hemangiopericytoma; Human Placenta; Human 8 Week Whole Embryo; NCI\_CGAP\_Br3; NCI\_CGAP\_Co9;  
30 NCI\_CGAP\_Co12; Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
35 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 820 of SEQ ID NO:12, b



is an integer of 15 to 834, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO: 12, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 3

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares retina N2b4HR and to a lesser extent in Soares fetal liver spleen 1NFLS; Stratagene colon (#937204); Breast Lymph node cDNA library; Human Normal Breast; Soares placenta Nb2HP; Soares infant brain 1NIB; Soares  
 10 ovary tumor NbHOT; Soares retina N2b5HR; pBMC stimulated w/ poly I/C; Soares\_parathyroid\_tumor\_NbHPA; Human Primary Breast Cancer Reexcision; Human Thymus Stromal Cells; Soares breast 2NbHBst; Human Adrenal Gland Tumor; Stratagene liver (#937224); breast lymph node CDNA library; Human Testes Tumor; Soares\_pregnant\_uterus\_NbHPU; Human Microvascular Endothelial Cells,  
 15 fract. A; Human Bone Marrow, treated; Activated T-cell(12h)/Thiouridine-re-excision; T cell helper II; Nine Week Old Early Stage Human; Human Cerebellum; Soares\_multiple\_sclerosis\_2NbHMSP; Primary Dendritic Cells, lib 1; Larynx Tumour; HSC172 cells; Human Lung; Cem cells cyclohexamide treated; B Cell lymphoma; Soares\_NhHMPu\_S1; Soares\_testis\_NHT; Myeloid Progenitor Cell Line;  
 20 Soares\_fetal\_lung\_NbHL19W; Soares\_fetal\_heart\_NbHH19W; Human Ovary; Brain Frontal Cortex, re-excision; Human Chronic Synovitis; Apoptotic T-cell; Human Activated T-Cells; Human Pancreas Tumor; Human Heart; Ulcerative Colitis; Human Whole Six Week Old Embryo; H. Frontal cortex,epileptic,re-excision; PC3 Prostate cell line; Clontech human aorta polyA+ mRNA (#6572); NCI\_CGAP\_Co8;  
 25 NCI\_CGAP\_GC2; NCI\_CGAP\_GCB1; Fetal Heart; Dendritic cells, pooled; Bone marrow; Human Fetal Heart; T Cell helper I; Endothelial-induced; Activated T-Cell (12hs)/Thiouridine labelledEco; CD34 depleted Buffy Coat (Cord Blood), re-excision; Soares\_fetal\_heart\_NbHH19W; Monocyte activated; NCI\_CGAP\_Pr4; Human Testes; Human Endometrial Tumor; Hodgkin's Lymphoma II; Osteoblasts;  
 30 Keratinocyte; Human 8 Week Whole Embryo.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 62 as residues: Asp-21 to Trp-34.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3357 of SEQ ID NO:13, b is an integer of 15 to 3371, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 4

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil2984584 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " P1.11659\_5 protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|2984584 (AC004472) P1.11659_5 [Homo sapiens] >sp|O60377|O60377
P1.11659_5
      (FRAGMENT).
      Length = 188

      Plus Strand HSPs:

      Score = 156 (54.9 bits), Expect = 3.3e-10, P = 3.3e-10
      Identities = 27/47 (57%), Positives = 41/47 (87%), Frame = +1

      Q:   100 QSSKSGKIYLRDVRLLFSRKSMEVDSCAAYELKSYTESPTNPQFSP 240
           +SS+SG++ LH D+RLLFSR+S+E+D+G YEL++ TE+P NP++SP
      S:   140 RSSRSGRLSLHGDIRLLFSRRSLELDTGLPYELQAVTEAPHNPRYSP 186

```

The segment of gil2984584 that is shown as "S" above is set out as sequence SEQ ID NO. 115. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such

activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 116 which corresponds to the Q  
5 sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

When tested against U937 Myeloid cell lines, supernatants removed from cells containing this gene activated the GAS assay. Thus, it is likely that this gene activates myeloid cells through the Jak-STAT signal transduction pathway. The gamma  
10 activating sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and  
15 differentiation of cells.

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_multiple\_sclerosis\_2NbHMSP and to a lesser extent in Soares retina N2b4HR; Soares placenta Nb2HP; Human Kidney Cortex, re-rescue; H. Atrophic Endometrium; Pancreatic Islet; Soares\_parathyroid\_tumor\_NbHPA;  
20 Messangial cell, frac 2; Human Adipose; Hepatocellular Tumor, re-excision; Stratagene fetal retina 937202; Resting T-Cell Library, II; Anergic T-cell.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:14 and may have been publicly available prior to conception of  
25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1080 of SEQ ID NO:14, b  
30 is an integer of 15 to 1094, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 5**

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_fetal\_heart\_NbHH19W and to a lesser extent in  
5 Soares\_NhHMPu\_S1; H. Atrophic Endometrium; Human Amygdala, re-excision;  
Human Fetal Brain; Smooth muscle, control.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of  
10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1229 of SEQ ID NO:15, b  
15 is an integer of 15 to 1243, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 6**

20 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Atrophic Endometrium; Soares infant brain 1NIB.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:16 and may have been publicly available prior to conception of  
25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 804 of SEQ ID NO:16, b  
30 is an integer of 15 to 818, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 7**

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Atrophic Endometrium; Brain frontal cortex.

5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence  
10 would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 833 of SEQ ID NO:17, b is an integer of 15 to 847, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a  
15 + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 8**

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Atrophic Endometrium; Neutrophils IL-1 and LPS  
20 induced.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
25 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 648 of SEQ ID NO:18, b is an integer of 15 to 662, where both a and b correspond to the positions of  
30 nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 9**

It has been discovered that this gene is expressed primarily in H. Atrophic Endometrium.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2240 of SEQ ID NO:19, b is an integer of 15 to 2254, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

15

**FEATURES OF PROTEIN ENCODED BY GENE NO: 10**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil563136 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "unassigned reading frame [Trypanoplasma borreli]." A partial alignment demonstrating the observed homology is shown immediately below.

```

25  >gil563136 unassigned reading frame [Trypanoplasma borreli]
      Length = 68

      Plus Strand HSPs:

30  Score = 63 (22.2 bits), Expect = 3.6, P = 0.97
      Identities = 16/45 (35%), Positives = 27/45 (60%), Frame = +2

      Q:   329 IHFFFFVDITFRIELFVIYCLD*YFVYLGIMKKVCVN-IALLKHMF 463
          ++F+F+  I      +F I+CL+ YF+  I  + +N IALL  +F
35  S:   15 LYFYFYFTI-----IFFIFCLNSYFLMCNIFIYIILNTIALLTQVF 55

```

The segment of gil563136 that is shown as "S" above is set out as sequence SEQ ID NO. 117.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 118 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

5           When tested against K562 leukemia cell lines, supernatants removed from cells containing this gene activated the ISRE assay. Thus, it is likely that this gene activates leukemia cells through the Jak-STAT signal transduction pathway. The interferon-sensitive response element is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a  
10   large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

          When tested against fibroblast cell lines, supernatants removed from cells  
15   containing this gene activated the EGR1 assay. Thus, it is likely that this gene activates fibroblast cells through a signal transduction pathway. Early growth response 1 (EGR1) is a promoter associated with certain genes that induces various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation.

20           It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Atrophic Endometrium; T cell helper II.

          Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of  
25   the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1065 of SEQ ID NO:20, b  
30   is an integer of 15 to 1079, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

**FEATURES OF PROTEIN ENCODED BY GENE NO: 11**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil3127066 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "type 2 orexin receptor [Sus scrofa]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|3127066 (AF059740) type 2 orexin receptor [Sus scrofa]
>sp|O62809|O62809
      TYPE 2 OREXIN RECEPTOR (FRAGMENT).
      Length = 77

Plus Strand HSPs:

Score = 60 (21.1 bits), Expect = 6.0, P = 1.0
Identities = 14/36 (38%), Positives = 23/36 (63%), Frame = +2

Q:  461 SWKIFFGKNVIKATPKIDTVS-GCTILTVICLHVFD 568
      +W  FFG+++ K  P + TVS  ++LT+ C+  DR
S:  14 TW--FFGQSLCKVIPYLQTVSVSVSVLTLSCI-ALDR 47

```

The segment of gil3127066 that is shown as "S" above is set out as sequence SEQ ID NO. 119. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 120 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in H. Atrophic Endometrium.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of



the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 618 of SEQ ID NO:21, b is an integer of 15 to 632, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

## 10 FEATURES OF PROTEIN ENCODED BY GENE NO: 12

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil182221 (all information available through the recited accession number is incorporated herein by reference) which is described therein as a human protein encoded by ORF3. A partial alignment demonstrating the observed homology is shown immediately below.

```

20  >gi|182221 ORF 3 [Homo sapiens] >pir|E41925|E41925 hypothetical protein 3
    -
      human
      Length = 143

25  Plus Strand HSPs:
      Score = 80 (28.2 bits), Expect = 0.00067, Sum P(2) = 0.00067
      Identities = 17/50 (34%), Positives = 30/50 (60%), Frame = +3

30  Q:  1122 NCHGKLEWYRQLWKTAWQFVTT-----YNPSVSLG IY LKEMKT*CSHKN 1256
      +C    +  + LWKT W+F+          ++P++ LLGIY K+ K+ C +K+
      S:   67 HCWWDCKL VQPLWKT VWRFLRDLQLEIPFDPAIPLLGIY PKDYKS-CCYKD 116
  
```

The segment of gil182221 that is shown as "S" above is set out as sequence SEQ ID NO. 121.

35 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 122 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Early Stage Human Brain; Soares infant brain 1NIB and to a lesser extent in Soares\_multiple\_sclerosis\_2NbHMSP; NTERA2 teratocarcinoma cell line+retinoic acid (14 days); Synovial hypoxia-RSF subtracted; Glioblastoma; Human  
 5 Infant Brain; Human Prostate; Stromal cell TF274; H. Frontal cortex,epileptic,re-excision; Soares\_parathyroid\_tumor\_NbHPA.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of  
 10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1478 of SEQ ID NO:22, b  
 15 is an integer of 15 to 1492, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 13

20 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil190475 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "salivary proline-rich  
 25 protein 1 [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

    >gi|190475 salivary proline-rich protein 1 [Homo sapiens]
    >sp|Q15215|Q15215
  30      SALIVARY PROLINE-RICH PROTEIN 1 (FRAGMENT).
        Length = 93

    Plus Strand HSPs:

  35      Score = 69 (24.3 bits), Expect = 0.00017, Sum P(2) = 0.00017
        Identities = 21/51 (41%), Positives = 26/51 (50%), Frame = +2
  
```

Q: 296 P\*GPPSAEAGGHPQSPLPCNP SWGHNGALSA--GGQCPSHLPSSPQQKRPATP 448  
 P GPP E G +PQ P P P+ G+ A GQ P P PQ +P+ P  
 S: 43 PQGFPQQE-GNNPQGPPF--PAGGNPQQPQAPPAGQ-PQGPPRPPQGDKPSRP 91

5

The segment of gil190475 that is shown as "S" above is set out as sequence SEQ ID NO. 123. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 124 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_pregnant\_uterus\_NbHPU; Early Stage Human Brain and to a lesser extent in Soares\_fetal\_heart\_NbHH19W; Nine Week Old Early Stage Human; Soares ovary tumor NbHOT; Human Primary Breast Cancer; Human Hypothalamus,schizophrenia, re-excision; NCI\_CGAP\_GCB1; Human endometrial stromal cells-treated with estradiol; Pancreas normal PCA4 No; Human endometrial stromal cells; Spinal Cord, re-excision; Human Chronic Synovitis; 12 Week Old Early Stage Human, II; Human Primary Breast Cancer Reexcision; Soares breast 2NbHBst; 12 Week Old Early Stage Human; Soares breast 3NbHBst; Soares\_senescent\_fibroblasts\_NbHSF; Human Fetal Lung III; Human Placenta; Endothelial cells-control; Smooth muscle,control; Bone Marrow Cell Line (RS4,11).

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 72 as residues: Gly-37 to Pro-45.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:23 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention

are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 624 of SEQ ID NO:23, b is an integer of 15 to 638, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 14

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen 1NFLS and to a lesser extent in Early Stage Human Brain; Soares infant brain 1NIB; Soares\_NhHMPu\_S1; Human Soleus; Stratagene lung (#937210); Soares placenta Nb2HP; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Stratagene HeLa cell s3 937216; Soares\_placenta\_8to9weeks\_2NbHP8to9W; Human Adult Pulmonary, re-excision; Stratagene hNT neuron (#937233); Stratagene NT2 neuronal precursor 937230; Human Fetal Brain, normalized CO; Human Quadriceps; Salivary Gland; Soares\_NhHMPu\_S1; Soares\_total\_fetus\_Nb2HF8\_9w; Human Amygdala, re-excision; Synovial IL-1/TNF stimulated; Human Umbilical Vein, Endo. remake; Healing groin wound, 7.5 hours post incision; Human endometrial stromal cells-treated with progesterone; Human Adult Small Intestine; Human Infant Brain; Human Prostate; H. Kidney Medulla, re-excision; Soares\_fetal\_heart\_NbHH19W; Macrophage-oxLDL; Human Hypothalamus, Schizophrenia; Soares adult brain N2b5HB55Y; Human Thymus Stromal Cells; Human Adrenal Gland Tumor; NTERA2, control; Soares\_fetal\_heart\_NbHH19W; Stratagene HeLa cell s3 937216; Smooth muscle, serum induced, re-exc; Soares\_NhHMPu\_S1; Human Placenta; H Macrophage (GM-CSF treated), re-excision; Soares melanocyte 2NbHM; Soares\_fetal\_lung\_NbHL19W.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention

are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 930 of SEQ ID NO:24, b is an integer of 15 to 944, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 15

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil1200224 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "coat protein [Bacillus subtilis]." A partial alignment demonstrating the observed homology is shown immediately below.

```

15      >gi|1200224 coat protein [Bacillus subtilis] >gi|1146180 spore coat
      protein
      [Bacillus subtilis] >gnl|PID|e1183666 spore coat protein
      (inner)
20      [Bacillus subtilis] >pir|D27393|D27393 11K inner spore coat
      protein
      cotD - Bacillus subtilis >sp|P07791|COTD_BACSU SPORE COAT
      PROTEIN
25      D.
      Length = 75

      Minus Strand HSPs:

      Score = 56 (19.7 bits), Expect = 5.8, P = 1.0
30      Identities = 9/28 (32%), Positives = 16/28 (57%), Frame = -1

      Q:   375 TLLINIHPYDDVDNHEHYSIPIHIIAH 292
            T++ +IHP      + NH+H+   +H   H
      S:   27  TIVPHIHPQHTTNVNHQHFQ-HVHYFPH 53
35

```

The segment of gil1200224 that is shown as "S" above is set out as sequence SEQ ID NO. 125.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 126 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen 1NFLS and to a lesser extent in Early Stage Human Brain; Soares infant brain 1NIB; Soares\_NhHMPu\_S1; Human Soleus; Stratagene lung (#937210); 12 Week Old Early Stage Human; Osteoblasts; Soares placenta Nb2HP; Soares retina N2b4HR; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Synovial IL-1/TNF stimulated; Stratagene HeLa cell s3 937216; Stratagene ovarian cancer (#937219); Myeloid Progenitor Cell Line; Synovial Fibroblasts (II1/TNF), subt; Human Adult Small Intestine; Human Placenta; Soares breast 3NbHBst; Soares melanocyte 2NbHM; Soares\_placenta\_8to9weeks\_2NbHP8to9W; Human Fetal Heart; Human Adult Pulmonary, re-excision; Spleen, Chronic lymphocytic leukemia; Nine Week Old Early Stage Human; Human Fetal Brain, normalized 50021F; Human Fetal Brain, normalized CO; Stratagene hNT neuron (#937233); Stratagene NT2 neuronal precursor 937230; Human Fetal Brain, normalized AC5002; Prostate; Human Kidney; Human White Adipose; human colon cancer; Human Lung; Human Quadriceps; Salivary Gland; HSA 172 Cells; Soares\_fetal\_heart\_NbHH19W; Soares\_NhHMPu\_S1; Soares\_total\_fetus\_Nb2HF8\_9w; Human Amygdala, re-excision; Human Hypothalamus, schizophrenia, re-excision; Stromal-Osetoclastoma; Hepatocellular Tumor; human corpus colosum; Alzheimers, spongy change; Human Umbilical Vein, Endo. remake; Healing groin wound, 7.5 hours post incision; Synovial hypoxia-RSF subtracted; LNCAP prostate cell line; Human Stomach, re-excision; Human endometrial stromal cells-treated with progesterone; Healing groin wound, 6.5 hours post incision; Synovial hypoxia; Soares\_NhHMPu\_S1; Human Ovary; wilm's tumor; Human Infant Brain; Human Prostate; H. Kidney Medulla, re-excision; T-Cell PHA 16 hrs; Human Brain, Striatum; Human Uterine Cancer; Human Ovarian Cancer Reexcision; Macrophage-oxLDL; Human Hypothalamus, Schizophrenia; Human Hippocampus; Soares adult brain N2b5HB55Y; Human Thymus Stromal Cells; Human Adrenal Gland Tumor; Human Whole Six Week Old Embryo; Soares\_fetal\_heart\_NbHH19W; H. Frontal cortex, epileptic, re-excision; NTERA2, control; Smooth muscle, serum induced, re-exc; Human Gall Bladder; PC3 Prostate cell line; H Macrophage (GM-CSF treated), re-excision; Human Fetal Lung III; Endothelial-induced; Human Endometrial Tumor; Stratagene HeLa cell s3 937216; Human 8 Week Whole Embryo; Soares\_fetal\_lung\_NbHL19W.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 394 of SEQ ID NO:25, b is an integer of 15 to 408, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 16

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil403460 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "transformation-related protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi|403460 transformation-related protein [Homo sapiens] >sp|Q15662|Q15662
TRANSFORMATION-RELATED PROTEIN (FRAGMENT).
Length = 368
```

Minus Strand HSPs:

```
Score = 114 (40.1 bits), Expect = 0.0074, P = 0.0074
Identities = 29/57 (50%), Positives = 33/57 (57%), Frame = -1
```

```
Q: 865 GPKQSSRLGLLGSDQGHTPPHANFFVEVESCY--VAQAGLELLGSSDPPTLA-PK 704
    G K+ S L L SWD H PP +F VE+ + QAGLELL SS PPT A PK
S: 26 GFKRFSCSLPSSWDYRHVPPRQVHFVFSVETGFHRAGQAGLELLTSSVPPTSAPFK 82
```

The segment of gil403460 that is shown as "S" above is set out as sequence SEQ ID NO. 127. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such

activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 128 which corresponds to the Q  
5 sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in Early Stage Human Brain.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.  
10 75 as residues: Leu-42 to Lys-50.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
15 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1673 of SEQ ID NO:26, b is an integer of 15 to 1687, where both a and b correspond to the positions of  
20 nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 17**

The computer algorithm BLASTX has been used to determine that the  
25 translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. pir|A23925|A23925 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " proline-rich phosphoprotein - crab-eating macaque." A partial alignment demonstrating the  
30 observed homology is shown immediately below.

```
>pir|A23925|A23925 proline-rich phosphoprotein - crab-eating macaque
```



Length = 115

Minus Strand HSPs:

5      Score = 83 (29.2 bits), Expect = 0.072, P = 0.070  
       Identities = 27/91 (29%), Positives = 36/91 (39%), Frame = -2

10      Q:    645 TEKEAGAQVLMPRHKGSPRQAEISS-PPGDGAQ---PQHSVLQSGGAQHGANPLHRHPWG 478  
               +++E    Q +    H+G P    +LS    P GDG +    PQ            GG    H    P    P  
       S:    15 SDEEDSDQFIDEAHQGPPLGGQLSKHPAGDGNKDDGPQQKPPHHGGHHHRPPPTGEPQN 74

      Q:    477 SS----HTSCRIPTSPTRGQRGRAKPSPSLKP 397  
               +            H        RPT P        G        PS        +P  
       S:    75 TQKPGGHHHRPTPPPGKPEGSPFPSQGSRP 105

15      The segment of pirlA23925|A23925 that is shown as "S" above is set out as sequence SEQ ID NO. 129. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for  
 20      determining such activities are also known in the art some of which have been described elsewhere herein.

      Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 130 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the  
 25      computer are, of course, removed).

      It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen 1NFLS and to a lesser extent in Human Infant Brain; Soares\_fetal\_heart\_NbHH19W; Primary Dendritic Cells, lib 1; Soares infant brain 1NIB; Human adult testis, large inserts; Colon Normal II; Early  
 30      Stage Human Brain; Human Synovial Sarcoma; Bone marrow; Spleen, Chronic lymphocytic leukemia.

      Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 76 as residues: Arg-18 to Lys-26, Gly-35 to Ala-42, Gln-61 to Gly-67.

      Many polynucleotide sequences, such as EST sequences, are publicly  
 35      available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention

are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1175 of SEQ ID NO:27, b is an integer of 15 to 1189, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

### **FEATURES OF PROTEIN ENCODED BY GENE NO: 18**

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Soares adult brain N2b4HB55Y; Soares\_multiple\_sclerosis\_2NbHMSP; Stratagene lung carcinoma 937218; Human Amygdala, re-excision; Glioblastoma; Early Stage Human Brain.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1806 of SEQ ID NO:28, b is an integer of 15 to 1820, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

### **FEATURES OF PROTEIN ENCODED BY GENE NO: 19**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil30099 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " collagen N-prepropeptide (aa -22 to 72) [Homo sapiens] " A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|30099 collagen N-prepropeptide (aa -22 to 72) [Homo sapiens]
>sp|Q14057|Q14057 PRO-ALPHA-2(I) COLLAGEN N-PREPROPEPTIDE
5  PRECURSOR
    (FRAGMENT) .
    Length = 94

    Minus Strand HSPs:
10  Score = 79 (27.8 bits), Expect = 0.17, P = 0.15
    Identities = 22/46 (47%), Positives = 24/46 (52%), Frame = -1

    Q:  235 GDRG--GEVGPASHPGDDKGMKVTPVVKLQPPIGWKGWGGKFSARY 101
        GDRG GE GP PG D G P PP G G GG F+A+Y
15  S:  37 GDRGPRGERGPPGPPGRDGGDGGPPGPPGPPGPPGPPGLGNFAAQY 82

```

The segment of gil30099 that is shown as "S" above is set out as sequence SEQ ID NO. 131. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 132 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human 8 Week Whole Embryo and to a lesser extent in Human Rhabdomyosarcoma; Early Stage Human Brain; normalized infant brain cDNA; Soares fetal liver spleen 1NFLS; Smooth muscle-ILb induced; Soares adult brain N2b4HB55Y; Human Hypothalamus, schizophrenia, re-excision; Human Fetal Brain; Human Whole Six Week Old Embryo; H. Frontal cortex, epileptic, re-excision; Stratagene fetal retina 937202; Human Synovial Sarcoma; Soares\_testis\_NHT; Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence

would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1559 of SEQ ID NO:29, b is an integer of 15 to 1573, where both a and b correspond to the positions of  
 5 nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 20

The computer algorithm BLASTX has been used to determine that the  
 10 translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil3150526 (all information available through the recited accession number is incorporated herein by reference) which is described therein as a *C. elegans* protein. A partial alignment demonstrating the observed homology is shown immediately  
 15 below.

```
>gi|3150526 (AF067219) No definition line found [Caenorhabditis elegans]
>sp|O61782|O61782 R12E2.14 PROTEIN.
Length = 74
```

20 Minus Strand HSPs:

```
Score = 64 (22.5 bits), Expect = 5.7, P = 1.0
Identities = 18/50 (36%), Positives = 24/50 (48%), Frame = -1
```

```
25 Q: 245 LGGAGITSARPCPSQEV*GDTMDGD*GPKVPASLPPQKPHSQAS PQSRA 96
    LGG G + CP + G+ G GP+ A+ PP P A+PQ A
S: 21 LGGGG-GAGGCCPPAQP GCGNPCQGA-GPQYAAAPPPPPQYAAAPQGGA 68
```

30 The segment of gil3150526 that is shown as "S" above is set out as sequence SEQ ID NO. 133.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 134 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the  
 35 computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares retina N2b4HR; Soares infant brain 1NIB and to a lesser extent in Soares\_fetal\_heart\_NbHH19W; Soares\_pregnant\_uterus\_NbHPU;

Human Chondrosarcoma; Soares adult brain N2b5HB55Y; Early Stage Human Brain; Soares melanocyte 2NbHM; Soares fetal liver spleen 1NFLS; Human Lung Cancer, subtracted; LNCAP + 30nM R1881; Whole 6 Week Old Embryo; Human T-cell lymphoma, re-excision; Human Soleus; Glioblastoma; Human Osteosarcoma; Human  
 5 Infant Brain; Human Brain, Striatum; Human Fetal Kidney; NCI\_CGAP\_GCB1; Apoptotic T-cell; Human Hippocampus; Human Fetal Brain; Macrophage-oxLDL, re-excision; Brain frontal cortex; Dendritic cells, pooled; Human Synovial Sarcoma; Human Fetal Lung III; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Spleen, Chronic lymphocytic leukemia; Human Cerebellum; Soares placenta Nb2HP.

10 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 79 as residues: Gln-35 to Gly-58, Asn-109 to Cys-118, Thr-122 to Ser-132.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:30 and may have been publicly available prior to conception of  
 15 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1314 of SEQ ID NO:30, b  
 20 is an integer of 15 to 1328, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 21

25 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. bbs1151373 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "influenza virus  
 30 hemagglutinin 5' epitope tag=fusion protein {N-terminal} [Saccharomyces cerevisiae]." A partial alignment demonstrating the observed homology is shown immediately below.

```

    >bbs|151373 influenza virus hemagglutinin 5' epitope tag=fusion protein
      (N-terminal) [Saccharomyces cerevisiae=yeast, YCpIF15,16,17
5  cloning      vector, Peptide Plasmid Synthetic Partial, 50 aa]
    [Saccharomyces
      cerevisiae]
      Length = 50

10      Plus Strand HSPs:

      Score = 72 (25.3 bits), Expect = 0.16, P = 0.15
      Identities = 15/18 (83%), Positives = 15/18 (83%), Frame = +1

15      Q:   520 QPGGXTSSRAAXNAVELQ 573
              QPGG TSSRAA AVELQ
              S:   31 QPGGSTSSRAAATAVELQ 48

```

The segment of bbs|151373 that is shown as "S" above is set out as sequence  
 20 SEQ ID NO. 135.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 136 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

25 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Early Stage Human Brain; Soares infant brain 1NIB.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:31 and may have been publicly available prior to conception of  
 30 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 576 of SEQ ID NO:31, b  
 35 is an integer of 15 to 590, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 22

40 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting

example, the sequence accessible through the following database accession no. bbsl151371 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "influenza virus hemagglutinin 5' epitope tag=fusion protein {N-terminal, frame 1} [Saccharomyces cerevisiae]." A partial alignment demonstrating the observed homology is shown immediately below.

```

10      >bbsl151371 influenza virus hemagglutinin 5' epitope tag=fusion protein
      {N-terminal, frame 1} [Saccharomyces cerevisiae=yeast,
      YCpIF15.16,17 cloning vector, Peptide Plasmid Synthetic
Partial, 45
      aa] [Saccharomyces cerevisiae]
      Length = 45

15      Minus Strand HSPs:

      Score = 48 (16.9 bits), Expect = 0.39, Sum P(2) = 0.32
      Identities = 9/9 (100%), Positives = 9/9 (100%), Frame = -2

20      Q:      28 GDPLVLERP 2
      GDPLVLERP
      S:      28 GDPLVLERP 36

      Score = 39 (13.7 bits), Expect = 0.39, Sum P(2) = 0.32
      Identities = 8/18 (44%), Positives = 12/18 (66%), Frame = -1

25      Q:      77 HLSSSME-EAVPNSCSPG 27
      + SS++      + NSCSPG
      S:      11 YASSTVSISLISNSCSPG 28

30

```

The segments of bbsl151371 that are shown as "S" above are set out as sequences SEQ ID NO. 137 and/or SEQ ID NO. 139.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 138 and/or SEQ ID NO. 140 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Early Stage Human Brain.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 81 as residues: Asp-31 to Pro-36.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 738 of SEQ ID NO:32, b is an integer of 15 to 752, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 23

10       The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. bbs1156631 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "

15       NF2=neurofibromatosis type 2 [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>bbs|156631 NF2=neurofibromatosis type 2 (alternatively spliced, form A4)
[human, colorectal cancer, Peptide Partial Mutant, 46 aa]
[ Homo sapiens] >pir|I54375|I54375 gene NF2 protein - human
(fragment)
>gi|463121 NF2 [Homo sapiens] (SUB 8-46)
Length = 46

Plus Strand HSPs:

Score = 103 (36.3 bits), Expect = 0.00046, P = 0.00046
Identities = 22/34 (64%), Positives = 24/34 (70%), Frame = +2

Q: 1625 FKTESHSVPQAGMQ*CDLGLSLQSPPPRFKHFSCSL 1726
      F ES SV AG+Q DLG LQP PP+FK FSCSL
S: 4 FNCESCSVTLAGVQWRDLGLLQPLPPKFKRFSCSL 37

Minus Strand HSPs:

Score = 123 (43.3 bits), Expect = 3.6e-06, P = 3.6e-06
Identities = 24/36 (66%), Positives = 25/36 (69%), Frame = -2

Q: 1830 IAQAGVQWRDLGSPQTPLPLGLKRFSCSLSLTSSSWDYR 1723
      + AGVQWRDLG Q P KRFSCSL SSWDYR
S: 11 VTLAGVQWRDLGLLQPLPPKFKRFSCSLSFSSWDYR 46

```

The segments of bbs|15663| that are shown as “S” above are set out as  
45 sequences SEQ ID NO. 141 and/or SEQ ID NO. 143. Based on the structural



similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

5 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 142 and/or SEQ ID NO. 144 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following  
10 tissues/cDNA libraries: Early Stage Human Brain.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:33 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically  
15 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1853 of SEQ ID NO:33, b is an integer of 15 to 1867, where both a and b correspond to the positions of  
20 nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 24

The computer algorithm BLASTX has been used to determine that the  
25 translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi473351 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " Ec protein  
homologue [Arabidopsis thaliana]." A partial alignment demonstrating the observed  
30 homology is shown immediately below.

```
>gi|473351 Ec protein homologue [Arabidopsis thaliana]  
>sp|Q42258|EC3_ARATH EC
```

PROTEIN HOMOLOG 3 (FRAGMENT).  
Length = 82

Plus Strand HSPs:

Score = 65 (22.9 bits), Expect = 0.44, P = 0.36  
Identities = 14/40 (35%), Positives = 19/40 (47%), Frame = +2

Q: 212 CTLPQARGGFSPHPLCPPNLGCTF\*EGDSPCPCPLPQNKT 331  
C + +A G H +CP C G +PC CP Q +T  
S: 29 CRMREASAGDQGHMVCPCGEHC-----GCNPCNCPKTQTQT 64

The segment of gil473351 that is shown as "S" above is set out as sequence SEQ ID NO. 145. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 146 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Early Stage Human Brain; Soares placenta Nb2HP and to a lesser extent in Hemangiopericytoma; Soares\_fetal\_heart\_NbHH19W; A1-CELL LINE; Soares infant brain 1NIB; Soares ovary tumor NbHOT; Pancreatic Islet; Soares\_NhHMPu\_S1; Stratagene hNT neuron (#937233); Human Pineal Gland; Human Neutrophil; Human Fetal Dura Mater; Human Hypothalamus, Schizophrenia; Human Chondrosarcoma; Human Whole Six Week Old Embryo; CD34 depleted Buffy Coat (Cord Blood), re-excision; Hodgkin's Lymphoma II; neutrophils control; Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention

are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 405 of SEQ ID NO:34, b is an integer of 15 to 419, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 25**

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Soares\_fetal\_heart\_NbHH19W; Soares\_NhHMPu\_S1; Early Stage Human Brain; PERM TF274; Pancreas Islet Cell Tumor; Soares\_multiple\_sclerosis\_2NbHMSP; prostate-edited; Soares\_NhHMPu\_S1; Stratagene hNT neuron (#937233).

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:35 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 935 of SEQ ID NO:35, b is an integer of 15 to 949, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 26**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gill1654230 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "nitrogen regulatory factor NRFA [*Penicillium urticae*]." A partial alignment demonstrating the observed homology is shown immediately below.

>gi|1654230 nitrogen regulatory factor NRFA [Penicillium urticae]  
 >sp|Q92269|NRFA\_PENUR NITROGEN REGULATORY PROTEIN NRFA.  
 Length = 865

5

Plus Strand HSPs:

Score = 80 (28.2 bits), Expect = 9.4, Sum P(2) = 1.0  
 Identities = 31/98 (31%), Positives = 43/98 (43%), Frame = +1

10

Q: 967 LDGPAEGRQG-----TQISSSRCPSIHRVISRLPHSKDIQNAEGAHRSAVSGVT 1116  
 +DGP E QG +S R N R ++P + NA R +++G  
 S: 562 MDGPPEWNQGGTLGRGHGSVASVSDVRNQNQDPRRYGKVPRTASTPNAAALLRQSLNGSA 621

15

Q: 1117 SQPPSL\*PLEAHSP--GLSF\*TPSXGLPGCPLGTA-GPPS 1227  
 S PP+ P + P GLS PS PG P G+ G P+  
 S: 622 SGPPTNHSPSTPPESGLSSAVPSR--PGSPGGSKNGDPN 659

The segment of gi1654230 that is shown as "S" above is set out as sequence  
 20 SEQ ID NO. 147.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 148 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

25 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares melanocyte 2NbHM and to a lesser extent in Soares fetal liver spleen 1NFLS; Soares placenta Nb2HP; Stratagene placenta (#937225); NCI\_CGAP\_Co3; Human Colon Cancer, re-excision; Soares\_fetal\_heart\_NbHH19W; Soares\_pregnant\_uterus\_NbHPU; Soares\_multiple\_sclerosis\_2NbHMSP; Nine Week  
 30 Old Early Stage Human; Soares infant brain 1NIB; Activated T-Cells, 4 hrs, subtracted; NCI\_CGAP\_Br2; NCI\_CGAP\_Pr22; NCI\_CGAP\_Br1.1; Tongue Tumour; NCI\_CGAP\_Lu5; NCI\_CGAP\_Kid6; HL-60, RA 4h, Subtracted; prostate-edited; Healing Abdomen wound, 70&90 min post incision; H. Epididymus, caput & corpus; Human Liver; Human Soleus; Human Tonsils, Lib 2; Human Lung Cancer, re-  
 35 excision; Hepatocellular Tumor; Breast Cancer cell line, MDA 36; Healing groin wound, 7.5 hours post incision; Human endometrial stromal cells-treated with progesterone; wilm's tumor; Human Adult Small Intestine; Spinal Cord, re-excision; Clontech human aorta polyA+ mRNA (#6572); Monocyte activated, re-excision; Human Fetal Kidney; Human Heart; Macrophage-oxLDL; Human Chondrosarcoma;  
 40 Human Adrenal Gland Tumor; Soares\_fetal\_lung\_NbHL19W; Smooth muscle, serum induced, re-exc; Pancreas Islet Cell Tumor; 12 Week Old Early Stage Human; Colon

Normal II; breast lymph node CDNA library; Human adult (K.Okubo); Human pancreatic islet; Human placenta polyA+ (TFujiwara); Normal colon; Human Fetal Heart; Activated T-Cell (12hs)/Thiouridine labelledEco; Endothelial cells-control; Monocyte activated; HUMAN B CELL LYMPHOMA; Spleen, Chronic lymphocytic leukemia; Human Testes; Hodgkin's Lymphoma II; Keratinocyte; Human 8 Week Whole Embryo.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1621 of SEQ ID NO:36, b is an integer of 15 to 1635, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 27

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil288145 (all information available through the recited accession number is incorporated herein by reference) which is described therein as a putative ORF encoding a human protein. A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|288145 put. ORF [Homo sapiens] >pir|I38022|I38022 hypothetical protein
-
      human
      Length = 196

      Plus Strand HSPs:

      Score = 251 (88.4 bits), Expect = 8.4e-20, P = 8.4e-20
      Identities = 51/82 (62%), Positives = 58/82 (70%), Frame = +2

      Q:  1424 HLKSGI*DQPGQHGETPSLLKIQKLARHGSRSL*SQLFGRLRQENHLNLGGGGCSELRSR 1603

```

HL+SG+ D PGQHG+ PSLLKIQ+LA HG R L SQL RLRQENHLN GG GCSE +S  
 S: 3 HLRSGVQDYPGQHGKIPSLKIQELAGHGGRCLQSQLLRRLRQENHLNSGGRGCSEPKSH 62  
 Q: 1604 NCAPAWATRAKLHLKE-KTKQD 1666  
 C PAW T K+ KTK +  
 S: 63 LCIPAWVTEGDSVSKQNKTKNE 84

The segment of gil288145 that is shown as "S" above is set out as sequence SEQ ID NO. 149.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 150 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Cerebellum; H. Epididymus, caput & corpus; 12 Week Old Early Stage Human, II.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2400 of SEQ ID NO:37, b is an integer of 15 to 2414, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 28

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil348245 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " protein

serine/threonine kinase [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

5  >gi|348245 protein serine/threonine kinase [Homo sapiens]
   >pir|I78885|I78885
       protein serine/threonine kinase - human
       Length = 841

10  Minus Strand HSPs:

   Score = 124 (43.7 bits), Expect = 0.00055, P = 0.00055
   Identities = 27/46 (58%), Positives = 32/46 (69%), Frame = -1

15  Q:   896 EQGLALLPRLE--GVITAHCILQLLGS RDPPTSASQSAGVTGTYHH 765
       +Q LAL P+LE  G I AH  L+LLGS D P SAS+ AG+TG  HH
   S:   455 DQSLALSPKLECSGTILAHSNLRLLGSSDSPASASRVAGITGVCHH 500

```

The segment of gil348245 that is shown as "S" above is set out as sequence SEQ ID NO. 151. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 152 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in *H. Epididymus*, cauda.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:38 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 915 of SEQ ID NO:38, b is an integer of 15 to 929, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:38, and where b is greater than or equal to a + 14.

## FEATURES OF PROTEIN ENCODED BY GENE NO: 29

5           The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil512350 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "ninF protein  
10 [Bacteriophage P22]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|512350 ninF protein [Bacteriophage P22] >sp|Q38666|Q38666 NINF
15 PROTEIN.
      Length = 58

      Plus Strand HSPs:

      Score = 66 (23.2 bits), Expect = 3.1, P = 0.96
20      Identities = 16/47 (34%), Positives = 24/47 (51%), Frame = +1

      Q:   361 PSQATPASLDQDKWAAQQPVFNKCGQKISTHSTEYHCCDGDCAVAVV 501
           PSQ   SL   K + ++ +   CGQK+  H  E H C+G C  ++
      S:    4 PSQ----SLQYQKESVERALTCANCGQKL--HVLEVHVCEGGCAELM 44
25

```

The segment of gil512350 that is shown as "S" above is set out as sequence SEQ ID NO. 153.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 154 which corresponds to the Q  
30 sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_fetal\_liver\_spleen\_1NFLS\_S1; H. Epididymus, cauda and to a lesser extent in Rejected Kidney, lib 4; Human B Cell Lymphoma.

35           Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:39 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically



excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1111 of SEQ ID NO:39, b is an integer of 15 to 1125, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 30

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil3822194 (all information available through the recited accession number is incorporated herein by reference) which is described therein as an E. coli protein. A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|3822194 (AF074613) unknown [Escherichia coli O157:H7]
>sp|G3822194|G3822194
      HYPOTHETICAL 9.4 KD PROTEIN.
      Length = 86

      Plus Strand HSPs:

      Score = 62 (21.8 bits), Expect = 6.0, P = 1.0
      Identities = 15/27 (55%), Positives = 15/27 (55%), Frame = +2

      Q:   611 FSMRLTSLPPCAGNATDFVSSVFFFP 691
           FS RL  LPP AG  T  V S F F C
      S:   2 FSFRL--LPPLAGRLTSSVKSAFQFEC 26

```

The segment of gil3822194 that is shown as "S" above is set out as sequence SEQ ID NO. 155.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 156 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Stratagene lung (#937210) and to a lesser extent in Soares adult brain N2b4HB55Y; Soares\_fetal\_heart\_NbHH19W; Human Bone Marrow,

treated; Soares placenta Nb2HP; Soares infant brain 1NIB; Synovial IL-1/TNF stimulated; Human Brain, Striatum; Early Stage Human Brain; Human Amygdala; NCI\_CGAP\_GCB1; NCI\_CGAP\_Brn23; Stratagene neuroepithelium (#937231); Keratinocyte.

- 5 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 89 as residues: Val-40 to Cys-45, Lys-58 to Thr-64.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1107 of SEQ ID NO:40, b is an integer of 15 to 1121, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 31

- 20 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil288145 (all information available through the recited accession number is incorporated herein by reference) which is described therein as a putative ORF encoding a human protein. A partial alignment demonstrating the observed homology is shown immediately below.

```

30 >gi|288145 put. ORF [Homo sapiens] >pir|I38022|I38022 hypothetical protein
      human
      Length = 196

      Plus Strand HSPs:

35 Score = 253 (89.1 bits), Expect = 5.0e-20, P = 5.0e-20
      Identities = 49/67 (73%), Positives = 55/67 (82%), Frame = +1

      Q: 1879 LRSGVRDQPGQYGETPSLLKIQKLARHGGICMCSQLLGRRLRQENCLNPGGRVCSEPRSHR 2058

```

S:           LRSGV+D PGQ+G+ PSLKIQ+LA HGG C+ SPLL RLRQEN LN GGR CSEP+SH  
           4 LRSGVQDYPGQHGKIPSLKIQELAGHGGRCLQSPLLRLRQENHLNSGGRGCSEPKSHL 63

Q: 2059 CTTAWVT 2079

5           C   AWVT  
           S:    64 CIPAWVT 70

The segment of gil288145 that is shown as "S" above is set out as sequence SEQ ID NO. 157.

10           Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 158 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

          It has been discovered that this gene is expressed primarily in the following  
 15 tissues/cDNA libraries: Human Whole Six Week Old Embryo; Endothelial cells-control.

          Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:41 and may have been publicly available prior to conception of  
 20 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2154 of SEQ ID NO:41, b  
 25 is an integer of 15 to 2168, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 32**

30           It has been discovered that this gene is expressed primarily in Endothelial cells-control.

          Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:42 and may have been publicly available prior to conception of  
 35 the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2091 of SEQ ID NO:42, b is an integer of 15 to 2105, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 33

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. dbjllD84488\_1 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "small GTP-binding protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>dbjllD84488_1 small GTP-binding protein [Homo sapiens] >sp|O14966|O14966
SMALL
      GTP-BINDING PROTEIN.
      Length = 203

      Minus Strand HSPs:

Score = 687 (241.8 bits), Expect = 5.7e-67, P = 5.7e-67
Identities = 131/131 (100%), Positives = 131/131 (100%), Frame = -3

Q: 1333 GQERFTSMTRLYYRDASACVIMFDVTNATTFSNSQQRWKQDLDSKLTLPNGEPVPCLLLAN 1154
      GQERFTSMTRLYYRDASACVIMFDVTNATTFSNSQQRWKQDLDSKLTLPNGEPVPCLLLAN
S: 66 GQERFTSMTRLYYRDASACVIMFDVTNATTFSNSQQRWKQDLDSKLTLPNGEPVPCLLLAN 125

Q: 1153 KCDLSPWAVSRDQIDRFSKENGFTGWTETSVKENKNINEAMRVLIEKMMRNSTEDIMSLS 974
      KCDLSPWAVSRDQIDRFSKENGFTGWTETSVKENKNINEAMRVLIEKMMRNSTEDIMSLS
S: 126 KCDLSPWAVSRDQIDRFSKENGFTGWTETSVKENKNINEAMRVLIEKMMRNSTEDIMSLS 185

Q: 973 TQGDYINLQTK 941
      TQGDYINLQTK
S: 186 TQGDYINLQTK 196

```

The segment of dbjllD84488\_1 that is shown as "S" above is set out as sequence SEQ ID NO. 159. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for

determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 160 which corresponds to the Q  
5 sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Endothelial cells-control; CD34 positive cells (Cord Blood); Osteoblasts and to a lesser extent in Human Umbilical Vein Endothelial Cells, fract.  
10 A; Human Osteoclastoma, re-excision; Macrophage (GM-CSF treated); NTERA2, control; Human Gall Bladder; Anergic T-cell; Human Amygdala; Human Microvascular Endothelial Cells, fract. A; Human Bone Marrow, treated; T cell helper II; NCI\_CGAP\_Kid3; Soares\_senescent\_fibroblasts\_NbHSF.

Many polynucleotide sequences, such as EST sequences, are publicly  
15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:43 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention  
20 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1477 of SEQ ID NO:43, b is an integer of 15 to 1491, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.

25

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 34**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.  
30 gil1181442 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " a279R [Paramecium

bursaria Chlorella virus 1]." A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >gi|1181442 a279R [Paramecium bursaria Chlorella virus 1] >gi|1181442
a279R
      [Paramecium bursaria Chlorella virus 1] >sp|Q84596|Q84596
GENOME,
      PARTIAL SEQUENCE.
      Length = 72

10      Plus Strand HSPs:
      Score = 52 (18.3 bits), Expect = 3.3, Sum P(2) = 0.96
      Identities = 10/36 (27%), Positives = 18/36 (50%), Frame = +3

15      Q:   363 VVHGWWSTATITYWNSDRFTVLTALKIKLDTAAVTF 470
              +   WWS      Y++S RF ++  + +   T  +TF
      S:   16 IYFSWWSKFLTIYFSS-RFYIICVSVTNSTFKLTF 50
  
```

20 The segment of gill1181442 that is shown as "S" above is set out as sequence  
SEQ ID NO. 161.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 162 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

25 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares placenta Nb2HP and to a lesser extent in Morton Fetal Cochlea; Soares\_fetal\_lung\_NbHL19W; Soares\_multiple\_sclerosis\_2NbHMSP; Soares\_pregnant\_uterus\_NbHPU; Human Ovarian Cancer Reexcision; Early Stage Human Brain; Human B Cell Lymphoma; NCI\_CGAP\_GCB1; Soares\_testis\_NHT;

30 Human Cerebellum; Human skin Tumor, subtracted; Human Adult Lymph Node, subtracted; Early Stage Human Brain, random primed; Human Placenta; Human Thyroid; Human Tonsils, Lib 2; Stratagene colon (#937204); Stratagene ovarian cancer (#937219); HEL cell line; H Female Bladder, Adult; NTERA2 + retinoic acid, 14 days; Human Osteoclastoma, re-excision; Human Prostate; Human Uterine

35 Cancer; T-Cell PHA 24 hrs; Merkel Cells; NTERA2, control; Hepatocellular Tumor, re-excision; Soares\_multiple\_sclerosis\_2NbHMSP; Human Testes, Reexcision; Human Fetal Heart; Colon Normal III; Human Amygdala; Human Bone Marrow, treated; Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:44 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2888 of SEQ ID NO:44, b is an integer of 15 to 2902, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 35

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. pirlA60196|A60196 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " N-ras 5'-region protein NRU - human." A partial alignment demonstrating the observed homology is shown immediately below.

```
>pir|A60196|A60196 N-ras 5'-region protein NRU - human
      Length = 767

25      Plus Strand HSPs:

      Score = 2223 (782.5 bits), Expect = 8.5e-230, P = 8.5e-230
      Identities = 428/429 (99%), Positives = 428/429 (99%), Frame = +3

30      Q:      3 VRMFFHFSEILDGNQLHIADEVEFTVVPDMLSAQRNHAIRIKKLPKGTVSFHSKSDHRFL 182
      VRMFFHFSEILDGNQLHIADEVEFTVVPDMLSAQRNHAIRIKKLPKGTVSFHSKSDHRFL
      S:      339 VRMFFHFSEILDGNQLHIADEVEFTVVPDMLSAQRNHAIRIKKLPKGTVSFHSKSDHRFL 398

35      Q:      183 GTVEKEATFSNPKTTSPNKGTEKEAEDGIIAYDDCGVKLTIAFQAKDVEGSTSPQIGDKV 362
      GTVEKEATFSNPKTTSPNKG EKEAEDGIIAYDDCGVKLTIAFQAKDVEGSTSPQIGDKV
      S:      399 GTVEKEATFSNPKTTSPNKGTEKEAEDGIIAYDDCGVKLTIAFQAKDVEGSTSPQIGDKV 458

40      Q:      363 EFSISDKQRPQQVATCVRLGRNSNSKRLGYSVATLKDNGFGFIETANHDKEIFFHYSEF 542
      EFSISDKQRPQQVATCVRLGRNSNSKRLGYSVATLKDNGFGFIETANHDKEIFFHYSEF
      S:      459 EFSISDKQRPQQVATCVRLGRNSNSKRLGYSVATLKDNGFGFIETANHDKEIFFHYSEF 518

      Q:      543 SGDVSLELGDMVEYSLSKGKGNKVS AEKVNKTHSVNGITEEADPTIYSGKVIRPLRSVD 722
      SGDVSLELGDMVEYSLSKGKGNKVS AEKVNKTHSVNGITEEADPTIYSGKVIRPLRSVD
      S:      519 SGDVSLELGDMVEYSLSKGKGNKVS AEKVNKTHSVNGITEEADPTIYSGKVIRPLRSVD 578
```

5 Q: 723 PTQTEYQGMIEIVEEGDMKGEVYPFGIVGMANKGDCLQKGESVKFQLCVLGQNAQTMAYN 902  
 S: 579 PTQTEYQGMIEIVEEGDMKGEVYPFGIVGMANKGDCLQKGESVKFQLCVLGQNAQTMAYN 638  
 Q: 903 ITPLRRATVECVKDQFGFINYEVDGSKKLFHFVKEVDGIELQAGDEVEFSVILNQRTGK 1082  
 S: 639 ITPLRRATVECVKDQFGFINYEVDGSKKLFHFVKEVDGIELQAGDEVEFSVILNQRTGK 698  
 10 Q: 1083 CSACNVWRVCEGPKAVAAPRPDRLVNRLKNITLDDASAPRLMVLRLQPRGPDNSMGFGAER 1262  
 S: 699 CSACNVWRVCEGPKAVAAPRPDRLVNRLKNITLDDASAPRLMVLRLQPRGPDNSMGFGAER 758  
 15 Q: 1263 KIRQAGVID 1289  
 KIRQAGVID  
 S: 759 KIRQAGVID 767  
 Score = 284 (100.0 bits), Expect = 2.9e-21, P = 2.9e-21  
 Identities = 79/258 (30%), Positives = 130/258 (50%), Frame = +3  
 20 Q: 303 IAFQAKDVEGSTSPQIGDKVEFSISDKQRPQQVATCVRLGRNSNSKRLLGIVATLKDN 482  
 + + +DVEG+ + GDK+ F I + + G A + LL + R G V +K+  
 S: 108 LTYTPEDVEGNVQLETGDKINFVIDNNKHTGAVSARNIMLLKKKQ--ARCQGVVCAKMEA 165  
 25 Q: 483 FGFIETANHDKIEIFFHYSEFSGDSDSLELGDMEVYSLSKGKGNKVSAAEKVNKTHSVNGIT 662  
 FGFIE + KEIFFHYSEF GD+++L+ GD VE+++ G +V+ + V I  
 S: 166 FGFIERGDVVKEIFFHYSEFKGDLETLPQDDVEFTIKDRNGKEVATD-VRLLPQGTVIF 224  
 30 Q: 663 EEADPTIYSGKVIRPLRSVDPTQTEYQGMIEIVEEGDMKGEVYPFGIVGMANKGDCLQKG 842  
 E+ + G V + + V P++ + + ++ + + PFG +K L+ G  
 S: 225 EDISIEHFEGTVTKVIPKV-PSKNQNDPLPGRIKVDVFIKELPFGDKDTSKVTLLLE-G 282  
 35 Q: 843 ESVKFQLCVLGQNAQTMAYNITPLRRA-----TVECVKDQFGFINYEVDGSKKL 989  
 + V+F + ++ A NI L + ++D FGFI V ++  
 S: 283 DHVRFNISTDRDKLERATNIEVLSNTFQFTNEAREMGVIAAMRDGFGFIKC-VDRDVRM 341  
 Q: 990 FFHVKEVDGIELQAGDEVEFSVI 1061  
 FFH E+ DG +L DEVEF+V+  
 S: 342 FFHFSEILDGNQLHIADEVEFTVV 365  
 40

The segments of pirlA60196/A60196 that are shown as "S" above are set out as sequences SEQ ID NO. 163 and/or SEQ ID NO. 165. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 164 and/or SEQ ID NO. 166 which correspond to the Q sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen INFLS and to a lesser extent in



- Human Testes; Human 8 Week Whole Embryo; Soares placenta Nb2HP; Soares melanocyte 2NbHM; Stratagene Human skeletal muscle cDNA library, cat. #936215.; Primary Dendritic Cells, lib 1; Stratagene HeLa cell s3 937216; Human Testes Tumor; Smooth muscle, serum treated; Soares breast 3NbHBst; Human
- 5 Osteoclastoma; Monocyte activated; Soares infant brain 1NIB; Human Adipose Tissue, re-excision; Smooth muscle, serum induced, re-exc; Bone marrow; Human B Cell Lymphoma; Spleen, Chronic lymphocytic leukemia; Human Endometrial Tumor; Nine Week Old Early Stage Human; Human Skin Tumor; Synovial hypoxia; Human Adipose; Human umbilical vein endothelial cells, IL-4 induced;
- 10 Soares\_fetal\_heart\_NbHH19W; Weizmann Olfactory Epithelium; Smooth Muscle-HASTE normalized; NTERA2 teratocarcinoma cell line+retinoic acid (14 days); Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Soares\_testis\_NHT; Soares\_parathyroid\_tumor\_NbHPA; Human Osteoblasts II; Rejected Kidney, lib 4; Human T-Cell Lymphoma; Human Placenta; Soares\_parathyroid\_tumor\_NbHPA;
- 15 Human Placenta; Colon Normal III; Human Microvascular Endothelial Cells, fract. A; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Human Bone Marrow, treated; Bone Marrow Cell Line (RS4,11); Soares\_pregnant\_uterus\_NbHPU; Osteoblasts; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Synovial IL-1/TNF stimulated; Healing groin wound, 7.5 hours post incision; Gessler Wilms tumor; Human Ovarian Cancer
- 20 Reexcision; Stratagene muscle 937209; Ulcerative Colitis; Human Thymus Stromal Cells; Soares\_pregnant\_uterus\_NbHPU; Human Gall Bladder; Human Substantia Nigra; Adipocytes; Normal colon; Human Fetal Heart; Endothelial cells-control; NCI\_CGAP\_Br2; NCI\_CGAP\_Alv1; NCI\_CGAP\_Co10; NCI\_CGAP\_GCB1; Human Amygdala; Smooth muscle, control; NCI\_CGAP\_GC4; Dermatofibrosarcoma
- 25 Protuberance; prostate-edited; Human Aortic Endothelium; Human Adult Retina; Human Fetal Spleen; Aorta endothelial cells + TNF-a; Human Quadriceps; Stratagene ovarian cancer (#937219); Human Tonsils, Lib 2; HEL cell line; Human Amygdala, re-excision; pBMC stimulated w/ poly I/C; Human Umbilical Vein, Endo. remake; Salivary Gland, Lib 2; Stratagene hNT neuron (#937233); Stratagene muscle
- 30 937209; Stratagene neuroepithelium (#937231); Stratagene NT2 neuronal precursor 937230; wilm's tumor; Human Manic Depression Tissue; Human Neutrophil; Human Chronic Synovitis; Soares\_fetal\_heart\_NbHH19W;

Soares\_NSF\_F8\_9W\_OT\_PA\_P\_S1; Mo7e Cell Line GM-CSF treated (1ng/ml);  
 Apoptotic T-cell; Human Fetal Dura Mater; Human Uterine Cancer;  
 Soares\_NhHMPu\_S1; Stromal cell TF274; Human pancreatic islet; Stratagene hNT  
 neuron (#937233); Stratagene endothelial cell 937223; Spinal cord; Human fetal  
 5 heart, Lambda ZAP Express; Jia bone marrow stroma; Soares\_testis\_NHT;  
 Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Stratagene fetal retina 937202; Human Testes  
 Tumor, re-excision; NTERA2, control; Stratagene lung (#937210); Hepatocellular  
 Tumor, re-excision; Soares\_pregnant\_uterus\_NbHPU; Pancreas Islet Cell Tumor;  
 PC3 Prostate cell line; Brain frontal cortex; Stratagene NT2 neuronal precursor  
 10 937230; Soares\_fetal\_heart\_NbHH19W; Human Synovial Sarcoma; human tonsils; T  
 Cell helper I; Endothelial-induced; NCI\_CGAP\_Co3; NCI\_CGAP\_Ew1;  
 NCI\_CGAP\_GCB1; Anergic T-cell; NCI\_CGAP\_Ew1; NCI\_CGAP\_GC3;  
 NCI\_CGAP\_Pr22; Soares\_fetal\_liver\_spleen\_1NFLS\_S1;  
 Soares\_fetal\_lung\_NbHL19W; T cell helper II; Human placenta cDNA (TFujiwara);  
 15 NCI\_CGAP\_Lu5; NCI\_CGAP\_Alv1; NCI\_CGAP\_GCB1; NCI\_CGAP\_Kid3;  
 Stratagene NT2 neuronal precursor 937230; Human Prostate Cancer, Stage C;  
 Neuroblastoma; Prostate-BPH subtracted II; Human colon carcinoma (HCC) cell line;  
 LNCAP, differential expression; Human Platelets; Prostate, normal, subtraction I;  
 Human Fetal Brain, normalized AC5002; Kidney Cortex; Larynx normal #10 261-  
 20 273; Human Adult Skeletal Muscle; Messangial cell, frac 1; Normal trachea; Human  
 Normal Cartilage, Fraction I; brain stem; CD40 activated monocyte dendritic cells;  
 Human Normal Cartilage Fraction II; Human Prostate Cancer, Stage B2; Prostate;  
 Human B Cell 8866; Tongue Tumour; LNCAP + 0.3nM R1881; CHME Cell  
 Line, untreated; Human Fetal Lung; Human (Caco-2) cell line, adenocarcinoma,  
 25 colon, remake; LNCAP untreated; Morton Fetal Cochlea; Human Tonsils, Fraction 2;  
 Human Primary Breast Cancer, re-excision; Human Cardiomyopathy, subtracted;  
 Brain-medulloblastoma; Healing Abdomen wound, 70&90 min post incision; Human  
 Neutrophils, Activated, re-excision; Frontal lobe, dementia, re-excision; Smooth  
 muscle, control, re-excision; Activated T-cells; H. Epididymus, cauda; Human T-cell  
 30 lymphoma, re-excision; Soares retina N2b4HR; Human Whole Brain, re-excision;  
 Human Epididymus; Pancreas Tumor PCA4 Tu; Human Synovium; Hepatocellular  
 Tumor; Human Prostate Cancer, Stage C fraction; human corpus colosum; H Female

- Bladder, Adult; NTERA2 + retinoic acid, 14 days; Synovial hypoxia-RSF subtracted; H. Kidney Cortex, subtracted; Ovarian Tumor 10-3-95; Human Stomach.re-excision; Stratagene ovarian cancer (#937219); Human Osteoclastoma, re-excision; H. Ovarian Tumor, II, OV5232; Jurkat T-cell G1 phase; Myeloid Progenitor Cell Line; Jurkat T-Cell, S phase; Human Ovary; Spleen metastatic melanoma; H. Lymph node breast Cancer; Brain Frontal Cortex, re-excision; Human Adult Small Intestine; KMH2; Human Infant Brain; Human Prostate; Human Umbilical Vein, Reexcision; T-Cell PHA 16 hrs; Barstead spleen HPLRB2; Heart; Human colon mucosa; Human pancreatic islet; Human adult lung 3' directed MboI cDNA; Jia bone marrow stroma;
- 10 Soares\_pregnant\_uterus\_NbHPU; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Soares\_senescent\_fibroblasts\_NbHSF; 12 Week Old Early Stage Human, II; Human Umbilical Vein Endothelial Cells, uninduced; Human Pancreas Tumor; Soares\_testis\_NHT; Human Heart; Macrophage-oxLDL; Stratagene HeLa cell s3 937216; Stratagene fibroblast (#937212); Stratagene neuroepithelium NT2RAMI
- 15 937234; Liver, Hepatoma; Soares\_fetal\_lung\_NbHL19W; Human Rhabdomyosarcoma; Synovial Fibroblasts (control); Epithelial-TNF $\alpha$  and INF induced; Gessler Wilms tumor; Human pancreatic islet; Soares\_NhHMPu\_S1; Stratagene hNT neuron (#937233); Stratagene lung carcinoma 937218; Hemangiopericytoma; Soares breast 2NbHBst; Macrophage (GM-CSF treated); H.
- 20 Frontal cortex,epileptic,re-excision; Soares\_NhHMPu\_S1; Human Liver, normal; Human placenta cDNA (TFujiwara); Macrophage-oxLDL, re-excision; Colon Tumor; 12 Week Old Early Stage Human; Colon Carcinoma; Dendritic cells, pooled; Colon Tumor II; Soares\_testis\_NHT; Stratagene HeLa cell s3 937216; Stratagene hNT neuron (#937233); Soares\_fetal\_lung\_NbHL19W;
- 25 Soares\_placenta\_8to9weeks\_2NbHP8to9W; Primary Dendritic cells,frac 2; Human Adult Pulmonary,re-excision; NCI\_CGAP\_AA1; NCI\_CGAP\_Co9; NCI\_CGAP\_GC1; NCI\_CGAP\_GCB0; CD34 positive cells (Cord Blood); Activated T-cell(12h)/Thiouridine-re-excision; NCI\_CGAP\_Br7; NCI\_CGAP\_GC3; NCI\_CGAP\_Ov2; NCI\_CGAP\_Kid6; Soares\_placenta\_8to9weeks\_2NbHP8to9W;
- 30 Stratagene neuroepithelium (#937231); Human Cerebellum; Soares\_fetal\_lung\_NbHL19W; Soares\_multiple\_sclerosis\_2NbHMSP; Soares\_fetal\_liver\_spleen\_1NFLS\_S1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1446 of SEQ ID NO:45, b is an integer of 15 to 1460, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 36

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil186804 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " L6 [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi|186804 L6 [Homo sapiens] >pir|A42926|A42926 L6 surface protein - human
Length = 202
```

Plus Strand HSPs:

```
Score = 841 (296.0 bits), Expect = 2.2e-83, P = 2.2e-83
Identities = 159/202 (78%), Positives = 162/202 (80%), Frame = +2
```

```

30  Q:   71 MCYGKCARCIGHSLVGXXXXXXXXXXXXYFPNGETKYASENHLSRFVWFFSGIVGGGLLM 250
      MCYGKCARCIGHSLVG                      YFPNGETKYASENHLSRFVWFFSGIVGGGLLM
      S:   1 MCYGKCARCIGHSLVGLALLCIAANILLYFPNGETKYASENHLSRFVWFFSGIVGGGLLM 60

35  Q:  251 LLPAFVFIGLEQDDCCGCCGHENCGKRCAMLSSVLAALIGLQDLATVSLWQPLA*QKDHY 430
      LLPAFVFIGLEQDDCCGCCGHENCGKRCAMLSSVLAALIG+          + L +
      S:   61 LLPAFVFIGLEQDDCCGCCGHENCGKRCAMLSSVLAALIGIAGSGYCVIVAALGLAEGPL 120

40  Q:  431 VLIPSASGTTPLPSTEGQYLLDTSTWSECTEPKHIVEWNVSLFSILLALGGIEFILCLIQ 610
      L                      STEGQYLLDTSTWSECTEPKHIVEWNVSLFSILLALGGIEFILCLIQ
      S:  121 CLDSLQWNYTFASTEGQYLLDTSTWSECTEPKHIVEWNVSLFSILLALGGIEFILCLIQ 180

      Q:  611 VINGVLGGICGFCCSHQQQYDC 676
      VINGVLGGICGFCCSHQQQYDC
      S:  181 VINGVLGGICGFCCSHQQQYDC 202
```

The segment of gil186804 that is shown as "S" above is set out as sequence SEQ ID NO. 167. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 168 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in Endothelial cells-control.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 95 as residues: Thr-49 to Pro-56.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:46 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 882 of SEQ ID NO:46, b is an integer of 15 to 896, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 37**

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil2982366 (all information available through the recited accession number is

incorporated herein by reference) which is described therein as "10kD protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >gi|2982366 (AF053470) 10kD protein [Homo sapiens]
          Length = 87

          Plus Strand HSPs:
10      Score = 493 (173.5 bits), Expect = 1.8e-46, P = 1.8e-46
          Identities = 87/87 (100%), Positives = 87/87 (100%), Frame = +1

          Q:   289 MYCLQWLLPVLLIPKPLNPALWFSHSMFMGFYLLSFLLERKPCTICALVFLAALFLICYS 468
                  MYCLQWLLPVLLIPKPLNPALWFSHSMFMGFYLLSFLLERKPCTICALVFLAALFLICYS
15      S:     1 MYCLQWLLPVLLIPKPLNPALWFSHSMFMGFYLLSFLLERKPCTICALVFLAALFLICYS 60

          Q:   469 CWGNCFlyHCSDSLPEsAHDPGVVGT 549
                  CWGNCFlyHCSDSLPEsAHDPGVVGT
          S:    61 CWGNCFlyHCSDSLPEsAHDPGVVGT 87
20

```

The segment of gi|2982366 that is shown as "S" above is set out as sequence SEQ ID NO. 169. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 170 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Nine Week Old Early Stage Human and to a lesser extent in Soares retina N2b4HR; Soares\_pregnant\_uterus\_NbHPU; CD34 positive cells (Cord Blood); Human Eosinophils; Human Cerebellum; 12 Week Old Early Stage Human; NCI\_CGAP\_GCB1; Human Amygdala; Stratagene colon (#937204); Human 8 Week Whole Embryo; Primary Dendritic Cells, lib 1; Human Heart; Human Fetal Brain; Soares\_testis\_NHT; Soares\_fetal\_heart\_NbHH19W; Soares\_total\_fetus\_Nb2HF8\_9w; NCI\_CGAP\_Lu5; normalized infant brain cDNA; Human Testes Tumor; Human Fetal Heart; Human Neutrophil, Activated; Activated T-Cell (12hs)/Thiouridine labelledEco; Human Soleus; Stratagene placenta

- (#937225); Soares\_fetal\_lung\_NbHL19W; Soares\_fetal\_heart\_NbHH19W; Ovarian Tumor 10-3-95; Human Whole Brain #2 - Oligo dT > 1.5Kb; Human Adipose Tissue, re-excision; Human Infant Brain; Human Fetal Dura Mater; Human Hippocampus; Human Chondrosarcoma; Hemangiopericytoma; Hepatocellular Tumor, re-excision;
- 5 Stratagene liver (#937224); Soares\_NFL\_T\_GBC\_S1; Stratagene hNT neuron (#937233); HM1; NCI\_CGAP\_Co9; 1-NIB; Stratagene Human skeletal muscle cDNA library, cat. #936215.; 12 Week Early Stage Human II, Reexcision; Early Stage Human Brain; Soares melanocyte 2NbHM; Primary Dendritic cells,frac 2; Human Adult Pulmonary,re-excision; Human Osteoclastoma; Human Microvascular
- 10 Endothelial Cells, fract. A; Soares\_pregnant\_uterus\_NbHPU; Stratagene ovarian cancer (#937219); Keratinocyte; Soares fetal liver spleen 1NFLS; Soares infant brain 1NIB; Human Kidney, normal Adult; Human Pancreatic Langerhans; Human Gastrocnemius; Testis, normal; brain stem; Salivary Gland, Lib 3; LNCAP + 30nM R1881; Frontal Lobe, Dementia; Human Adult Pulmonary; H. Atrophic
- 15 Endometrium; Human Stomach; Human Fetal Bone; Human Thyroid; Early Stage Human Lung, subtracted; HSA 172 Cells; Soares adult brain N2b4HB55Y; Smooth muscle, IL1b induced; Soares\_senescent\_fibroblasts\_NbHSF; Alzheimers, spongy change; Glioblastoma; Human Stomach,re-excision; Human Frontal Cortex, Schizophrenia; Stratagene muscle 937209; Human Osteosarcoma; Jurkat T-cell G1
- 20 phase; Human Manic Depression Tissue; H. Lymph node breast Cancer; Human Fetal Kidney; Human Primary Breast Cancer Reexcision; Human Pancreas Tumor; Human umbilical vein endothelial cells, IL-4 induced; Soares\_fetal\_heart\_NbHH19W; Soares\_fetal\_liver\_spleen\_1NFLS\_S1; Human Whole Six Week Old Embryo; H. Frontal cortex,epileptic,re-excision; Stratagene lung (#937210); Smooth muscle,
- 25 serum induced,re-exc; Pancreas Islet Cell Tumor; Macrophage-oxLDL, re-excision; Human Gall Bladder; PC3 Prostate cell line; HM3; Soares\_senescent\_fibroblasts\_NbHSF; Stratagene fetal retina 937202; Fetal Heart; Human adult (K.Okubo); Human heart cDNA (YNakamura); Human pancreatic cancer cell line Patu 8988t; Human retina cDNA randomly primed sublibrary;
- 30 NCI\_CGAP\_AA1; NCI\_CGAP\_AR1; NCI\_CGAP\_Br2; NCI\_CGAP\_Co2; NCI\_CGAP\_HN3; NCI\_CGAP\_HN4; NCI\_CGAP\_Ov8; NCI\_CGAP\_Kid3; NCI\_CGAP\_Lar1; NCI\_CGAP\_Pr22; NCI\_CGAP\_Pr23; NCI\_CGAP\_Brn23;

Soares\_fetal\_lung\_NbHL19W; Soares breast 3NbHBst; Human colon mucosa;  
 Soares\_fetal\_heart\_NbHH19W; H Macrophage (GM-CSF treated), re-excision;  
 Soares\_pineal\_gland\_N3HPG; Human Placenta; Bone marrow; human tonsils;  
 Endothelial cells-control; Soares\_fetal\_heart\_NbHH19W; Human B Cell Lymphoma;  
 5 Spleen, Chronic lymphocytic leukemia; Neutrophils IL-1 and LPS induced;  
 Hodgkin's Lymphoma II; Osteoblasts; Stratagene HeLa cell s3 937216; Stratagene  
 neuroepithelium NT2RAMI 937234; Stratagene colon (#937204).

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.  
 96 as residues: Ser-11 to Ser-17.

10 Many polynucleotide sequences, such as EST sequences, are publicly  
 available and accessible through sequence databases. Some of these sequences are  
 related to SEQ ID NO:47 and may have been publicly available prior to conception of  
 the present invention. Preferably, such related polynucleotides are specifically  
 excluded from the scope of the present invention. To list every related sequence  
 15 would be cumbersome. Accordingly, preferably excluded from the present invention  
 are one or more polynucleotides comprising a nucleotide sequence described by the  
 general formula of a-b, where a is any integer between 1 to 2056 of SEQ ID NO:47, b  
 is an integer of 15 to 2070, where both a and b correspond to the positions of  
 nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a  
 20 + 14.

### FEATURES OF PROTEIN ENCODED BY GENE NO: 38

The computer algorithm BLASTX has been used to determine that the  
 translation product of this gene shares sequence homology with, as a non-limiting  
 25 example, the sequence accessible through the following database accession no.  
 gil829423 (all information available through the recited accession number is  
 incorporated herein by reference) which is described therein as "env gene product  
 [Human immunodeficiency virus type 1]." A partial alignment demonstrating the  
 observed homology is shown immediately below.

30

```
>gi|829423 env gene product [Human immunodeficiency virus type 1]
>sp|Q71908|Q71908 CNTRL 1, CLONE 2, ACCESSORY REGION GENES,
COMPLETE
CDS (FRAGMENT).
```



Length = 40

Plus Strand HSPs:

5      Score = 64 (22.5 bits), Expect = 1.6, P = 0.80  
       Identities = 10/35 (28%), Positives = 18/35 (51%), Frame = +3

Q:      48 LKSHKKNYNMLWIFKWQVL\*LFFVCKILSNHWENI 152  
           +K +KNY LW +    +L +    +C + N W +  
 10      S:      3 VKGTRKNYQRLWRWGTMLLGMLMICSVAGNLWVTV 37

The segment of gil829423 that is shown as "S" above is set out as sequence SEQ ID NO. 171.

Preferred polypeptides of the invention comprise a polypeptide having the  
 15 amino acid sequence set out as SEQ ID NO. 172 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_pregnant\_uterus\_NbHPU and to a lesser extent in  
 20 Morton Fetal Cochlea; Stratagene fibroblast (#937212); Human Normal Breast; Human Osteoclastoma, re-excision; Colon Tumor; Endothelial cells-control.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of  
 25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 555 of SEQ ID NO:48, b  
 30 is an integer of 15 to 569, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 39

35      It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Endothelial-induced; Endothelial cells-control and to a lesser extent in Human Brain, Striatum; H. Frontal cortex,epileptic,re-excision;

Macrophage-oxLDL, re-excision; Human Amygdala; Human endometrial stromal cells-treated with progesterone; Human Whole Six Week Old Embryo; Human Fetal Lung III; Human Placenta; Human Old Ovary, subtracted; Skin, burned; Pharynx Carcinoma; H. Frontal Cortex, Epileptic; Striatum Depression; Human Fetal Lung;

5 Human (Caco-2) cell line, adenocarcinoma, colon, remake; LNCAP untreated; Human Umbilical Vein Endothelial Cells, fract. A; H. Epididymus, caput & corpus; H. Epididymus, cauda; H. Whole Brain #2, re-excision; Soares\_total\_fetus\_Nb2HF8\_9w; B Cell lymphoma; Stratagene endothelial cell 937223; Human endometrial stromal cells-treated with estradiol; Stratagene hNT

10 neuron (#937233); Breast Cancer cell line, MDA 36; Alzheimers, spongy change; Stratagene endothelial cell 937223; Human Frontal Cortex, Schizophrenia; Human Adipose Tissue, re-excision; Human endometrial stromal cells; Stratagene endothelial cell 937223; Spinal Cord, re-excision; Fetal Liver, subtraction II; Human Bone Marrow, re-excision; Human Hypothalamus, Schizophrenia; Human Adipose; Human

15 Adrenal Gland Tumor; Pancreas Islet Cell Tumor; Colon Carcinoma; Human Substantia Nigra; Colon Normal II; Soares\_multiple\_sclerosis\_2NbHMSP; Bone marrow; Human Fetal Heart; human tonsils; Colon Normal III; Human Microvascular Endothelial Cells, fract. A; Monocyte activated; T cell helper II; Human 8 Week Whole Embryo; Soares\_fetal\_lung\_NbHL19W; Soares fetal liver spleen 1NFLS.

20 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 98 as residues: Thr-45 to Asn-62, Thr-108 to Phe-117, Asp-124 to Arg-135, Thr-187 to Gly-193.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

25 related to SEQ ID NO:49 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the

30 general formula of a-b, where a is any integer between 1 to 1241 of SEQ ID NO:49, b is an integer of 15 to 1255, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 40

5 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_parathyroid\_tumor\_NbHPA and to a lesser extent in Soares\_pregnant\_uterus\_NbHPU; Human Osteoclastoma, re-excision; Human Osteoclastoma; Soares\_total\_fetus\_Nb2HF8\_9w; Osteoclastoma-normalized A; Soares\_fetal\_heart\_NbHH19W; Soares\_multiple\_sclerosis\_2NbHMSP; Human  
10 umbilical vein endothelial cells, IL-4 induced; Soares melanocyte 2NbHM; Soares\_multiple\_sclerosis\_2NbHMSP; Human aorta polyA+ (TFujiwara); Soares infant brain 1NIB; Healing Abdomen Wound, 15 days post incision; Osteoclastoma-normalized B; Ea.hy.926 cell line; Human Prostate Cancer, Stage B2; HEL cell line; Human Osteoclastoma Stromal Cells - unamplified; Healing groin wound, 7.5 hours  
15 post incision; Myeloid Progenitor Cell Line; Human Chronic Synovitis; Stromal cell TF274; Soares\_pregnant\_uterus\_NbHPU; PERM TF274; NCI\_CGAP\_Pr22; Human Thymus Stromal Cells; Pancreas Islet Cell Tumor; Endothelial cells-control; Human adult (K.Okubo); Nine Week Old Early Stage Human.

Many polynucleotide sequences, such as EST sequences, are publicly  
20 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention  
25 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2340 of SEQ ID NO:50, b is an integer of 15 to 2354, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

30

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 41

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil841318 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "mutant sterol regulatory element binding protein-2 [Cricetulus griseus]." A partial alignment demonstrating the observed homology is shown immediately below.

```

10  >gi|841318 mutant sterol regulatory element binding protein-2 [Cricetulus
      griseus]
      Length = 839

      Minus Strand HSPs:

15  Score = 973 (342.5 bits), Expect = 2.8e-97, P = 2.8e-97
      Identities = 195/299 (65%), Positives = 225/299 (75%), Frame = -2

      Q:  2236  EGFGRPHKDLWASKNENEEILERPAQLANARETPHSPGVEDAPIAKVGVLAASMEAKAS 2057
      S:  534    EGFGRPHKDLWA+K+ENEE ER Q + E PH+ GVEDAPI+KVGVLAA+MEAKAS
20      S:  534    EGFGRPHKDLWANKSENEETSERSPQSPDVGEIPHNSGVEDAPISKVGVLAANMEAKAS 593

      Q:  2056  SQQEKEDKPAETKKLRIAWPPPTLGGSSGSALEEGIKMSKPKWPPPEDEISKPEVPEDVXX 1877
      S:  594    Q+EKEDKPAETKKLRIAWPPP ELGSSGS +EEG K SKPKWPPPEDE SK E DV
25      S:  594    PQREKEDKPAETKKLRIAWPPPAELGSSGSTMEEGFKGSKPKWPPPEDETSKTEALGDVDL 653

      Q:  1876  XXXXXXXXXXXXXXXX PFTVAASFQSTSVKSPKTVSPPIRKGWSMSEQSEESVGGRVAER 1697
      S:  654    DLKKLRRSSSLKERSRPFTVAASFRTSSLKSPKTLSP-LRKGWSMSEQNEESGGGIVTAG 712

30      Q:  1696  KQVENAKASKKNGNVGKTTWQNKES-KGETGKRSKEGHSLEMENENLVENGADSDDEDDNS 1520
      S:  713    KQ ENAKA K NV K+ WQ ++S + E +RSK+ H +E+E+EN +ENGA+ E D+
      S:  713    KQ TENAKAPGKKENVRKSRWQREDSAREEVARRSKDIHGVELESENFIENGANIYEGDSC 772

35      Q:  1519  FLKQQSPQEPKSLNWSSFVDNTFAEEFTTQNQKSQDVELWEGEVVKELSVEEQIKRNRYY 1340
      S:  773    L QQSP +PKS +WS FVD T EEFTTQNQKS DV EGEVV+ELSVEEQIKRNRYY
      S:  773    DLVQQSPLDPKSPSWSGFVDTTPTTEFTTQNQKSHDVGFRGEVVRELSVEEQIKRNRYY 832

```

The segment of gil841318 that is shown as "S" above is set out as sequence SEQ ID NO. 173. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 174 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares placenta Nb2HP and to a lesser extent in Soares melanocyte 2NbHM; Soares\_pregnant\_uterus\_NbHPU; 12 Week Old Early Stage Human, II; Keratinocyte; Nine Week Old Early Stage Human; Stratagene colon (#937204); Stratagene ovarian cancer (#937219); Messangial cell, frac 2; Soares\_fetal\_lung\_NbHL19W; Human Osteoblasts II; Human Chondrosarcoma; Synovial Fibroblasts (control); Soares\_placenta\_8to9weeks\_2NbHP8to9W; Human Adrenal Gland Tumor; Stratagene lung (#937210); 12 Week Old Early Stage Human; Colon Normal III; Soares\_pregnant\_uterus\_NbHPU; H. Kidney Medulla, subtracted; Normal trachea; Human Normal Cartilage Fraction IV; Nasal polyps; Human Prostate, subtracted; Saos2 Cells, Vitamin D3 Treated; Human Adult Spleen; Human Adult Pulmonary; human colon cancer; B Cell lymphoma; Human Synovium; Amniotic Cells - Primary Culture; Synovial hypoxia-RSF subtracted; Glioblastoma; Ovarian Tumor 10-3-95; Human Adult Small Intestine; Soares\_parathyroid\_tumor\_NbHPA; Soares\_NhHMPu\_S1; Soares\_total\_fetus\_Nb2HF8\_9w; Human Fetal Dura Mater; Human Pancreas Tumor; Stromal cell TF274; Hemangiopericytoma; Human Placenta; Soares breast 3NbHBst; Human Testes, Reexcision; Human Fetal Heart; Human Adult Pulmonary, re-excision; Activated T-Cell (12hs)/Thiouridine labelledEco; Endothelial cells-control; Human Amygdala; Human Bone Marrow, treated; Human Testes; Stratagene colon (#937204); T cell helper II; Osteoblasts; Stratagene colon (#937204); Human Cerebellum; Primary Dendritic Cells, lib 1.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 100 as residues: Trp-14 to Ser-33.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:51 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2253 of SEQ ID NO:51, b

is an integer of 15 to 2267, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 42

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares melanocyte 2NbHM;

Soares\_pregnant\_uterus\_NbHPU; Soares fetal liver spleen 1NFLS and to a lesser extent in Stratagene lung (#937210); breast lymph node CDNA library;

- 10 Soares\_pregnant\_uterus\_NbHPU; Human Placenta; Human Endometrial Tumor; Keratinocyte; Soares placenta Nb2HP; prostate-edited; Human Lung; Human Colon, re-excision; Human Prostate; NCI\_CGAP\_GCB1; Human Placenta; Soares breast 3NbHBst; Soares\_fetal\_heart\_NbHH19W; Endothelial cells-control; Activated T-cell(12h)/Thiouridine-re-excision; Stratagene muscle 937209; HM1; Human adult
- 15 (K.Okubo); Human retina cDNA randomly primed sublibrary; NCI\_CGAP\_Co3; NCI\_CGAP\_GC4; NCI\_CGAP\_GCB1; NCI\_CGAP\_Kid3; Nine Week Old Early Stage Human; Soares infant brain 1NIB.

- Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
- 20 related to SEQ ID NO:52 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
- 25 general formula of a-b, where a is any integer between 1 to 2284 of SEQ ID NO:52, b is an integer of 15 to 2298, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

## 30 FEATURES OF PROTEIN ENCODED BY GENE NO: 43

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting

example, the sequence accessible through the following database accession no. gil862343 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " Gcap1 gene product [Mus musculus]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi|862343 Gcap1 gene product [Mus musculus]
      Length = 85
```

Minus Strand HSPs:

```
Score = 85 (29.9 bits), Expect = 0.019, P = 0.018
Identities = 20/48 (41%), Positives = 30/48 (62%), Frame = -3
```

```
15  Q:   396 YICVYLYVCTHPCIIYILSVQT-LHMYTHYLCKI*YMYNAYKF*QVYLC 253
      YIC+ +Y+ + I IYLS+ T +H +TH I Y+Y Y + +YLC
      S:   22 YICISYILSYLSISYILSIYTYIHTHTHTTYI-YIY-IYIYIYLYLC 68
```

The segment of gil862343 that is shown as "S" above is set out as sequence SEQ ID NO. 175. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 176 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares\_fetal\_lung\_NbHL19W; Endothelial cells-control and to a lesser extent in Human Astrocyte; Morton Fetal Cochlea; Frontal lobe,dementia,re-excision; Human Brain, Striatum; Bone Marrow Stromal Cell, untreated; Human Amygdala; Human Microvascular Endothelial Cells, fract. A.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence

would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1095 of SEQ ID NO:53, b is an integer of 15 to 1109, where both a and b correspond to the positions of  
 5 nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 44

The computer algorithm BLASTX has been used to determine that the  
 10 translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil535097 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " anion exchanger [Homo sapiens]." A partial alignment demonstrating the observed homology is shown  
 15 immediately below.

```

    >gi|535097 anion exchanger [Homo sapiens] >sp|Q15292|Q15292 ANION
    EXCHANGER.
          Length = 36
20      Plus Strand HSPs:
          Score = 69 (24.3 bits), Expect = 1.7, P = 0.82
          Identities = 13/20 (65%), Positives = 14/20 (70%), Frame = +2
25      Q:  1172 WLGVMACTCNSSTSGGQGGR 1231
           WLG +A  CN ST GGQ GR
           S:  15  WLGAVAHVCNPSTLGGQVGR 34
  
```

30 The segment of gil535097 that is shown as "S" above is set out as sequence SEQ ID NO. 177. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere  
 35 herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 178 which corresponds to the Q



sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: human tonsils; Endothelial cells-control.

5 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 103 as residues: Met-1 to Lys-6.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:54 and may have been publicly available prior to conception of  
10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1282 of SEQ ID NO:54, b  
15 is an integer of 15 to 1296, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 45**

20 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: CD34 positive cells (Cord Blood) and to a lesser extent in Spleen metastatic melanoma; Endothelial cells-control.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are  
25 related to SEQ ID NO:55 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the  
30 general formula of a-b, where a is any integer between 1 to 1418 of SEQ ID NO:55, b is an integer of 15 to 1432, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 46

5 It has been discovered that this gene is expressed primarily in Endothelial cells-control.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 965 of SEQ ID NO:56, b is an integer of 15 to 979, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

#### FEATURES OF PROTEIN ENCODED BY GENE NO: 47

20 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil2773183 (all information available through the recited accession number is incorporated herein by reference) which is described therein as " a C. elegans protein." A partial alignment demonstrating the observed homology is shown immediately below.

30 >gi|2773183 (AF039720) No definition line found [Caenorhabditis elegans]  
>gi|2773182 (AF039720) No definition line found  
[Caenorhabditis  
elegans] (SUB 657-885)  
Length = 885

35 Minus Strand HSPs:

Score = 322 (113.3 bits), Expect = 6.5e-42, Sum P(4) = 6.5e-42  
Identities = 54/92 (58%), Positives = 67/92 (72%), Frame = -2

Q: 2504 EVIYXXXXXXXXXERAHHC SICKRCIRKMDHHC PWVNNCVGEKNQRFVLFTMYIALSS 2325  
 + IY +RAHHC S+C+RCIR+MDHHC PWVNNCVGE NQ+FFVLFTMYIAL S  
 S: 81 QTIYKCQKCASIKPDRAHHC SVCERCIRMDHHC PWVNNCVGEGNQKFFVLFTMYIALLS 140  
 5 Q: 2324 VHALILCGFQFISCVRGQWTECSDFSPITVI 2229  
 +HAL +QF+ CV +W +CS +PP T +  
 S: 141 MHALYWG VQFVLCVGA EWQQCSSLTPPATTL 172  
 10 Score = 143 (50.3 bits), Expect = 6.5e-42, Sum P(4) = 6.5e-42  
 Identities = 33/72 (45%), Positives = 43/72 (59%), Frame = -3  
 Q: 2194 FSLSLAVMFGTQIHSICNDETEIERLKSEKPTWER--RLR---WEGMKS VFGGPPSXLWM 2030  
 F++ +VMFGTQI SICNDET IE ++S + R R W+ ++ VFGGP S W  
 15 S: 184 FAIFTSVMFGTQISSICNDETTIESMRSRNANLDEDESRNNAWKNLQLVFGGPPFSVRWF 243  
 Q: 2029 NPFVGFRRRLPT 1991  
 NP + F PT  
 S: 244 NP-LAMPFVARPT 255

20 The segments of gil2773183 that are shown as "S" above are set out as  
 sequences SEQ ID NO. 179 and/or SEQ ID NO. 181.

Preferred polypeptides of the invention comprise a polypeptide having the  
 amino acid sequence set out as SEQ ID NO. 180 and/or SEQ ID NO. 182 which  
 correspond to the Q sequences in the alignment shown above (gaps introduced in a  
 25 sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following  
 tissues/cDNA libraries: Primary Dendritic Cells, lib 1; Soares infant brain 1NIB and  
 to a lesser extent in Human Tonsils, Lib 2; Endothelial-induced;  
 Soares\_fetal\_heart\_NbHH19W; Stratagene placenta (#937225); Jurkat T-cell G1  
 30 phase; T-Cell PHA 16 hrs; Human Bone Marrow, re-excision; L428; Stromal cell  
 TF274; Human Gall Bladder; Soares melanocyte 2NbHM; Human Microvascular  
 Endothelial Cells, fract. A; Activated T-cell(12h)/Thiouridine-re-excision; T cell  
 helper II; Soares\_total\_fetus\_Nb2HF8\_9w.

Many polynucleotide sequences, such as EST sequences, are publicly  
 35 available and accessible through sequence databases. Some of these sequences are  
 related to SEQ ID NO:57 and may have been publicly available prior to conception of  
 the present invention. Preferably, such related polynucleotides are specifically  
 excluded from the scope of the present invention. To list every related sequence  
 would be cumbersome. Accordingly, preferably excluded from the present invention  
 40 are one or more polynucleotides comprising a nucleotide sequence described by the  
 general formula of a-b, where a is any integer between 1 to 2653 of SEQ ID NO:57, b

is an integer of 15 to 2667, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

## 5 FEATURES OF PROTEIN ENCODED BY GENE NO: 48

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|d1033002 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "D4 dopamine receptor (D4DR) [Lemur catta]." A partial alignment demonstrating the observed homology is shown immediately below.

```

15  >gnl|PID|d1033002 (AB016202) D4 dopamine receptor (D4DR) [Lemur catta]
    >sp|O77493|O77493 D4 DOPAMINE RECEPTOR (D4DR) (FRAGMENT).
    Length = 32

    Minus Strand HSPs:

20  Score = 42 (14.8 bits), Expect = 5.3, Sum P(2) = 0.99
    Identities = 8/20 (40%), Positives = 13/20 (65%), Frame = -2

    Q:   689 PSIGVDPKSGSCGKEIQSQSP 630
        P+ G+ P+G CG + Q +P
25  S:   1 PAPGL-PQGPGADCQPPAP 19

```

The segment of gnl|PID|d1033002 that is shown as "S" above is set out as sequence SEQ ID NO. 183. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out as SEQ ID NO. 184 which corresponds to the Q sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in Endothelial-induced.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:58 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1342 of SEQ ID NO:58, b is an integer of 15 to 1356, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to a + 14.

#### **FEATURES OF PROTEIN ENCODED BY GENE NO: 49**

It has been discovered that this gene is expressed primarily in Endothelial-induced.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 108 as residues: Met-26 to Glu-34.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:59 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 836 of SEQ ID NO:59, b is an integer of 15 to 850, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:59, and where b is greater than or equal to a + 14.

Table 1

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HE9NB82	203960 4/26/99	Uni-ZAP XR	11	1329	10	1329	60	60	60	1	26	27	109
2	HE9RN58	203960 4/26/99	Uni-ZAP XR	12	834	1	834	515	515	61	1	23	24	26
3	HE9TA42	203960 4/26/99	Uni-ZAP XR	13	3371	2187	3359	2540	2540	62	1	28	29	36
4	HEAAC21	203960 4/26/99	Uni-ZAP XR	14	1094	1	1094	68	68	63	1	15	16	63
5	HEAAC39	203960 4/26/99	Uni-ZAP XR	15	1243	1	1243		299	64	1			5
6	HEAAC48	203960 4/26/99	Uni-ZAP XR	16	818	1	818	178	178	65	1			30
7	HEAAD63	203960 4/26/99	Uni-ZAP XR	17	847	1	847	107	107	66	1			10
8	HEAAE19	203960 4/26/99	Uni-ZAP XR	18	662	1	662	296	296	67	1			14
9	HEAAM54	203960 4/26/99	Uni-ZAP XR	19	2254	1	2254		219	68	1	18	19	41
10	HEAAM96	203960 4/26/99	Uni-ZAP XR	20	1079	1	1079		217	69	1			15

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
11	HEAAU28	203960 4/26/99	Uni-ZAP XR	21	632	1	632	73	73	70	1	35	36	40
12	HEBAP51	203960 4/26/99	Uni-ZAP XR	22	1492	868	1492		1060	71	1	36	37	37
13	HEBAT05	203960 4/26/99	Uni-ZAP XR	23	638	1	638	216	216	72	1	25	26	55
14	HEBBF78	203960 4/26/99	Uni-ZAP XR	24	944	298	933		590	73	1			19
15	HEBBK04	203960 4/26/99	Uni-ZAP XR	25	408	1	408		280	74	1	16	17	29
16	HEBCN80	203960 4/26/99	Uni-ZAP XR	26	1687	1	1687		322	75	1	15	16	50
17	HEBCY54	203960 4/26/99	Uni-ZAP XR	27	1189	1	1189	172	172	76	1	25	26	118
18	HEBDF77	203960 4/26/99	Uni-ZAP XR	28	1820	1	1820	681	681	77	1	30	31	36
19	HEBDQ91	203960 4/26/99	Uni-ZAP XR	29	1573	1007	1573		1211	78	1	29	30	41
20	HEBDW31	203960 4/26/99	Uni-ZAP XR	30	1328	486	1305	734	734	79	1	22	23	142
21	HEBFL36	203960 4/26/99	Uni-ZAP XR	31	590	1	590	381	381	80	1	25	26	43

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
22	HEBGC01	203960 4/26/99	Uni-ZAP XR	32	752	1	752	201	201	81	1	22	23	36
23	HEBGE07	203960 4/26/99	Uni-ZAP XR	33	1867	1	1867	106	106	82	1	25	26	42
24	HEBGE23	203960 4/26/99	Uni-ZAP XR	34	419	1	419	153	153	83	1	31	32	81
25	HEBGE85	203960 4/26/99	Uni-ZAP XR	35	949	380	949	615	615	84	1			7
26	HEEAB58	203960 4/26/99	Uni-ZAP XR	36	1635	942	1607	1157	1157	85	1			17
27	HEEAJ46	203960 4/26/99	Uni-ZAP XR	37	2414	1	2414	291	291	86	1			7
28	HEGAI20	203960 4/26/99	Uni-ZAP XR	38	929	1	929	111	111	87	1	34	35	57
29	HEGAU15	203960 4/26/99	Uni-ZAP XR	39	1125	1	1125	59	59	88	1	30	31	34
30	HEBCI18	203960 4/26/99	Uni-ZAP XR	40	1121	713	1050	855	855	89	1	43	44	69
31	HELAT35	203960 4/26/99	Uni-ZAP XR	41	2168	1	2168	215	215	90	1			20
32	HELAT58	203960 4/26/99	Uni-ZAP XR	42	2105	1	2105	215	215	91	1			22



Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
33	HELAW94	203960 4/26/99	Uni-ZAP XR	43	1491	1	1491		182	92	1			15
34	HEBDF90	203960 4/26/99	Uni-ZAP XR	44	2902	1273	2710	1461	1461	93	1	19	20	40
35	HELDF80	203960 4/26/99	Uni-ZAP XR	45	1460	1	1460	37	37	94	1			10
36	HELDK79	203960 4/26/99	Uni-ZAP XR	46	896	1	896		316	95	1	26	27	68
37	HELDQ42	203960 4/26/99	Uni-ZAP XR	47	2070	49	2070	712	712	96	1			26
38	HELEE85	203960 4/26/99	Uni-ZAP XR	48	569	1	569	75	75	97	1			10
39	HELEL76	203960 4/26/99	Uni-ZAP XR	49	1255	1	1255	42	42	98	1	21	22	195
40	HELFO30	203960 4/26/99	Uni-ZAP XR	50	2354	565	2354	607	607	99	1	16	17	41
41	HTEGH03	203960 4/26/99	Uni-ZAP XR	51	2267	91	2248	156	156	100	1	20	21	33
42	HELGF28	203960 4/26/99	Uni-ZAP XR	52	2298	1	2298	255	255	101	1			21
43	HELGG84	203960 4/26/99	Uni-ZAP XR	53	1109	12	1109	147	147	102	1	16	17	22

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
44	HELGP60	203960 4/26/99	Uni-ZAP XR	54	1296	1	1296	171	171	103	1	21	22	29
45	HELHP11	203960 4/26/99	Uni-ZAP XR	55	1432	1	1302	293	293	104	1	21	22	38
46	HELAC55	203960 4/26/99	Uni-ZAP XR	56	979	1	979	188	188	105	1	35	36	38
47	HEMCJ80	203960 4/26/99	Uni-ZAP XR	57	2667	195	2512	323	323	106	1	24	25	78
48	HEMCL55	203960 4/26/99	Uni-ZAP XR	58	1356	1	1356	270	270	107	1			8
49	HEMDB07	203960 4/26/99	Uni-ZAP XR	59	850	1	850	99	99	108	1	25	26	34

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing) and the translated SEQ ID NO:Y (where Y may be any of the polypeptide sequences disclosed in the sequence listing) are sufficiently

- accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also
- 5 hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used, for example, to generate antibodies which bind specifically to proteins containing the polypeptides and the secreted proteins encoded by the cDNA clones identified in Table 1.
- 10 Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the
- 15 actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).
- Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the
- 20 generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods.
- 25 The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.
- 30 The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed

herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or a deposited clone, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the secreted protein.

The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or a cDNA contained in ATCC deposit Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide encoded by the cDNA contained in ATCC deposit Z. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide sequence encoded by the cDNA contained in ATCC deposit Z are also encompassed by the invention.

### 10 Signal Sequences

The present invention also encompasses mature forms of the polypeptide having the polypeptide sequence of SEQ ID NO:Y and/or the polypeptide sequence encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species of the protein. Further, it has long been known that cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, Virus Res. 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, Nucleic Acids Res. 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of

predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, supra.) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

5 In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., Protein Engineering 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis  
10 of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence  
15 shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present  
20 invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of  
25 directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present  
30 invention.

#### **Polynucleotide and Polypeptide Variants**

The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X, the complementary strand thereto, and/or the cDNA sequence contained in a deposited clone.

5 The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y and/or encoded by a deposited clone.

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

10 The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence contained in a deposited cDNA clone or the  
15 complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited clone, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein).

Polynucleotides which hybridize to these nucleic acid molecules under stringent  
20 hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

The present invention is also directed to polypeptides which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% identical to, for example, the polypeptide sequence  
25 shown in SEQ ID NO:Y, the polypeptide sequence encoded by the cDNA contained in a deposited clone, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described herein).

By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended  
30 that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In



other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using

the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, an amino acid sequences shown in Table 1 (SEQ ID NO:Y) or to the amino acid sequence encoded by cDNA contained in a deposited clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10  
5 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100  
10 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not  
15 matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter  
20 the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host  
25 (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These  
30 allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., J. Biol. Chem. 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions,

inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues,

where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a peptide or polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

### **Polynucleotide and Polypeptide Fragments**

The present invention is also directed to polynucleotide fragments of the polynucleotides of the invention.

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence which: is a portion of that contained in a deposited clone, or encoding the polypeptide encoded by the cDNA in a deposited clone; is a portion of that shown in SEQ ID NO:X or the complementary strand thereto, or is a portion of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:Y. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in a deposited clone or the nucleotide sequence shown in SEQ ID NO:X. In this context "about" includes the particularly recited value, a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X, or the complementary strand thereto, or the cDNA contained in a deposited clone. In this context "about" includes the particularly recited ranges, and ranges larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to these nucleic acid molecules under



stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, and ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are

specifically contemplated by the present invention. Moreover, polynucleotides encoding these domains are also contemplated.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

Preferably, the polynucleotide fragments of the invention encode a polypeptide which demonstrates a functional activity. By a polypeptide demonstrating a "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) polypeptide of invention protein. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide of the invention for binding) to an antibody to the polypeptide of the invention], immunogenicity (ability to generate antibody which binds to a polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

The functional activity of polypeptides of the invention, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the invention for binding to an antibody of the polypeptide of the invention, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another

embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

5 In another embodiment, where a ligand for a polypeptide of the invention identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See  
10 generally, Phizicky, E., et al., 1995, Microbiol. Rev. 59:94-123. In another embodiment, physiological correlates of binding of a polypeptide of the invention to its substrates (signal transduction) can be assayed.

In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the  
15 invention and fragments, variants derivatives and analogs thereof to elicit related biological activity related to that of the polypeptide of the invention (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope of the invention.

## 20 **Epitopes and Antibodies**

The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of the polypeptide having an amino acid sequence of SEQ ID NO:Y, or an epitope of the polypeptide sequence encoded by a polynucleotide sequence contained in ATCC deposit No. Z or encoded by a polynucleotide that  
25 hybridizes to the complement of the sequence of SEQ ID NO:X or contained in ATCC deposit No. Z under stringent hybridization conditions or lower stringency hybridization conditions as defined supra. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X),  
30 polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize

to the complementary strand under stringent hybridization conditions or lower stringency hybridization conditions defined supra.

The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described infra. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998-4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross-reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985), further described in U.S. Patent No. 4,631,211).

In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in

immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid

surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

5           As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination  
10 thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*,  
15 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion desulfide bonds have also been found to be  
20 more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For  
25 example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine  
30 residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni<sup>2+</sup>

nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo and Blasco, *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

## Antibodies

Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of SEQ ID NO:Y, and/or an epitope, of the present invention (as determined by immunoassays well known in the art for assaying specific antibody-antigen binding). Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a

Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules  
5 that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

Most preferably the antibodies are human antigen-binding antibody fragments  
10 of the present invention and include, but are not limited to, Fab, Fab' and F(ab')<sub>2</sub>, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region,  
15 CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, ship rabbit, goat, guinea pig, camel, horse, or  
20 chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

25 The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO  
30 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).



Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in  
5 contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in  
10 terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described  
15 herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%,  
20 less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific  
25 antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide  
30 of the invention. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-2}$  M,  $10^{-2}$  M,  $5 \times 10^{-3}$  M,  $10^{-3}$  M,  $5 \times 10^{-4}$  M,  $10^{-4}$  M,  $5 \times 10^{-5}$  M,  $10^{-5}$  M,  $5 \times 10^{-6}$  M,  $10^{-6}$  M,  $5 \times 10^{-7}$  M,  $10^{-7}$  M,  $5 \times 10^{-8}$  M,  $10^{-8}$  M,  $5 \times 10^{-9}$

M,  $10^{-9}$  M,  $5 \times 10^{-10}$  M,  $10^{-10}$  M,  $5 \times 10^{-11}$  M,  $10^{-11}$  M,  $5 \times 10^{-12}$  M,  $10^{-12}$  M,  $5 \times 10^{-13}$  M,  $10^{-13}$  M,  $5 \times 10^{-14}$  M,  $10^{-14}$  M,  $5 \times 10^{-15}$  M, or  $10^{-15}$  M.

The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described  
5 herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes  
10 antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand  
15 binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described  
20 supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

25 The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as  
30 well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor

agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., *Blood* 92(6):1981-1988 (1998); Chen et al., *Cancer Res.* 58(16):3668-3678 (1998); Harrop et al., *J. Immunol.* 161(4):1786-1794 (1998); Zhu et al., *Cancer Res.* 58(15):3209-3214 (1998); Yoon et al., *J. Immunol.* 160(7):3170-3179 (1998); Prat et al., *J. Cell. Sci.* 111(Pt2):237-247 (1998); Pitard et al., *J. Immunol. Methods* 205(2):177-190 (1997); Liautard et al., *Cytokine* 9(4):233-241 (1997); Carlson et al., *J. Biol. Chem.* 272(17):11295-11301 (1997); Taryman et al., *Neuron* 14(4):755-762 (1995); Muller et al., *Structure* 6(9):1153-1167 (1998); Bartunek et al., *Cytokine* 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

Antibodies of the present invention may be used, for example, but not limited to, to purify, detect, and target the polypeptides of the present invention, including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include

5 antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation,

10 metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of-interest can be produced by various procedures well known in the art. For example, a polypeptide of

15 the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum

20 hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques

25 known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and*

30 *T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term

"monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma  
5 technology are routine and well known in the art and are discussed in detail in the Examples (e.g., Example 16). In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum; the mouse spleen is harvested and splenocytes isolated. The  
10 splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of  
15 antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized  
20 with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')<sub>2</sub> fragments of the invention may be  
25 produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). F(ab')<sub>2</sub> fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated  
30 using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such

phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')<sub>2</sub> fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and Sawai et al., AJRI 34:26-34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu et al., PNAS 90:7995-7999 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including

in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO

98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can  
5 express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes.

10 The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce  
15 chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human  
20 immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a  
25 detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are  
30 incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to



provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a  
5 selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., *Bio/technology* 12:899-903 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using  
10 techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, *FASEB J.* 7(5):437-444; (1989) and Nissinoff, J. *Immunol.* 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or  
15 binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

20

#### Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or lower  
25 stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y.

The polynucleotides may be obtained, and the nucleotide sequence of the  
30 polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described

in Kutmeier et al., *BioTechniques* 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

5           Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library  
10           generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a  
15           cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

          Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated  
20           using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, *Current Protocols in Molecular Biology*, John  
25           Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

          In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the  
30           complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using

routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in E. coli may also be used (Skerra et al., Science 242:1038-1041 (1988)).

### Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably,  
5 by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a  
10 polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by  
15 expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo  
20 genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT  
25 Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an  
30 antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous

promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene* 45:101 (1986); Cockett et al., *Bio/Technology* 8:2 (1990)).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the

generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding  
5 sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such  
10 fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

15 In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin  
20 promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric  
25 gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for  
30 efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure

translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al.,  
5 Methods in Enzymol. 153:51-544 (1987)).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein.

10 Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript,  
15 glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

20 For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences,  
25 transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to  
30 form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule.

Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

- A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell 11:223 (1977)),
- 5 hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., Cell 22:817 (1980)) genes can be employed in tk-, hgp<sup>rt</sup>- or ap<sup>rt</sup>- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., Natl. Acad. Sci.
- 10 USA 77:357 (1980); O'Hare et al., Proc. Natl. Acad. Sci. USA 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 Clinical Pharmacy 12:488-505; Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science
- 15 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, 1993, TIB TECH 11(5):155-215); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre et al., Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al.
- 20 (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13; Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., J. Mol. Biol. 150:1 (1981), which are incorporated by reference herein in their entireties.
- 25 The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in
- 30 culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).



The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g.,

Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are incorporated by reference in their entireties.

5           The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant  
10   region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be  
15   made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991);  
20   Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337- 11341(1992) (said references incorporated by reference in their entireties).

          As discussed, supra, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to  
25   the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various  
30   domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., Nature 331:84-86 (1988). The polypeptides of the present invention fused or conjugated to an antibody having

disulfide- linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., J. Molecular Recognition 8:52-58 (1995); Johanson et al., J. Biol. Chem. 270:9459-9471 (1995)).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., Cell 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment

thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish  
5 peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material  
10 includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{111}\text{In}$  or  $^{99}\text{Tc}$ .

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or  
15 a radioactive metal ion, e.g., alpha-emitters such as, for example,  $^{213}\text{Bi}$ . A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-  
20 dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU),  
25 cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

30 The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein

or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor,  $\alpha$ -interferon,  $\beta$ -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- $\alpha$ , TNF- $\beta$ , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGI (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16 (Academic Press 1985), and Thorpe *et al.*, "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62:119-58 (1982).

Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

## 5 Immunophenotyping

The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or  
10 maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with  
15 antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to  
20 prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

## Assays For Antibody Binding

25 The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions,  
30 gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well

known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

5           Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X- 100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell  
10   lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the  
15   art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

20           Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-  
25   fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or  
30   alkaline phosphatase) or radioactive molecule (e.g., <sup>32</sup>P or <sup>125</sup>I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can

be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

5 ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be  
10 conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the  
15 coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

20 The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g.,  $^3\text{H}$  or  $^{125}\text{I}$ ) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the  
25 antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g.,  $^3\text{H}$  or  $^{125}\text{I}$ ) in the presence of  
30 increasing amounts of an unlabeled second antibody.

#### Therapeutic Uses



The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of

products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

- 5 It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will
- 10 preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-2}$  M,  $10^{-2}$  M,  $5 \times 10^{-3}$  M,  $10^{-3}$  M,  $5 \times 10^{-4}$  M,  $10^{-4}$  M,  $5 \times 10^{-5}$  M,  $10^{-5}$  M,  $5 \times 10^{-6}$  M,  $10^{-6}$  M,  $5 \times 10^{-7}$  M,  $10^{-7}$  M,  $5 \times 10^{-8}$  M,  $10^{-8}$  M,  $5 \times 10^{-9}$  M,  $10^{-9}$  M,  $5 \times 10^{-10}$  M,  $10^{-10}$  M,  $5 \times 10^{-11}$  M,  $10^{-11}$  M,  $5 \times 10^{-12}$  M,  $10^{-12}$  M,  $5 \times 10^{-13}$  M,  $10^{-13}$  M,  $5 \times 10^{-14}$  M,  $10^{-14}$  M,  $5 \times 10^{-15}$  M, and  $10^{-15}$  M.
- 15

### Gene Therapy

- In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or
- 20 prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

- 25 Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

- For general reviews of the methods of gene therapy, see Goldspiel et al., Clinical Pharmacy 12:488-505 (1993); Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science
- 30 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, TIBTECH 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al.

(eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990).

In a preferred aspect, the compound comprises nucleic acid sequences  
5 encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue- specific. In another particular embodiment,  
10 nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989). In specific  
15 embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid- carrying vectors, or  
20 indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be  
25 accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun;  
30 Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by

administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current*

Opinion in Genetics and Development 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., Human Gene Therapy 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., Science 252:431-434 (1991); Rosenfeld et al., Cell 68:143- 155 (1992); Mastrangeli et al., J. Clin. Invest. 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., Gene Therapy 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be  
5 determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages,  
10 neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

15 In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and  
20 maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

In a specific embodiment, the nucleic acid to be introduced for purposes of  
25 gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription. Demonstration of Therapeutic or Prophylactic Activity

The compounds or pharmaceutical compositions of the invention are  
30 preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition

include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the  
5 invention, in vitro assays which can be used to determine whether administration of a specific compound is indicated, include in vitro cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

#### 10 Therapeutic/Prophylactic Administration and Composition

The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably an antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from  
15 substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above;  
20 additional appropriate formulations and routes of administration can be selected from among those described herein below.

Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-  
25 mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion  
30 or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In

addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, polymeric materials can be used (see Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., Macromol. Sci. Rev. Macromol. Chem. 23:61 (1983); see also Levy et al., Science 228:190 (1985); During et al., Ann. Neurol. 25:351 (1989); Howard et al., J. Neurosurg. 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose



(see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

5           In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered in vivo to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by  
10 use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA  
15 for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term  
"pharmaceutically acceptable" means approved by a regulatory agency of the Federal  
20 or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as  
25 peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel,  
30 sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering

agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as  
5 pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier  
10 so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous  
15 administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such  
20 as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

25 The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol,  
30 histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant

expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. **Diagnosis and Imaging**

Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or disorders associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an

increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99}\text{Tc}$ ); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for

unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of <sup>99m</sup>Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body

scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI). Kits

The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the

kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may  
5 also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The  
10 diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled  
15 monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing  
20 unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is detected by  
25 incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include  
30 non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group.

Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

### **Fusion Proteins**

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), resulting in



chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian

5 immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).)

10 Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively,  
15 deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See,  
20 D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine  
25 peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope  
30 derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

### **Vectors, Host Cells, and Protein Production**

5           The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host  
10   cells.

          The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line  
15   and then transduced into host cells.

          The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The  
20   expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

25           As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in E. coli and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as E. coli,  
30   Streptomyces and Salmonella typhimurium cells; fungal cells, such as yeast cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as

CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, 5 pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

10 Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that the 15 polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, 20 phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can 25 also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant 30 production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-

mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some  
5 proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have  
10 been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control  
15 regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination, resulting in the formation of a new transcription unit (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; U.S. Patent No. 5,733,761, issued March 31, 1998; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650,  
20 published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra et al., *Nature* 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, *Proteins: Structures and*  
25 *Molecular Principles*, W.H. Freeman & Co., N.Y., and Hunkapiller et al., *Nature*, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a polypeptide sequence of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence.  
30 Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid,  $\alpha$ -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid,  $\gamma$ -Abu,  $\epsilon$ -Ahx, 6-amino hexanoic acid,

Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoro-amino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptides which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, *e.g.*, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent NO: 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated  
5 molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

10 The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., Exp. Hematol. 20:1028-1035 (1992) (reporting  
15 pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues;  
20 those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

25 One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be  
30 performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the

N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for  
5 derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention  
10 relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, *Therapeutics*) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

15 Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone (including fragments, variants, splice variants, and fusion proteins, corresponding to these polypeptides as described herein). These  
20 homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the  
25 multimer of the invention is a homodimer (*e.g.*, containing polypeptides having identical or different amino acid sequences) or a homotrimer (*e.g.*, containing polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

30 As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (*i.e.*, polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the

invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, 5 ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed 10 when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid 15 residues contained in the polypeptide sequence ( e.g., that recited in the sequence listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the 20 consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 25 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for 30 example, osteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide



linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

5           Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al.,  
10   Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant  
15   fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

          Trimeric polypeptides of the invention may offer the advantage of enhanced  
20   biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be  
25   employed in preparing trimeric polypeptides of the invention.

          In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, associations  
30   proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent  
5 Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by  
10 reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally,  
15 techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic  
20 engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are  
25 generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by  
30 reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal

peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

## 5 **Uses of the Polynucleotides**

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

10 The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

15 Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an  
20 amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping  
25 strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread.  
30 This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see

Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the measured gene expression level with a standard level of polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder.

In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the present invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the present invention, where each probe has one strand containing a 31' mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

Where a diagnosis of a disorder, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the present invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

By "measuring the expression level of polynucleotide of the present invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the present invention or the level of the mRNA encoding the polypeptide in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder. As will be appreciated in the art, once a standard polypeptide

level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains the polypeptide of the present invention or mRNA. As indicated, biological samples include body fluids (such as semen, lymph, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and other tissue sources found to express the polypeptide of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides are attached to a solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene chip with polynucleotides of the present invention attached may be used to identify polymorphisms between the polynucleotide sequences, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, including cancerous diseases and conditions. Such a method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced supra are hereby incorporated by reference in their entirety herein.

The present invention encompasses polynucleotides of the present invention that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, Science 254, 1497 (1991); and M. Egholm, O. Buchardt,

L.Christensen, C. Behrens, S. M. Freier, D. A. Driver, R. H. Berg, S. K. Kim, B. Norden, and P. E. Nielsen, *Nature* 365, 666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because  
5 there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined  
10 with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ( $T_{sub.m}$ ) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

15 The present invention is useful for detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic  
20 leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

Pathological cell proliferative disorders are often associated with inappropriate  
25 activation of proto-oncogenes. (Germann, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in *Neoplastic Diseases of the Blood*, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now believed to result from the qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the  
30 chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Germann et al., supra) It is likely that mutated or altered expression of specific genes is involved in the

pathogenesis of some leukemias, among other tissues and cell types. (Gelman et al., supra) Indeed, the human counterparts of the oncogenes involved in some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Gelman et al., supra)

- 5 For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580) However, it has been shown that exposure of HL-60 cells to a DNA  
10 construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness would not be  
15 limited to treatment of proliferative disorders of hematopoietic cells and tissues, in light of the numerous cells and cell types of varying origins which are known to exhibit proliferative phenotypes.

- In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Antisense  
20 techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the  
25 polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991) ) or to the mRNA itself (antisense  
30 - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while



antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

5           Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome,  
10       thereby producing a new trait in the host cell.

          The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more  
15       restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA  
20       markers for RFLP.

          The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using  
25       this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

          Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological  
30       samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art

technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an  
5 identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with  
10 tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

15 In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA  
20 antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

### **Uses of the Polypeptides**

Each of the polypeptides identified herein can be used in numerous ways. The  
25 following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods: (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-  
30 3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known

in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99\text{m}}\text{Tc}$ ), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

5           In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers  
10   for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

          A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example,  $^{131}\text{I}$ ,  
15    $^{112}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ ), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a  
20   human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99\text{m}}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging:  
25   The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

          Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a  
30   standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high

amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ  
5 preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Moreover, polypeptides of the present invention can be used to treat disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to  
10 supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used  
15 in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease. For example, administration of an antibody directed to a  
20 polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration  
25 columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

30

### **Gene Therapy Methods**

Another aspect of the present invention is to gene therapy methods for treating disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the invention that operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Belldgrun et al., J. Natl. Cancer Inst., 85:207-216 (1993); Ferrantini et al., Cancer Research, 53:107-1112 (1993); Ferrantini et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura et al., Cancer Research 50: 5102-5106 (1990); Santodonato, et al., Human Gene Therapy 7:1-10 (1996); Santodonato, et al., Gene Therapy 4:1246-1255 (1997); and Zhang, et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the polynucleotide of the invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations,

lipofectin or precipitating agents and the like. However, the polynucleotides of the invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and  
5 5,580,859, which are herein incorporated by reference.

The polynucleotide vector constructs of the invention used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene;  
10 pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of polynucleotide sequence of the invention. Suitable promoters  
15 include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters,  
20 such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotides of the invention.

Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the  
25 polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the invention can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver,  
30 spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular,

fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph  
5 fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated  
10 cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked *nucleic acid* sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20  
15 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

20 The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the  
25 procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

30 The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

In certain embodiments, the polynucleotide constructs of the invention are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a  
5 tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA , 86:6077-6081 (1989), which is herein incorporated  
10 by reference); and purified transcription factors (Debs et al., J. Biol. Chem., 265:10189-10192 (1990), which is herein incorporated by reference), in functional form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are  
15 particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

20 Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication NO: WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., Felgner et al., Proc.  
25 Natl. Acad. Sci. USA, 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol,  
30 phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP



starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC),  
dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine  
5 (DOPE) can be used in various combinations to make conventional liposomes, with or  
without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be  
prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas  
into a sonication vial. The sample is placed under a vacuum pump overnight and is  
hydrated the following day with deionized water. The sample is then sonicated for 2  
10 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an  
inverted cup (bath type) probe at the maximum setting while the bath is circulated at  
15EC. Alternatively, negatively charged vesicles can be prepared without sonication  
to produce multilamellar vesicles or by extrusion through nucleopore membranes to  
produce unilamellar vesicles of discrete size. Other methods are known and available  
15 to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar  
vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred.  
The various liposome-nucleic acid complexes are prepared using methods well known  
in the art. See, e.g., Straubinger et al., *Methods of Immunology*, 101:512-527 (1983),  
20 which is herein incorporated by reference. For example, MLVs containing nucleic  
acid can be prepared by depositing a thin film of phospholipid on the walls of a glass  
tube and subsequently hydrating with a solution of the material to be encapsulated.  
SUVs are prepared by extended sonication of MLVs to produce a homogeneous  
population of unilamellar liposomes. The material to be entrapped is added to a  
25 suspension of preformed MLVs and then sonicated. When using liposomes containing  
cationic lipids, the dried lipid film is resuspended in an appropriate solution such as  
sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and  
then the preformed liposomes are mixed directly with the DNA. The liposome and  
DNA form a very stable complex due to binding of the positively charged liposomes  
30 to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are  
prepared by a number of methods, well known in the art. Commonly used methods  
include  $\text{Ca}^{2+}$ -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta*,

394:483 (1975); Wilson et al., Cell , 17:77 (1979)); ether injection (Deamer et al., Biochim. Biophys. Acta, 443:629 (1976); Ostro et al., Biochem. Biophys. Res. Commun., 76:836 (1977); Fraley et al., Proc. Natl. Acad. Sci. USA, 76:3348 (1979)); detergent dialysis (Enoch et al., Proc. Natl. Acad. Sci. USA , 76:145 (1979)); and  
5 reverse-phase evaporation (REV) (Fraley et al., J. Biol. Chem., 255:10431 (1980); Szoka et al., Proc. Natl. Acad. Sci. USA , 75:145 (1978); Schaefer-Ridder et al., Science, 215:166 (1982)), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the  
10 ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent NO: 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469  
15 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

20 In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding polypeptides of the invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis  
25 virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-  
30 19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy , 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any

means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and  $\text{CaPO}_4$  precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

5           The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding polypeptides of the invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express polypeptides of the invention.

          In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with  
10   polynucleotides of the invention contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses polypeptides of the invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional  
15   mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al., Am. Rev. Respir. Dis., 109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld et al., Science, 252:431-434 (1991);  
20   Rosenfeld et al., Cell, 68:143-155 (1992)). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al. Proc. Natl. Acad. Sci. USA, 76:6606 (1979)).

          Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel., 3:499-503 (1993);  
25   Rosenfeld et al., Cell, 68:143-155 (1992); Engelhardt et al., Human Genet. Ther., 4:759-769 (1993); Yang et al., Nature Genet., 7:362-369 (1994); Wilson et al., Nature, 365:691-692 (1993); and U.S. Patent NO: 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and  
30   constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other

varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or  
5 packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

10 In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, Curr. Topics in Microbiol. Immunol., 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base  
15 pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will  
20 include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct containing polynucleotides of the invention is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells  
25 which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct of the invention.  
30 These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express the desired gene product.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding the polypeptide sequence of interest) via homologous recombination (see, e.g., U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA, 86:8932-8935 (1989); and Zijlstra et al., Nature, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

5       The polynucleotides encoding polypeptides of the present invention may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, VEGF-2 (VEGF-C), VEGF-3 (VEGF-B), epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth  
10   factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

      Preferably, the polynucleotide encoding a polypeptide of the invention contains a secretory signal sequence that facilitates secretion of the protein.  
15   Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

20       Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available  
25   depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppository solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the  
30   rat livers. (Kaneda et al., Science, 243:375 (1989)).

      A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is

administered by direct injection into or locally within the area of arteries.

Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide  
5 construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

Therapeutic compositions useful in systemic administration, include  
10 recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection,  
15 aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA , 189:11277-11281 (1992), which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present  
20 invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

25 Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of  
30 polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian. Therapeutic compositions of

the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly

## 5 **Biological Activities**

The polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological  
10 activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists could be used to treat the associated disease.

## **Immune Activity**

The polynucleotides or polypeptides, or agonists or antagonists of the present  
15 invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells.  
20 The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

25 A polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. A polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those  
30 disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia



telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary

Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitis, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a  
5 polynucleotides or polypeptides, or agonists or antagonists of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host  
10 disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of a polynucleotides or polypeptides, or  
agonists or antagonists of the present invention that inhibits an immune response,  
15 particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polynucleotides or  
polypeptides, or agonists or antagonists of the present invention may also be used to  
modulate inflammation. For example, the polypeptide or polynucleotide or agonists  
20 or antagonist may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-  
25 mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

### **Hyperproliferative Disorders**

30 A polynucleotides or polypeptides, or agonists or antagonists of the invention can be used to treat or detect hyperproliferative disorders, including neoplasms. A polynucleotides or polypeptides, or agonists or antagonists of the present invention

may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

- 5           For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an  
10   immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

- Examples of hyperproliferative disorders that can be treated or detected by a polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast,  
15   digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

- Similarly, other hyperproliferative disorders can also be treated or detected by  
20   a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's  
Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia,  
25   located in an organ system listed above.

          One preferred embodiment utilizes polynucleotides of the present invention to inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

- Thus, the present invention provides a method for treating cell proliferative  
30   disorders by inserting into an abnormally proliferating cell a polynucleotide of the present invention, wherein said polynucleotide represses said expression.

Another embodiment of the present invention provides a method of treating cell-proliferative disorders in individuals comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA  
5 construct comprising a recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is  
10 hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic,  
15 specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated (i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

20 Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes " is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational  
25 modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of  
30 cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as; but not

limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference. In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as decreases in the rate of proliferation or growth of the cells. The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

The present invention is further directed to antibody-based therapies which involve administering of anti-polypeptides and anti-polynucleotide antibodies to a mammalian, preferably human, patient for treating one or more of the described disorders. Methods for producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere  
5    herein. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the  
10    present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without  
15    undue experimentation.

In particular, the antibodies, fragments and derivatives of the present invention are useful for treating a subject having or developing cell proliferative and/or differentiation disorders as described herein. Such treatment comprises administering a single or multiple doses of the antibody, or a fragment, derivative, or a conjugate  
20    thereof.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors, for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will  
25    preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-6}M$ ,  $10^{-6}M$ ,  $5 \times 10^{-7}M$ ,  $10^{-7}M$ ,  $5 \times 10^{-8}M$ ,  $10^{-8}M$ ,  $5 \times 10^{-9}M$ ,  $10^{-9}M$ ,  $5 \times 10^{-10}M$ ,  
30    or  $10^{-10}M$ .

$10^{-10}$ M,  $10^{-10}$ M,  $5 \times 10^{-11}$ M,  $10^{-11}$ M,  $5 \times 10^{-12}$ M,  $10^{-12}$ M,  $5 \times 10^{-13}$ M,  $10^{-13}$ M,  $5 \times 10^{-14}$ M,  $10^{-14}$ M,  $5 \times 10^{-15}$ M, and  $10^{-15}$ M.

Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference).

Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference).

Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12 (1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by reference).

Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues.

Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby  
5 incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

In another embodiment, the invention provides a method of delivering compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or polypeptide antibodies associated with heterologous  
10 polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

15 Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to  
20 enhance the immune response (e.g. chemokines), to said antigens and immunogens.

#### **Cardiovascular Disorders**

Polynucleotides or polypeptides, or agonists or antagonists of the invention may be used to treat cardiovascular disorders, including peripheral artery disease,  
25 such as limb ischemia.

Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart,  
30 dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus,



and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilog of Fallot, ventricular heart septal defects.

Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, 5 angiodyplasia, angiomas, bacillary angiomas, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, 10 hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, ataxia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous 15 insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, 20 carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral 25 arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, 30 migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and

thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment  
5 syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

10 Polynucleotides or polypeptides, or agonists or antagonists of the invention, are especially effective for the treatment of critical limb ischemia and coronary disease.

Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous  
15 injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppository solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides of the invention may be administered as part of a *Therapeutic*,  
20 described in more detail below. Methods of delivering polynucleotides of the invention are described in more detail herein.

#### **Anti-Angiogenesis Activity**

The naturally occurring balance between endogenous stimulators and  
25 inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally  
30 delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases.

A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 5 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, 10 *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can be treated with the 15 polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating an angiogenesis-related disease and/or 20 disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat a cancer or tumor. Cancers which may be treated with polynucleotides, polypeptides, 25 antagonists and/or agonists include, but are not limited to solid tumors, including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced 30 malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically,

in order to treat cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating other disorders, besides cancers, which involve angiogenesis. These disorders include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

For example, within one aspect of the present invention methods are provided for treating hypertrophic scars and keloids, comprising the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably

initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

Moreover, Ocular disorders associated with neovascularization which can be treated with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalmol.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalmol.* 22:291-312 (1978).

Thus, within one aspect of the present invention methods are provided for treating neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of disorders can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and

administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the goal of the administration would be to place the composition at the advancing front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

Within another aspect of the present invention, methods are provided for treating neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the

compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

Within another aspect of the present invention, methods are provided for treating retrolental fibroplasia, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

Additionally, disorders which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

Moreover, disorders and/or states, which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb



angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have  
5 angiogenesis as a pathologic consequence such as cat scratch disease (Rochele minalia quintosa), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and  
10 fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

15 Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present  
20 invention a compositions (in the form of, for example, a spray or film) may be utilized to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat  
25 tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized  
30 during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide, agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration. Within particularly preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

The polynucleotides, polypeptides, agonists and/or agonists of the present invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl

complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten  
5 oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its  
10 hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and  
15 sugars.

A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan  
20 Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin;  
25 Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum;  
30 alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-

316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolimidazole; and metalloproteinase inhibitors such as BB94.

### **Diseases at the Cellular Level**

5 Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by the polynucleotides or polypeptides and/or antagonists or agonists of the invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer,  
10 melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary  
15 cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred  
embodiments, the polynucleotides or polypeptides, and/or agonists or antagonists of  
20 the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by the polynucleotides or polypeptides, or agonists or  
antagonists of the invention, include, but are not limited to, progression, and/or  
25 metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic  
lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and  
30 non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and  
carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma,

osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal  
 5 cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder  
 10 carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by the polynucleotides or polypeptides, and/or agonists or antagonists of the  
 15 invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus  
 20 erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestosis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by  
 25 alcohol), septic shock, cachexia and anorexia.

### **Wound Healing and Epithelial Cell Proliferation**

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing the polynucleotides or polypeptides, and/or agonists  
 30 or antagonists of the invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds.

Polynucleotides or polypeptides, as well as agonists or antagonists of the invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal  
5 ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, and/or agonists or antagonists of  
10 the invention, could be used to promote dermal reestablishment subsequent to dermal loss

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are a non-  
15 exhaustive list of grafts that polynucleotides or polypeptides, agonists or antagonists of the invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepidermic grafts, avascular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft, epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft,  
20 homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omentum graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, can be used to promote skin strength and to improve the appearance of aged skin.

25 It is believed that the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could promote proliferation of epithelial cells such as  
30 sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. The polynucleotides or polypeptides, and/or agonists

or antagonists of the invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may have a cytoprotective effect on the small intestine mucosa. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention,

could be used to treat diseases associate with the under expression of the polynucleotides of the invention.

Moreover, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to prevent and heal damage to the lungs due to various pathological states. A growth factor such as the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of aveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using the polynucleotides or polypeptides, and/or agonists or antagonists of the invention. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary displasia, in premature infants.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetraholoride and other hepatotoxins known in the art).

In addition, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

### **Neurological Diseases**



Nervous system disorders, which can be treated with the compositions of the invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to,

multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

In a preferred embodiment, the polypeptides, polynucleotides, or agonists or  
5 antagonists of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat or prevent neural cell injury associated with cerebral hypoxia. In one aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury  
10 associated with cerebral ischemia. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with cerebral infarction. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a stroke.  
15 In a further aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a heart attack.

The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in  
20 promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture; (2) increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, e.g., choline acetyltransferase or  
25 acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may routinely be measured using a method set forth herein or otherwise known in the art, such as, for example, the method set forth in Arakawa et al. (J. Neurosci.  
30 10:3507-3515 (1990)); increased sprouting of neurons may be detected by methods known in the art, such as, for example, the methods set forth in Pestronk et al. (Exp. Neurol. 70:65-82 (1980)) or Brown et al. (Ann. Rev. Neurosci. 4:17-42 (1981));

increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known in the art and depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include, but are not limited to, disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including, but not limited to, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

### **Infectious Disease**

A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polypeptide or polynucleotide and/or agonist or antagonist of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae,

Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papiloma virus, Papovaviridae, Parvoviridae, 5 Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiollitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue 10 syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and 15 viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists 20 or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat AIDS.

Similarly, bacterial or fungal agents that can cause disease or symptoms and 25 that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Cryptococcus neoformans, Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), 30 Bacteroidaceae, Blastomycosis, Bordetella, Borrelia (e.g., Borrelia burgdorferi), Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, E. coli (e.g., Enterotoxigenic E. coli and Enterohemorrhagic E.

coli), Enterobacteriaceae (Klebsiella, Salmonella (e.g., Salmonella typhi, and Salmonella paratyphi), Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Mycobacterium leprae, Vibrio cholerae, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal),

5 Meisseria meningitidis, Pasteurellacea Infections (e.g., Actinobacillus, Haemophilus (e.g., Haemophilus influenza type B), Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, Shigella spp., Staphylococcal, Meningococcal, Pneumococcal and Streptococcal (e.g., Streptococcus pneumoniae and Group B Streptococcus). These bacterial or fungal families can cause the following diseases or

10 symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning,

15 Typhoid, pneumonia, Gonorrhea, meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. Polynucleotides or polypeptides, agonists or

20 antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, Polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat: tetanus, Diphtheria, botulism, and/or meningitis type B.

Moreover, parasitic agents causing disease or symptoms that can be treated or

25 detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax,

30 Plasmodium falciparum, Plasmodium malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis),

liver disease, lung disease, opportunistic infections (e.g., AIDS related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or  
5 agonists or antagonists of the invention are used to treat malaria.

Preferably, treatment using a polypeptide or polynucleotide and/or agonist or antagonist of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the  
10 engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

### **Regeneration**

15 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis,  
20 osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic,  
25 and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage.  
30 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or

ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a  
5 polynucleotide or polypeptide and/or agonist or antagonist of the present invention to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral  
10 nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide and/or agonist or antagonist of the present invention.

15

### **Chemotaxis**

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils,  
20 epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase chemotactic activity of particular cells. These chemotactic  
25 molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts,  
30 which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may inhibit chemotactic activity. These molecules

could also be used to treat disorders. Thus, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention could be used as an inhibitor of chemotaxis.

5

### **Binding Activity**

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit  
10 (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural  
15 or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).). Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

20 Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially  
25 containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the  
30 candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product



mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

- 5            Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

- Additionally, the receptor to which a polypeptide of the invention binds can be  
10 identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF  
15 family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition  
20 site for a site-specific protein kinase.

            Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

- 25            As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein  
30 microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of polypeptides of the invention thereby effectively generating agonists and antagonists of polypeptides of the invention. See  
5 generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., et al., *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S. *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, L. O., et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated  
10 by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides of the invention may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired polynucleotide sequence of the invention molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides of the  
15 invention may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptides of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more  
20 heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B,  
25 decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the  
30 polypeptides of the invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide. The

biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention.

5 An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and 3[H] thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the  
10 compound to determine if the compound stimulates proliferation by determining the uptake of 3[H] thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography which measures the incorporation of 3[H] thymidine. Both agonist and antagonist compounds may be identified by this procedure.

15 In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following  
20 interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

25 All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from  
30 suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to the polypeptides of the invention comprising the steps of: (a) incubating a candidate binding compound with the polypeptide; and

(b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with the polypeptide, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

5           Also, one could identify molecules bind a polypeptide of the invention experimentally by using the beta-pleated sheet regions contained in the polypeptide sequence of the protein. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, the amino acid sequence of each beta pleated sheet regions in a disclosed  
10           polypeptide sequence. Additional embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, any combination or all of contained in the polypeptide sequences of the invention. Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the amino acid sequence of each of the beta  
15           pleated sheet regions in one of the polypeptide sequences of the invention. Additional embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions in one of the polypeptide sequences of the invention.

## 20           Targeted Delivery

          In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

          As discussed herein, polypeptides or antibodies of the invention may be  
25           associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic  
30           acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double

stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

### **Drug Screening**

Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any

of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids  
5 expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

Thus, the present invention provides methods of screening for drugs or any  
10 other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically  
15 labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present  
20 invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and  
25 washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays  
30 in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of

any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

### **Antisense And Ribozyme (Antagonists)**

5 In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained a deposited clone. In one embodiment, antisense sequence is generated internally by the organism, in another embodiment, the antisense sequence is separately administered  
10 (see, for example, O'Connor, Neurochem., 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense  
15 Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research, 6:3073 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

20 For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed in vitro by incubating cells with the oligoribonucleotide. A similar procedure for in vivo use is described in WO 91/15580. Briefly, a pair of  
25 oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl<sub>2</sub>, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then  
30 ligated to the EcoRI/Hind III site of the retroviral vector PMV7 (WO 91/15580).

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA

oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid of the invention. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding a polypeptide of the invention, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature*, 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell*, 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature*, 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of interest. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids of the invention, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA sequence of the



invention it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, *e.g.*,  
5 the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., *Nature*, 372:333-335 (1994). Thus, oligonucleotides complementary to either the 5' - or 3' -  
10 non- translated, non-coding regions of a polynucleotide sequence of the invention could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but  
15 could be used in accordance with the invention. Whether designed to hybridize to the 5' -, 3' - or coding region of mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

20 The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*,  
25 for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci.*, 84:648-652 (1987); PCT Publication NO: WO88/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication NO: WO89/10134, published April 25, 1988), hybridization-  
30 triggered cleavage agents. (See, *e.g.*, Krol et al., *BioTechniques*, 6:958-976 (1988)) or intercalating agents. (See, *e.g.*, Zon, *Pharm. Res.*, 5:539-549 (1988)). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide,

hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., Nucl. Acids Res., 15:6625-6641 (1987)). The oligonucleotide is a 2-O-methylribonucleotide (Inoue et al., Nucl. Acids Res., 15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett. 215:327-330 (1987)).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (Nucl. Acids Res., 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., Proc. Natl. Acad. Sci. U.S.A., 85:7448-7451 (1988)), etc.

While antisense nucleotides complementary to the coding region sequence of the invention could be used, those complementary to the transcribed untranslated region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science, 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs corresponding to the polynucleotides of the invention, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5' -UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature, 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within each nucleotide sequence disclosed in the sequence listing. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA corresponding to the polynucleotides of the invention; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the polynucleotides of the invention in vivo. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the

ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular  
5 concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation  
10 or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirous in cases such as restenosis after balloon  
15 angioplasty.

The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat the diseases described herein.

20 Thus, the invention provides a method of treating disorders or diseases, including but not limited to the disorders or diseases listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present  
25 invention

### **Other Activities**

The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as  
30 thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide

may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

The polypeptide of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. The polypeptide of the invention may have the ability to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

The polypeptide of the invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, the polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

The polypeptide of the invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues.

The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, polypeptides or

polynucleotides and/or agonist or antagonists of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

5 Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

10 Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

#### 15 **Other Preferred Embodiments**

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

20 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

25 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

30 Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the

range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

- 5           Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

- Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500  
10 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

- A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about  
15 the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

          A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

- 20           Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

- 25           Also preferred is a composition of matter comprising a DNA molecule which comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

- 30           Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA

Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the  
5 complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule  
10 comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete  
15 nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X  
20 wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and  
25 determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence  
30 selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected



from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as

defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least  
5 one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence  
10 selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in  
15 Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide  
20 molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample; if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected  
25 from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

30 Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid

sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone

identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

In specific embodiments of the invention, for each "Contig ID" listed in the fourth column of Table 2, preferably excluded are one or more polynucleotides comprising, or alternatively consisting of, a nucleotide sequence referenced in the fifth column of Table 2 and described by the general formula of a-b, whereas a and b are uniquely determined for the corresponding SEQ ID NO:X referred to in column 3 of Table 2. Further specific embodiments are directed to polynucleotide sequences excluding one, two, three, four, or more of the specific polynucleotide sequences referred to in the fifth column of Table 2. In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby incorporated by reference in their entirety.

**TABLE 2**

Gene No.	cDNA Clone ID	NT SEQ ID NO: X	Contig ID	Public Accession Numbers
1	HE9NB82	11	790925	T48271, T48272, T84138, R00180, R00181, R01779, R01778, R49669, R49669, H42920, R93248, H52219, H62631, H75401, H75956, N58710, AA011389, AA011426
3	HE9TA42	13	834972	T74758, T74846, R50277, R50737, R52504, R52505, R76987, H05272, H12752, H12793, R98348, R98349, H68391, H84958, H91439, N49227, N50277, N52913, N54419, N55604, W32435, W45010, W56052, W72113, W94473, AA011029, AA011151, AA017109, AA017110, AA017244, AA017280, AA019069, AA019071, AA019150, AA019152, AA056090, AA056117, AA056151, AA058595, AA088508, AA088509, AA132442, AA132610, AA150049, AA150034, AA459829, AA463911, AA419089, AA419062, AA557492, H85529, AA766136, AA928412, AA995492, C15582
12	HEBAP51	22	631248	T75214, R45278, R45278, N45592, N48800, W56090
14	HEBBF78	24	742915	T39268, T60160, T60220, R35222, R35223, R44510, R44510, R59970, H00372, H17643, H17757, H21224, R99021, H59535, H60407, H63093, H63178, H70834, H72562, H72964, H82418, H94858, H94913, H97157, N32571, N41911, N53689, N69676, N73052, N78595, W78722, W80529, AA024462, AA054989, AA100617, AA187655, AA191585, AA194889.

				AA223950, AA227494, AA242922, AA253197, AA252244, AA417824, AA427942
18	HEBDF77	28	692347	R12748, R39800, H06705, H06754, H29013, R88622, N46729, AA133829
19	HEBDQ91	29	840288	T71210, T81885, R85100, AA167586, AA824367, Z41111, Z45398
21	HEBFL36	31	604638	R38261
25	HEBGE85	35	834478	R42024, R43307, R52913, R42024, R43307, R67278, H19248, N62208, N76506, W72237, W74222, W92450, W95152, W95249, AA047719, AA057767, AA191673, AA232863, AA256929, AA255438, AA258256, AA262360, AA421841
26	HEEAB58	36	834398	T39482, T40618, T50039, T50096, R20546, R20546, R66341, R66342, R67483, R82598, H01566, H54508, H66304, H67729, H79785, H82064, H82164, H84518, H97729, N20279, N29188, N29395, N35601, N59339, N70870, N71706, N77147, AA029343, AA029344, AA040150, AA040814, AA507067, AA507956, AA534713, AA534743, AA535536, AA688253, AA906644, AA961588, D78835, C01311, C06150, C17188
29	HEGAU15	39	834379	W92008, W92009
30	HEBCI18	40	831464	T63400, T63545, T94026, T94778, T75424, R39278, R82600, R82599, R87540, R87937, R87938, AA039787, AA039788, AA176798, AA836133, AI057154
33	HELAW94	43	834796	AI003772, W45510
34	HEBDF90	44	835835	R21211, R74193, R74282, R81220, H02254, H02352, H88945, H89178, N53357, N63150, N66663, N92903, N98946, W21109, AA044128, AA044302, AA058311, AA167044, AA805148, AA810643, AA249069, AA398057, AA399111
37	HELDQ42	47	695716	T49898, H19034, H38523, H83345, AA015878, AA018778, AA018035, AA045767, AA099965
40	HELFO30	50	828202	R54828, R55169, N35347, N44715, N52355, N52362, N75125, N75133, N94993, W04599, W15275, W19462, W20219, W31499, W32433, W39235, W45008, W56137, W55956, N90579, AA028004, AA029907, AA030036, AA039829, AA044387, AA042839, AA056023, AA056071, AA424153, AA424192, AA512882, D79420, D79933, C00297
41	HTEGH03	51	815562	T60134, T92770, R33256, R33257, R36501, R37497, R63928, R64019, R81559, R81560, R82368, R82427, H01029, H01778, H24484, H95209, H95210, N24611, N25469, N26138, N26846, N27847, N33750, N91796, W21524, W56252, AA031345, AA031389, AA054690, AA056448, AA057272, AA058717, AA075396, AA075446, AA102851, AA102852, AA132449, AA157836, AA426489, AA430130
43	HELGG84	53	674456	N66810, N70665, N75004, W05747
47	HEMCJ80	57	765041	T54337, R20254, R43710, R43710, N23462, AA039793, AA056228, AA429186



Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

5

### Examples

#### Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited</u>
	<u>Plasmid</u>	
	Lambda Zap	pBluescript (pBS)
	Uni-Zap XR	pBluescript (pBS)
20	Zap Express	pBK
	lafmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
	pCMVSPORT 3.0	pCMVSPORT 3.0
25	pCR <sup>®</sup> 2.1	pCR <sup>®</sup> 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and

pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.

Vectors pSport1, pCMVSport 2.0 and pCMVSport 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR<sup>®</sup>2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly

isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported.

- 5 The oligonucleotide is labeled, for instance, with  $^{32}\text{P}$ - $\gamma$ -ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as
- 10 those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., Molecular Cloning: A Laboratory
- 15 Manual, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

- Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the
- 20 desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25  $\mu\text{l}$  of reaction mixture with 0.5  $\mu\text{g}$  of the above cDNA template. A convenient reaction mixture is 1.5-5 mM  $\text{MgCl}_2$ , 0.01% (w/v) gelatin, 20  $\mu\text{M}$  each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR
- 25 (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the
- 30 DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods

include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

**Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide**

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

5

**Example 3: Tissue Distribution of Polypeptide**

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P<sup>32</sup> using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

15

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70 degree C overnight, and the films developed according to standard procedures.

20

**Example 4: Chromosomal Mapping of the Polynucleotides**

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose

25

30

gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

**Example 5: Bacterial Expression of a Polypeptide**

5           A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product  
10   into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance ( $Amp^r$ ), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning  
15   sites.

          The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which  
20   expresses the lacI repressor and also confers kanamycin resistance ( $Kan^r$ ). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

          Clones containing the desired constructs are grown overnight (O/N) in liquid  
25   culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.<sup>600</sup>) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to  
30   increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4 degree C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded  
5 onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH  
10 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl.  
15 Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250  
20 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4 degree C or frozen at -80 degree C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements  
25 operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains:  
1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The  
30 origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR  
5 primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

10

**Example 6: Purification of a Polypeptide from an Inclusion Body**

The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10 degree C.

15 Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10 degree C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution  
20 containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by  
25 centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4 degree  
30 C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with



20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4 degree C without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential  
5 filtration unit equipped with 0.16 um membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a  
10 stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak  
15 anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are  
20 collected under constant  $A_{280}$  monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 ug of purified  
25 protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

**Example 7: Cloning and Expression of a Polypeptide in a Baculovirus**  
30 **Expression System**

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine

procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200  $\mu$ l of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded

in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with  
5 the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of <sup>35</sup>S-methionine and 5 uCi <sup>35</sup>S-cysteine (available from Amersham) are added.  
10 The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the  
15 produced protein.

#### **Example 8: Expression of a Polypeptide in Mammalian Cells**

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and  
20 signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early  
25 promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109),  
30 pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human HeLa, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1,

Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a  
5 selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of  
10 interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279  
(1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers,  
15 the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the  
20 expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the  
25 cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by  
30 procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to

produce the secreted protein, the vector does not need a second signal peptide.

Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a  
5 commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector  
10 are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five  $\mu$ g of the expression plasmid pC6 a pC4 is cotransfected with 0.5  
15  $\mu$ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in  
20 alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing  
25 even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

### 30 **Example 9: Protein Fusions**

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For

example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization  
5 signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused  
10 protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These  
15 primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with  
20 BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted  
25 protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

30 GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCCACCGTGC  
CCAGCACCTGAATTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAAA  
CCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGT

GGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG  
 ACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA  
 CAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACT  
 GGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA  
 5 ACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC  
 CACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAG  
 GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGT  
 GGAGTGGGAGAGCAATGGGCAGCCGGAGAACAATAAGACCACGCCT  
 CCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTG  
 10 GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCA  
 TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG  
 GTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

#### **Example 10: Production of an Antibody from a Polypeptide**

15 The antibodies of the present invention can be prepared by a variety of  
 methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells  
 expressing a polypeptide of the present invention is administered to an animal to  
 induce the production of sera containing polyclonal antibodies. In a preferred  
 method, a preparation of the secreted protein is prepared and purified to render it  
 20 substantially free of natural contaminants. Such a preparation is then introduced into  
 an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are  
 monoclonal antibodies (or protein binding fragments thereof). Such monoclonal  
 antibodies can be prepared using hybridoma technology. (Köhler et al., Nature  
 25 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J.  
 Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell  
 Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures  
 involve immunizing an animal (preferably a mouse) with polypeptide or, more  
 preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in  
 30 any suitable tissue culture medium; however, it is preferable to culture cells in Earle's  
 modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at



about 56 degrees C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

It will be appreciated that Fab and F(ab')<sub>2</sub> and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in

the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 5 314:268 (1985).)

**Example 11: Production Of Secreted Protein For High-Throughput Screening Assays**

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well 15 plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

20 Plate 293T cells (do not carry cells past P+20) at  $2 \times 10^5$  cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine 25 (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I 30 mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem

I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degrees C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of  $\text{CaCl}_2$  (anhyd); 0.00130 mg/L  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ; 0.050 mg/L of  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ ; 0.417 mg/L of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ; 311.80 mg/L of KCl; 28.64 mg/L of  $\text{MgCl}_2$ ; 48.84 mg/L of  $\text{MgSO}_4$ ; 6995.50 mg/L of NaCl; 2400.0 mg/L of  $\text{NaHCO}_3$ ; 62.50 mg/L of  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ ; 71.02 mg/L of  $\text{Na}_2\text{HPO}_4$ ; .4320 mg/L of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine- $\text{H}_2\text{O}$ ; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL- $\text{H}_2\text{O}$ ; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL- $\text{H}_2\text{O}$ ; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalainine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2 $\text{H}_2\text{O}$ ; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and

0.680 mg/L of Vitamin B<sub>12</sub>; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin  
5 complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

10 The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degrees C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml  
15 deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing  
20 expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

### **Example 12: Construction of GAS Reporter Construct**

25 One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

30 GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six

members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at  
5 higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2,  
10 Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two  
15 groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

20 Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the  
25 proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATS</u>	<u>GAS(elements) or ISRE</u>
<u>IFN family</u>							
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
<u>gp130 family</u>							
10	IL-6 (Pleiotrophic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrophic)	?	+	?	?	1,3	
	OnM(Pleiotrophic)	?	+	+	?	1,3	
	LIF(Pleiotrophic)	?	+	+	?	1,3	
	CNTF(Pleiotrophic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrophic)	?	+	?	?	1,3	
	IL-12(Pleiotrophic)	+	-	+	+	1,3	
<u>g-C family</u>							
	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
20	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25							
<u>gp140 family</u>							
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30							
<u>Growth hormone family</u>							
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35							
<u>Receptor Tyrosine Kinases</u>							
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5' :GCGCCTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5' :CTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

### **Example 13: High-Throughput Screening Assay for T-cell Activity.**

The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.



Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1% Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells ( $10^7$  per transfection), and resuspend in OPTI-MEM to a final concentration of  $10^7$  cells/ml. Then add 1ml of  $1 \times 10^7$  cells in OPTI-MEM to T25 flask and incubate at 37 degrees C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing polypeptides of the invention and/or induced polypeptides of the invention as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100,000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the  
5 assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophane covers) and  
10 stored at -20 degrees C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4 degrees C and serve as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the  
15 positive control wells.

The above protocol may be used in the generation of both transient, as well as, stable transfected cells, which would be apparent to those of skill in the art.

#### **Example 14: High-Throughput Screening Assay Identifying Myeloid Activity**

20 The following protocol is used to assess myeloid activity by determining whether polypeptides of the invention proliferates and/or differentiates myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is  
25 U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest  $2 \times 10^7$  U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing  
30 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM Na<sub>2</sub>HPO<sub>4</sub>·7H<sub>2</sub>O, 1 mM MgCl<sub>2</sub>, and 675 uM CaCl<sub>2</sub>. Incubate at 37 degrees C for 45 min.

- 5 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degrees C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 10 These cells are tested by harvesting  $1 \times 10^8$  cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of  $5 \times 10^5$  cells/ml. Plate 200 ul cells per well in the 96-well plate (or  $1 \times 10^5$  cells/well).

- Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37 degrees C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

20 **Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.**

- When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat phenochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor).

- 30 The EGR1 gene expression is activated during this treatment. Thus, by stably

transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene  
5 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified  
10 product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30  
15 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-  
20 inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine  
25 protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80%  
30 confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as  $5 \times 10^5$  cells/ml.

- 5           Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to  $1 \times 10^5$  cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the
- 10           supernatant according to Example 17.

**Example 16: High-Throughput Screening Assay for T-cell Activity**

- NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and
- 15           CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

- 20           In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.

- 25           Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF-KB would be useful in treating diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid
- 30           arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC  
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

10 5':GCGGCAAGCTTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene)

15 Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5' : CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCCATCTGCCATCTCAATTA  
GTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATT  
TCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATT  
20 CCAGAAGTAGTGAGGAGGCTTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT:3' (SEQ ID  
NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and  
25 HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly,  
30 the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly,

the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

#### 5 **Example 17: Assay for SEAP Activity**

As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

- 10 Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

- Cool the samples to room temperature for 15 minutes. Empty the dispenser  
15 and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at  
20 each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

#### **Reaction Buffer Formulation:**

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6

23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

**Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability**

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.



For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO<sub>2</sub> incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

- 5        A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO<sub>2</sub> incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

- 10       For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10<sup>6</sup> cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10<sup>6</sup> cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is  
15 then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

- 20       To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular  
25 Ca<sup>++</sup> concentration.

#### **Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity**

- 30       The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies.

In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor  
5 dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily  
10 of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins  
15 capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr  
20 with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of  
25 cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

30 To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20

minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na<sub>3</sub>VO<sub>4</sub>, 2 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub> and a cocktail of protease inhibitors (# 1836170) obtained from

- 5 Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4 degrees C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on  
10 ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degrees C at 16,000 x g.

- Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described  
15 here.

- Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and  
20 PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

- The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg<sub>2</sub><sup>+</sup> (5mM ATP/50mM MgCl<sub>2</sub>), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride,  
25 pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl<sub>2</sub>, 5 mM MnCl<sub>2</sub>, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degrees C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

- The tyrosine kinase assay reaction is then terminated by adding 10 ul of  
30 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degrees C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times.

- 5 Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degrees C for one hour. Wash the well as above.

- Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the  
10 absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

#### **Example 20: High-Throughput Screening Assay Identifying Phosphorylation**

##### **Activity**

- As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine  
20 phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

- 25 Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules,  
30 this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degrees C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts  
5 filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the  
10 Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

15

**Example 21: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide**

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from  
20 these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

25 PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct  
30 sequencing.

PCR products is cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United

States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

#### **Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample**

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in

Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

### **Example 23: Formulation**

The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a Therapeutic. By therapeutic is meant a polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

The Therapeutic will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the Therapeutic administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Therapeutics can be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray.

"Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials



(for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al.,  
5 Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release Therapeutics also include liposomally entrapped  
10 Therapeutics of the invention (*see* generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang  
15 et al., Proc. Natl. Acad. Sci.(USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the  
20 optimal Therapeutic.

In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

25 Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically  
30 acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both.

5 Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

10 The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten  
15 residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar  
20 alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or  
25 stabilizers will result in the formation of polypeptide salts.

Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a  
30 stopper pierceable by a hypodermic injection needle.

Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized

formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

5           The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use  
10       or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

          The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, alum, alum plus  
15       deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG, and MPL. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to,  
20       Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B,  
25       whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic  
30       mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate

administration of one of the compounds or agents given first, followed by the second.

The Therapeutics of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not  
5 limited to, other members of the TNF family, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are  
10 administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

15 In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta),  
20 OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF),  
25 and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International  
30 Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors.

Nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

In other embodiments, Therapeutics of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In

another specific embodiment, Therapeutics of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment,

5 Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat

10 or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with

15 ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment,

20 Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial infection.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered

25 with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin,

30 beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin,

erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

Conventional nonspecific immunosuppressive agents, that may be  
5 administered in combination with the Therapeutics of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

In specific embodiments, Therapeutics of the invention are administered in  
10 combination with immunosuppressants. Immunosuppressants preparations that may be administered with the Therapeutics of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMMUNE™/NEORAL™/SANGDYA™ (cyclosporin), PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucocorticosteroids, and RAPAMUNE™ (sirolimus). In a specific  
15 embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the  
20 Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, and GAMIMUNE™. In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

25 In an additional embodiment, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives,  
30 arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid,

amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

In another embodiment, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., mephallen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

In a specific embodiment, Therapeutics of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, Therapeutics of the invention are administered in combination with Rituximab. In a further embodiment, Therapeutics of the invention are administered with Rituxmab and CHOP, or Rituxmab and any combination of the components of CHOP.

In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3,



IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

In an additional embodiment, the Therapeutics of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PlGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the Therapeutics of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

In an additional embodiment, the Therapeutics of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but

are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

In additional embodiments, the Therapeutics of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example,  
5 radiation therapy.

**Example 24: Method of Treating Decreased Levels of the Polypeptide**

The present invention relates to a method for treating an individual in need of an increased level of a polypeptide of the invention in the body comprising  
10 administering to such an individual a composition comprising a therapeutically effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the  
15 secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily  
20 dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

**Example 25: Method of Treating Increased Levels of the Polypeptide**

25 The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

30 In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of

decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer. For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

#### **Example 26: Method of Treatment Using Gene Therapy-Ex Vivo**

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is

maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

5           The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the  
10 packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is  
15 removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been  
20 efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

#### 25   **Example 27: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention**

Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent  
30 NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935

(1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting  
5 sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the  
10 amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

15 The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size  
20 fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral  
25 sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the  
30 polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining  
5 cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na<sub>2</sub> HPO<sub>4</sub>, 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains  
10 approximately 3X10<sup>6</sup> cells/ml. Electroporation should be performed immediately following resuspension.

Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with  
15 HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3'end. Two non-coding sequences are amplified via PCR: one non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3'end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5'end and a HindIII site at the 3'end. The CMV promoter and the  
20 fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap  
25 (Bio-Rad). The final DNA concentration is generally at least 120 µg/ml. 0.5 ml of the cell suspension (containing approximately 1.5X10<sup>6</sup> cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 µF and 250-300 V, respectively. As voltage increases, cell survival decreases, but  
30 the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following  
5 day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a  
10 patient as described above.

#### **Example 28: Method of Treatment Using Gene Therapy - In Vivo**

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the  
15 introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in  
20 the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 94(12):3281-3290 (1996) (incorporated herein by reference).

25 The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

30 The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or

precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg



body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence  
5 can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or  
10 mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard  
15 recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on  
20 the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

25 After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA  
30 in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper

dosages and other treatment parameters in humans and other animals using naked DNA.

**Example 29: Transgenic Animals.**

5           The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, *e.g.*, baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are  
10       used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

          Any technique known in the art may be used to introduce the transgene (*i.e.*, polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear  
15       microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic  
20       stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, *e.g.*, Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-  
25       mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

          Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into  
30       enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding

strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of  
5 heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of  
10 interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating  
15 such conditions and/or disorders.

### **Example 30: Knock-Out Animals.**

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (*E.g.*,  
20 see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding  
25 regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination,  
30 results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas &

Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

5 In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not  
10 limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention,  
15 e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve  
20 expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the  
25 body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

30 When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For

example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

### 10 **Example 31: Production of an Antibody**

#### a) Hybridoma Technology

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing polypeptide(s) of the invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide(s) of the invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

Monoclonal antibodies specific for polypeptide(s) of the invention are prepared using hybridoma technology. (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized with polypeptide(s) of the invention or, more preferably, with a secreted polypeptide-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line

(SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which  
5 secrete antibodies capable of binding the polypeptide(s) of the invention.

Alternatively, additional antibodies capable of binding to polypeptide(s) of the invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In  
10 accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by polypeptide(s) of the invention. Such antibodies comprise anti-  
15 idiotypic antibodies to the protein-specific antibody and are used to immunize an animal to induce formation of further protein-specific antibodies.

For in vivo use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing  
20 chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268  
25 (1985).)

b) Isolation Of Antibody Fragments Directed Against  
Polypeptide(s) From A Library Of scFvs

Naturally occurring V-genes isolated from human PBLs are constructed into a  
30 library of antibody fragments which contain reactivities against polypeptide(s) of the invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

### Rescue of the Library.

A library of scFvs is constructed from the RNA of human PBLs as described in PCT publication WO 92/01047. To rescue phage displaying antibody fragments, approximately 10<sup>9</sup> E. coli harboring the phagemid are used to inoculate 50 ml of  
5 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2 x 10<sup>8</sup> TU of delta gene 3 helper (M13 delta gene III, see PCT publication WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is  
10 centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 µg/ml ampicillin and 50 µg/ml kanamycin and grown overnight. Phage are prepared as described in PCT publication WO 92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody  
15 fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in  
20 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 µm filter (Minisart NML; Sartorius) to give a final concentration of approximately 10<sup>13</sup> transducing units/ml  
25 (ampicillin-resistant clones).

### Panning of the Library.

Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 µg/ml or 10 µg/ml of a polypeptide of the present invention. Tubes are blocked with  
30 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 10<sup>13</sup> TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with



PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 µg/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

#### Characterization of Binders.

Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., PCT publication WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

#### **Example 32: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation**

Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can,

in combination with various co-stimulatory proteins. induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective  
5 ligands CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the  
10 detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

In Vitro Assay- Purified polypeptides of the invention, or truncated forms thereof, is assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the polypeptides of  
15 the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed Staphylococcus aureus Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as IL-2 and IL-15 synergize with SAC and IgM  
20 crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation. Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

25 Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added  $10^5$  B-cells suspended in culture medium (RPMI 1640 containing 10% FBS,  $5 \times 10^{-5}$ M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and  $10^{-5}$  dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The  
30 positive and negative controls are IL2 and medium respectively.

In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or 2 mg/Kg of a polypeptide of the invention, or truncated forms thereof. Mice receive this

treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with polypeptides of the invention identify the results of the activity of the polypeptides on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths, and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from mice treated with polypeptide is used to indicate whether the polypeptide specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

Likewise, a predicted consequence of increased mature B-cell representation in vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and polypeptide-treated mice.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

### **Example 33: T Cell Proliferation Assay**

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of  $^3\text{H}$ -thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100  $\mu\text{l}$ /well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1  $\mu\text{g}/\text{ml}$  in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10<sup>4</sup>/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of polypeptides of the invention (total volume 200  $\mu\text{l}$ ). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100  $\mu\text{l}$  of supernatant is removed and stored -20 degrees C for measurement of

IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 ul of medium containing 0.5 uCi of  $^3\text{H}$ -thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of  $^3\text{H}$ -thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative controls for the effects of polypeptides of the invention.

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

**Example 34: Effect of Polypeptides of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells**

Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and MHC class II antigens). Treatment with activating factors, such as TNF- $\alpha$ , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC $\gamma$ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of polypeptides of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells ( $10^6/\text{ml}$ ) are treated with increasing concentrations of polypeptides of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of polypeptides of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degreesC. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. Polypeptides, agonists, or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American

Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of  $2 \times 10^6$ /ml in PBS containing PI at a final concentration of 5  $\mu$ g/ml, and then incubated at room temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

15. Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of  $5 \times 10^5$  cells/ml with increasing concentrations of the a polypeptide of the invention and under the same conditions, but in the absence of the polypeptide. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of a polypeptide of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g, R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

Oxidative burst. Purified monocytes are plated in 96-w plate at  $2 \times 10^5$  cell/well. Increasing concentrations of polypeptides of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM

NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 µl 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H<sub>2</sub>O<sub>2</sub> produced by the macrophages, a standard curve of a H<sub>2</sub>O<sub>2</sub> solution of known molarity is performed for each experiment.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polypeptides, polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

### **Example 35: Biological Effects of Polypeptides of the Invention**

#### **Astrocyte and Neuronal Assays**

Recombinant polypeptides of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate a polypeptide of the invention's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of a polypeptide of the invention to

induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

#### Fibroblast and endothelial cell assays

5 Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated  
10 for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE<sub>2</sub> assays, the human lung fibroblasts are cultured at  
15 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or polypeptides of the invention with or without IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for PGE<sub>2</sub> by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to  
20 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without polypeptides of the invention IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

Human lung fibroblasts are cultured with FGF-2 or polypeptides of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth  
25 of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with polypeptides of the invention.

#### Parkinson Models

The loss of motor function in Parkinson's disease is attributed to a deficiency of  
30 striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic



projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP<sup>+</sup>) and released.

- 5 Subsequently, MPP<sup>+</sup> is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP<sup>+</sup> is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

- 10 It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989). Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

- Based on the data with FGF-2, polypeptides of the invention can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of a polypeptide of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm<sup>2</sup> on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

- 30 Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons

would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if a polypeptide of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the polypeptide may be involved in Parkinson's Disease.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 36: The Effect of Polypeptides of the Invention on the Growth of Vascular Endothelial Cells**

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at  $2-5 \times 10^4$  cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. A polypeptide having the amino acid sequence of SEQ ID NO:Y, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

An increase in the number of HUVEC cells indicates that the polypeptide of the invention may proliferate vascular endothelial cells.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 37: Stimulatory Effect of Polypeptides of the Invention on the Proliferation of Vascular Endothelial Cells**

For evaluation of mitogenic activity of growth factors, the colorimetric MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was performed (CellTiter 96 AQ, Promega). Cells are seeded in a 96-well plate (5,000 cells/well) in 0.1 mL serum-supplemented medium and are allowed to attach overnight.

After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF<sub>165</sub> or a polypeptide of the invention in 0.5% FBS) with or without Heparin (8 U/ml) are added to wells for 48 hours. 20 mg of MTS/PMS mixture (1:0.05) are added per well and allowed to incubate for 1 hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader. Background absorbance from control wells (some media, no cells) is subtracted, and seven wells are performed in parallel for each condition. See, Leak *et al. In Vitro Cell. Dev. Biol.* 30A:512-518 (1994).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 38: Inhibition of PDGF-induced Vascular Smooth Muscle Cell Proliferation Stimulatory Effect**

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation. Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6 mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse anti-BrdUrd antibody at 4 degrees C for 2 h after being exposed to denaturing solution and then incubated with the streptavidin-peroxidase and diaminobenzidine. After counterstaining with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-positive cells to the total cell number. In addition, the simultaneous detection of the BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for individual cells by the concomitant use of bright field illumination and dark field-UV fluorescent illumination. See, Hayashida et al., J. Biol. Chem. 6:271(36):21985-21992 (1996).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 39: Stimulation of Endothelial Migration**

This example will be used to explore the possibility that a polypeptide of the invention may stimulate lymphatic endothelial cell migration.

- 5           Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD; Falk, W., et al., J. Immunological Methods 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8 um (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours at room temperature and dried under sterile air. Test substances are diluted to appropriate
- 10           concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25 ul of the final dilution is placed in the lower chamber of the modified Boyden apparatus. Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber,  $2.5 \times 10^5$  cells suspended in 50 ul M199 containing 1%
- 15           FBS are seeded in the upper compartment. The apparatus is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO<sub>2</sub> to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by
- 20           counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.


- The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
- 25           invention.

**Example 40: Stimulation of Nitric Oxide Production by Endothelial Cells**

- Nitric oxide released by the vascular endothelium is believed to be a mediator of vascular endothelium relaxation. Thus, activity of a polypeptide of the invention can be
- 30           assayed by determining nitric oxide production by endothelial cells in response to the polypeptide.

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of a positive control (such as VEGF-1) and the polypeptide of the invention. Nitric oxide in the medium is determined by use of the Griess reagent to measure total nitrite after  
 5 reduction of nitric oxide-derived nitrate by nitrate reductase. The effect of the polypeptide of the invention on nitric oxide release is examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer is measured with a NO-specific polarographic electrode connected to a NO meter (Iso-NO, World Precision Instruments Inc.) (1049). Calibration of the NO elements is performed according to the  
 10 following equation:



The standard calibration curve is obtained by adding graded concentrations of  $\text{KNO}_2$  (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing KI and  $\text{H}_2\text{SO}_4$ . The specificity of the Iso-NO electrode to NO is previously determined by  
 15 measurement of NO from authentic NO gas (1050). The culture medium is removed and HUVECs are washed twice with Dulbecco's phosphate buffered saline. The cells are then bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates are kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C. The NO sensor probe is inserted vertically into the wells, keeping the tip of the electrode 2  
 20 mm under the surface of the solution, before addition of the different conditions.

S-nitroso acetyl penicillamin (SNAP) is used as a positive control. The amount of released NO is expressed as picomoles per  $1 \times 10^6$  endothelial cells. All values reported are means of four to six measurements in each group (number of cell culture wells). See, Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

25 The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 41: Effect of Polypeptides of the Invention on Cord Formation in Angiogenesis**

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to  
5 form capillary-like structures (hollow structures) when cultured *in vitro*.

CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a  
10 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 µl/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 µg Cell Applications' Chord Formation Medium containing control buffer or a polypeptide of the invention (0.1 to 100 ng/ml) and the cells are cultured for an  
15 additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

Commercial (R&D) VEGF (50 ng/ml) is used as a positive control.  $\beta$ -estradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

20 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

25 **Example 42: Angiogenic Effect on Chick Chorioallantoic Membrane**

Chick chorioallantoic membrane (CAM) is a well-established system to examine angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of polypeptides of the invention to stimulate angiogenesis in CAM can be examined.

30 Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese quail (*Coturnix coturnix*) are incubated at 37.8°C and 80% humidity. Differentiated CAM of 16-day-old chick and 13-day-old quail embryos is studied with the following methods.

On Day 4 of development, a window is made into the egg shell of chick eggs. The embryos are checked for normal development and the eggs sealed with cellotape. They are further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) are cut into disks of about 5 mm in diameter. Sterile and salt-free growth factors are dissolved in distilled water and about 3.3 mg/ 5 ml are pipetted on the disks. After air-drying, the inverted disks are applied on CAM. After 3 days, the specimens are fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They are photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described above. Controls are performed with carrier disks alone.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

#### **Example 43: Angiogenesis Assay Using a Matrigel Implant in Mouse**

*In vivo* angiogenesis assay of a polypeptide of the invention measures the ability of an existing capillary network to form new vessels in an implanted capsule of murine extracellular matrix material (Matrigel). The protein is mixed with the liquid Matrigel at 4 degree C and the mixture is then injected subcutaneously in mice where it solidifies. After 7 days, the solid "plug" of Matrigel is removed and examined for the presence of new blood vessels. Matrigel is purchased from Becton Dickinson Labware/Collaborative Biomedical Products.

When thawed at 4 degree C the Matrigel material is a liquid. The Matrigel is mixed with a polypeptide of the invention at 150 ng/ml at 4 degrees C and drawn into cold 3 ml syringes. Female C57Bl/6 mice approximately 8 weeks old are injected with the mixture of Matrigel and experimental protein at 2 sites at the midventral aspect of the abdomen (0.5 ml/site). After 7 days, the mice are sacrificed by cervical dislocation, the Matrigel plugs are removed and cleaned (i.e., all clinging membranes and fibrous tissue is removed). Replicate whole plugs are fixed in neutral buffered 10% formaldehyde, embedded in paraffin and used to produce sections for histological examination after staining with Masson's Trichrome. Cross sections from 3 different regions of each plug

are processed. Selected sections are stained for the presence of vWF. The positive control for this assay is bovine basic FGF (150 ng/ml). Matrigel alone is used to determine basal levels of angiogenesis.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

#### **Example 44: Rescue of Ischemia in Rabbit Lower Limb Model**

To study the in vivo effects of polynucleotides and polypeptides of the invention on ischemia, a rabbit hindlimb ischemia model is created by surgical removal of one femoral arteries as described previously (Takeshita *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). The excision of the femoral artery results in retrograde propagation of thrombus and occlusion of the external iliac artery. Consequently, blood flow to the ischemic limb is dependent upon collateral vessels originating from the internal iliac artery (Takeshita *et al.* *Am J. Pathol* 147:1649-1660 (1995)). An interval of 10 days is allowed for post-operative recovery of rabbits and development of endogenous collateral vessels. At 10 day post-operatively (day 0), after performing a baseline angiogram, the internal iliac artery of the ischemic limb is transfected with 500 mg naked expression plasmid containing a polynucleotide of the invention by arterial gene transfer technology using a hydrogel-coated balloon catheter as described (Riessen *et al.* *Hum Gene Ther.* 4:749-758 (1993); Leclerc *et al.* *J. Clin. Invest.* 90: 936-944 (1992)). When a polypeptide of the invention is used in the treatment, a single bolus of 500 mg polypeptide of the invention or control is delivered into the internal iliac artery of the ischemic limb over a period of 1 min. through an infusion catheter. On day 30, various parameters are measured in these rabbits: (a) BP ratio - The blood pressure ratio of systolic pressure of the ischemic limb to that of normal limb; (b) Blood Flow and Flow Reserve - Resting FL: the blood flow during undilated condition and Max FL: the blood flow during fully dilated condition (also an indirect measure of the blood vessel amount) and Flow Reserve is reflected by the ratio of max FL: resting FL; (c) Angiographic Score - This is measured by the angiogram of collateral vessels. A score is determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh; (d)



Capillary density - The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

The studies described in this example tested activity of polynucleotides and polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the agonists, and/or antagonists of the invention.

#### **Example 45: Effect of Polypeptides of the Invention on Vasodilation**

Since dilation of vascular endothelium is important in reducing blood pressure, the ability of polypeptides of the invention to affect the blood pressure in spontaneously hypertensive rats (SHR) is examined. Increasing doses (0, 10, 30, 100, 300, and 900 mg/kg) of the polypeptides of the invention are administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM. Statistical analysis are performed with a paired t-test and statistical significance is defined as  $p < 0.05$  vs. the response to buffer alone.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

#### **Example 46: Rat Ischemic Skin Flap Model**

The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. Expression of polypeptides of the invention, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- a) Ischemic skin
- b) Ischemic skin wounds
- c) Normal wounds

The experimental protocol includes:

- a) Raising a 3x4 cm, single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).
- b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).

c) Topical treatment with a polypeptide of the invention of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.

d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

5           The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

#### 10    **Example 47: Peripheral Arterial Disease Model**

Angiogenic therapy using a polypeptide of the invention is a novel therapeutic strategy to obtain restoration of blood flow around the ischemia in case of peripheral arterial diseases. The experimental protocol includes:

15           a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.

            b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3 weeks.

20           c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of expression of a polypeptide of the invention and histology. Biopsy is also performed on the other side of normal muscle of the contralateral hindlimb.

            The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

#### **Example 48: Ischemic Myocardial Disease Model**

30           A polypeptide of the invention is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after

coronary artery occlusion. Alteration of expression of the polypeptide is investigated in situ. The experimental protocol includes:

a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.

5        b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4 weeks.

c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

10        The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

15        **Example 49: Rat Corneal Wound Healing Model**

This animal model shows the effect of a polypeptide of the invention on neovascularization. The experimental protocol includes:

a)        Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.

20        b)        Inserting a spatula below the lip of the incision facing the outer corner of the eye.

c)        Making a pocket (its base is 1-1.5 mm from the edge of the eye).

d) Positioning a pellet, containing 50ng- 5ug of a polypeptide of the invention, within the pocket.

25        e)        Treatment with a polypeptide of the invention can also be applied topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

30        The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

**Example 50: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing****Models****A. Diabetic db+/db+ Mouse Model.**

To demonstrate that a polypeptide of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All

manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

5           Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and  
10           iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to  
15           treatment, wounds are gently cleansed with sterile saline and gauze sponges.

          Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is  
20           no longer visible and the wound is covered by a continuous epithelium.

          A polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

          Animals are euthanized on day 8 with an intraperitoneal injection of sodium  
25           pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

          Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

30           Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post

treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

5

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with a polypeptide of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

10

15

Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

20

Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer can serve as a positive tissue control and human brain tissue can be used as a negative tissue control. Each specimen includes a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

25

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

30

### ***B. Steroid Impaired Rat Model***

The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-

Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)).

Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *Am. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", *In: Antiinflammatory Steroid Action: Basic and Clinical Aspects*, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", *In: Antiinflammatory Steroid Action: Basic and Clinical Aspects*, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that a polypeptide of the invention can accelerate the healing process, the effects of multiple topical applications of the polypeptide on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue

punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

5           Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

10           The polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

          Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for  
15 histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

          Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

20           Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:

25

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

          Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus  
30 microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by



treatment with a polypeptide of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of  $< 0.05$  is considered significant.

- 5        The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

10    **Example 51: Lymphadema Animal Model**

- The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of a polypeptide of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, 15 quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

- Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. 20 Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's 25 Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

- Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the 30 lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places. Analysis is performed in a blind manner.

**Circumference Measurements:** Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

**Volumetric Measurements:** On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

**Blood-plasma protein measurements:** Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and  $\text{Ca}^{2+}$  comparison.

Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

- 5        Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

- 10       The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

15       **Example 52: Suppression of TNF alpha-induced adhesion molecule expression by a Polypeptide of the Invention**

- 20       The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The
- 25       local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

- 30       Tumor necrosis factor alpha (TNF-a), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

- 30       The potential of a polypeptide of the invention to mediate a suppression of TNF-a induced CAM expression can be examined. A modified ELISA assay which uses ECs as a

solid phase absorbent is employed to measure the amount of CAM expression on TNF- $\alpha$  treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO<sub>2</sub>. HUVECs are seeded in 96-well plates at concentrations of  $1 \times 10^4$  cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 ul of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 ul volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100  $\mu$ l of 0.1% paraformaldehyde-PBS(with Ca<sup>++</sup> and Mg<sup>++</sup>) is added to each well. Plates are held at 4°C for 30 min.

Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10  $\mu$ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10  $\mu$ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

Then add 20  $\mu$ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100  $\mu$ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 ( $10^0$ ) >  $10^{0.5}$  >  $10^{-1}$  >  $10^{-1.5}$ . 5  $\mu$ l of each dilution is added to triplicate wells and the resulting AP content in each well is

5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl of pNPN reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties.

320

Applicant's or agent's file reference number	PS514PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>72</u> , line <u>N/A</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>26 April 1999</u>	Accession Number <u>203960</u>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States) <u>Europe</u> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")     	

For receiving Office use only	
<input type="checkbox"/>	This sheet was received with the international application
Authorized officer	

For International Bureau use only	
<input type="checkbox"/>	This sheet was received by the International Bureau on:
Authorized officer	

**ATCC Deposit No.: 203960**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: 203960****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.



***What Is Claimed Is:***

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
  - (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
  - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X, having biological activity;
  - (f) a polynucleotide which is a variant of SEQ ID NO:X;
  - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
  - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
  - (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.
- 5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 20 6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 25 7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
- 30 10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:
- (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
  - 5 (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
  - (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
  - (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included  
10 in ATCC Deposit No:Z;
  - (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
  - (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
  - 15 (g) a variant of SEQ ID NO:Y;
  - (h) an allelic variant of SEQ ID NO:Y; or
  - (i) a species homologue of the SEQ ID NO:Y.
12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-  
20 terminus or the N-terminus.
13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.
- 25 14. A recombinant host cell that expresses the isolated polypeptide of claim 11.
15. A method of making an isolated polypeptide comprising:
- (a) culturing the recombinant host cell of claim 14 under conditions such that  
30 said polypeptide is expressed; and
  - (b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount  
5 of the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of  
10 claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to  
15 a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.  
20

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

(a) contacting the polypeptide of claim 11 with a binding partner; and

(b) determining whether the binding partner effects an activity of the  
25 polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

22. A method of identifying an activity in a biological assay, wherein the  
30 method comprises:

(a) expressing SEQ ID NO:X in a cell;

(b) isolating the supernatant;

- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

<110> Human Genome Sciences, Inc.

<120> 49 Human Secreted Proteins

<130> PS514PCT

<140> Unassigned

<141> 2000-03-16

<150> 60/125,361

<151> 1999-03-19

<150> 60/169,910

<151> 1999-12-10

<160> 184

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

```

gggataccgga gcccaaatct tctgacaaaa ctcacacatg cccaccgtgc ccagcacctg      60
aattcgaggg tgcaccgtca gtcttctctt tcccccaaaa acccaaggac accctcatga      120
tctcccgga ccttgaggtc acatgcgtgg tgggtggacgt aagccacgaa gaccctgagg      180
tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg      240
aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact      300
ggctgaatgg caaggagtac aagtgcgaagg tctccaacaa agccctccca acccccatcg      360
agaaaaccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac accctgcccc      420
catcccgga tgagctgacc aagaaccagg tcagcctgac ctgcctgggc aaaggcttct      480
atccaagcga catcgccgtg gagtgggaga gcaatgggca gccggagaac aactacaaga      540
ccacgcctcc cgtgctggac tccgacggct ccttcttctt ctacagcaag ctcaccgtgg      600
acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat gaggtctctg      660
acaaccacta cagcagaag agcctctccc tgtctccggg taaatgagtg cgacggccgc      720
gactctagag gat                                     733

```

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

```

Trp Ser Xaa Trp Ser
1                               5

```

<210> 3  
 <211> 86  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60  
 cccgaaatat ctgccatctc aattag 86

<210> 4  
 <211> 27  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 gcggcaagct ttttgcaaag cctaggc 27

<210> 5  
 <211> 271  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 ctcgagatatt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60  
 aaatatctgc catctcaatt agtcagcaac catagtcctcg cccctaactc cgcccatccc 120  
 gcccctaact ccgcccagtt ccgcccattc tccgccccat ggctgactaa ttttttttat 180  
 ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240  
 ttttgagggc ctaggctttt gcaaaaagct t 271

<210> 6  
 <211> 32  
 <212> DNA  
 <213> Homo sapiens

<400> 6  
 gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7  
 <211> 31  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8  
 <211> 12  
 <212> DNA  
 <213> Homo sapiens

<400> 8  
ggggactttc cc

12

<210> 9  
<211> 73  
<212> DNA  
<213> Homo sapiens

<400> 9  
gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg  
ccatctcaat tag

60

73

<210> 10  
<211> 256  
<212> DNA  
<213> Homo sapiens

<400> 10  
ctcgagggga ctttcccggg gactttccgg ggactttccg ggactttcca tctgccatct  
caattagtca gcaaccatag tcccggccct aactccgccc atcccggccc taactccgccc  
cagttccgccc cattctccgc cccatggctg actaattttt tttatttatg cagaggccga  
ggccgcctcg gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg  
cttttgcaaa aagctt

60

120

180

240

256

<210> 11  
<211> 1329  
<212> DNA  
<213> Homo sapiens

<400> 11  
aatacaggcg ctgcacagcc tggccccacct gttcattaat gcactcaatt tagtactgaa  
tgggtctttct cccagcccat tcccagccct tcctatttcc tttcctattt tttttttctc  
cccacacttt cttgggactc ccaccttgga aggaggaagg gctgacctgg gttctctcca  
gcccccaggt gcgcggggtc acccgtgccc cttcattatg gacctgggccc ctaccggaac  
ccctgcccc a gttaccacaa ctcaggccgg ctggccccgg ccatgggctg cgcaaatcac  
cagcccccaa cccaggggagg aactggcccc tcctagggag cctcttcgac ttttttagaa  
aaatgatctc cttttcttcc cagccatgat gtttagtaaa ttttttagt accgcactta  
gcagacagct ttccaagtgt gctttcttgc cacaaaagtg tcctggcaag agccccttat  
ttttaagaca tcaggaagcc agaccgctt gagttgggag aattttgtag ctcaacatat  
caagtctctg atggtatctg agctgcccac acccccacct gccagggcc cacagagccc  
aaaacagaag ggggctgccc cagccagcag agcacagagt ttctggagct cccatccaca  
gatgcaggag ggggtactga tggtaacccc catgtggatt tgagggcagc agtcctggc  
ctcaccttag ccagcctggg tggctcccta gcccgaagag gccaggaagg gctggaaggc  
agggcctgca ggtgctcccc gccctgagac ccaggcccca aatcagcaat aatgaacaaa  
cccttgggccc agcctgggct ggtgacctgg gcaccagaga ccttgcatcc ctccctatcc  
taggaggccc ctaggggtgc cccatctcag tgtcccctga actctttatt tgccataatt  
atatatatat atatgagata tataaatata tataaaatag ctattttgct taaatttcta  
cagtatgtaa aagtgaaaaa atgatgaaga cgggtgcacc tgtctgagtt tggccctcat  
gtgagctgtg cccttccctc tcctcatgcc cccttccagc ggcttctgcc aaccatgggg  
ggctggacca ccatggccac tgacccagcc cctcagaatc ccacactcca atcctttcca  
tttcagttta gtcctaaaag ttcatacacag ggtctttctt tctactccag gactgggttt

60

120

180

240

300

360

420

480

540

600

660

720

780

840

900

960

1020

1080

1140

1200

1260



gttttttatat atataaaaaa aaaaaaaaaa aactcgaggg ggggcccggg acgcaattcc 1320  
caaggagtt 1329

<210> 12

<211> 834

<212> DNA

<213> Homo sapiens

<400> 12

ggcagcagct	ctcttgctga	gagtcacaggc	ctgcctgcct	gccagccaga	ggtgcccctt	60
atgcacatcc	agtcctacca	gcctgcttcc	aggagcctag	agtgggaaag	gcacctgggg	120
gagccctggt	gaagggctgg	acctgaccca	tggaggcagg	aggaagacct	ggctttcttg	180
gtttgtccgg	ccctggggag	cagccttccc	tactcacttt	ccccttgccg	caccaaattg	240
tcactccttc	cagagcaaaa	actggcactc	atagcaagcc	agaaccccca	tccccatcct	300
catgggtctg	tggctgggtg	caggccatca	aaatggacac	cacgagacag	aagtggggac	360
tgcctggcca	cctagcgcc	tcccactcct	taagcaagca	caaagaagat	gaggcagaga	420
attgtcagag	ctgaaagtgc	ctttgggttg	tcacaactca	ggtatgcacc	catgtggcag	480
ccaggcagcg	gagccctact	agactgccag	tctcatgcac	ccagacctgt	ggccagattg	540
tggcctctgg	ctgcctcagg	cattttcctc	tttgcatctg	gttttctctg	attagtaact	600
acagctgcaa	cagacatcct	ccacattgtg	cacactggtt	ttgcctttgt	cctcaagagt	660
tgatacttgg	cattagcatg	aaacttgtgg	gtgtgggagg	atggagagag	aactctaaca	720
caaaacatct	tattaaaatt	gtacttgaga	gacaaaaaaa	aaaactctgt	tgtattttga	780
cagaattatt	tttattaaaa	tacacatcca	tgagcaaaaa	aaaaaaaaaa	aaaa	834

<210> 13

<211> 3371

<212> DNA

<213> Homo sapiens

<400> 13

aattcggcag	agtgacagga	acaaccgggg	tacagagcca	tttktttactg	gagtacgggg	60
ccagggtgcc	cccttagtca	ctaccaactt	cctgggtgaa	gaccaaggga	atgcaagtcc	120
ccgatacatc	cgatgtacat	cctataatat	cccttgccca	tctgacatgg	ctaagcaggc	180
tcagggtgcc	ctggcagcag	tcatacaaac	gctggcaagg	ctgccccccag	aggaggcttc	240
accgtatggt	gtggaccatg	gggaatctgg	ccctttgcgc	tgcaaccgct	gcaaagcata	300
catgtgtccc	ttcatgcagt	tcattgaagg	agggaggcgt	ttccagtgtc	gtttttgcag	360
ctgtatcaat	gatgttcccc	cccagtatct	tcagcacctg	gatcataccg	gcaaaccgtg	420
ggatgcttat	gaccgcccct	agctatccct	gggctcttat	gaattcttgg	ccactgtaga	480
ttactgcaag	aacaataaag	tccccagccc	tcttgccctt	atcttcatga	ttgacgtctc	540
ctacaatgcc	atcaggactg	gtcttggttag	gctcctctgt	gaggagctca	agtcactgtt	600
agacttttcta	cctagggagg	gtggggcaga	agagtcagca	atccgcgttg	gctttgtcac	660
ctacaataag	gtgctccact	tctataatgt	gaagagctca	ttggcccagc	cacagatgat	720
ggttgtgtct	gatgtggctg	acatgtttgt	gccactgctg	gatggcttcc	tggtaaacgt	780
caatgagtct	cgggcagtta	tcaccagctt	attggatcag	attccagaaa	tgtttgcaga	840
cacaagggaa	acagagacag	tatttgtagc	agttatccag	gctggaatgg	aggctctgaa	900
ggctgctgag	tgtgcaggga	agctctttct	attccataca	tccctgcccc	ttgcagaggc	960
cccagggaaa	ctgaagaaca	gagatgacag	gaagctgac	aatacagaca	aggagaagac	1020
tctgtttccag	cctcagacag	gtgcctatca	gaccctggcc	aaagagtgtg	tggcccaagg	1080
ctgctgtgta	gatctctttc	tcttccctaa	ccagtatgtg	gatgtggcca	cactctctgt	1140
tgtgccccag	ctcactgggtg	gctctgtcta	caaatatgct	tccttttcagg	tggagaacga	1200
ccaggagcgg	ttcctgagtg	acctgcgtcg	tgatgtccag	aagggttgtg	gctttgatgc	1260
tgtgatgcgg	gtccggacaa	gcaactggat	ccgtgctgta	gatttctttg	gagctttcta	1320
catgagcaac	acgacagatg	tggagctggc	tgggctagat	ggggacaaaa	cagtgactgt	1380

ggagttcaag	catgacgac	ggctcaatga	agagagcgga	ctctyctgca	gtgtgcccctg	1440
ctttayacca	gctgtgcagg	gcagcgctgg	ctccgcatcc	ataatctggc	cctgaactgc	1500
tgcacccagc	tggctgatct	atatcgaaac	tgtgagactg	acacgctcat	caactacatg	1560
gccaaagtgtg	catatcgggg	agtcctgaat	agccctgtga	aggctgttcg	tgacacgctc	1620
atcacccagt	gtgccagat	cctggcctgt	tacagaaaga	actgtgctag	ccccttcctg	1680
agtgcacgaa	gctactccca	gtttacctga	actgtgtgtt	gaagagtgat	gtcctgcagc	1740
ctggagctga	agtcactact	gatgaccgtg	cctatgtccg	acagctagtt	acctccatgg	1800
atgtgactga	gaccaatgtc	ttcttctacc	ctcggctctt	acctttgaca	aagtctccccg	1860
ttgagagtac	taccgaacca	ccagcagttc	gagcctctga	agagcgtcta	agcaatgggg	1920
atatatatatt	actggagaat	gggtcaacc	tcttcctctg	ggtgggagca	agcgtccaac	1980
agggtgttgt	ccagagcctt	ttcagcgtct	cctccttcag	tcagatcacc	agtggtttga	2040
gtgttctgcc	agttctggat	aatccactgt	ccaagaaggt	tcgaggcctc	attgatagct	2100
tacgggcaca	gagatcccgg	tacatgaagc	ttaccgtggt	gaaacaggaa	gacaagatgg	2160
agatgctgtt	caagcacttc	ctgggtggaag	acaagagtct	gagtggggga	gcatcttatg	2220
tggactttct	ctgtcatatg	cacaaggaga	ttcggcagct	actgagctaa	agcaagtggg	2280
taaatggcat	agggcccagg	ctagcttcca	gaaagcaccc	caggatgtca	gagaaattgg	2340
gacagtaaca	tatcttatgt	aaagtgcact	cagtctctct	ggggggaggg	ggagatataa	2400
ggagacacct	tctttctggg	ctcaagtatc	ctgccactct	gtcatgtcct	gctgatggaa	2460
ggtgcccctg	ttccctcatt	ctaccctctt	tttctctgta	atcctgtcat	aatgaatgta	2520
gcttctcagt	tcactgtata	tgattcggtg	ttgggggttt	ggaggcaccc	agaccctggc	2580
aatattatgt	gtccctttgg	accagtctcc	caagaggaga	ggggcaggca	ggaaagagtg	2640
gggatcctaa	ggttactaca	gggggctcag	tgtcatccac	aacttcctat	attagggata	2700
aaacatatag	gtgcacaaga	gctgggggtat	agcccatagg	tgggtggagag	aaaagtggct	2760
agtccttctt	gggcctggag	gttagcagtc	aagtttctct	gctttcactg	ctcgtctcgt	2820
ctctcctgca	atgattgatg	atcactccgt	ggatagagag	gcacactgtc	agaggtgacc	2880
ggagaactga	gttgcaaaat	atattaagat	ctggtagagg	taccagcttc	ctttccagct	2940
ggagaggccc	caacactgga	tggttctgta	gggagcctag	ggagcctggt	catcaacttg	3000
caatacctca	cagagccagt	tcacatccca	ctctgagctc	ccacgagaaa	cactgcttct	3060
ccaggcccgg	ggttggtggg	gagagaggca	gaggcagctg	gagcgccgtt	ctctcctgct	3120
gggacaccgc	ttgggctttg	gtattgactg	agtggctgac	agttatcttc	caaccccaac	3180
tggcttgggg	gcaggacaag	ggctaggctt	gatgggtggc	aggcttgcc	gctccccacc	3240
tgggatgccc	ctgctctgga	cctctcattt	ctcttcattg	gtttattttt	caatgcacat	3300
ttaattttgta	aagaaataaa	ataaattaag	atgtaaaaaa	aaaaaaaaaa	aaaaaaaaaa	3360
aaaaaaaaaa	a					3371

&lt;210&gt; 14

&lt;211&gt; 1094

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

ggcacgaggt	tcttcctact	ttcaaatttt	aattatttcta	gtttcttttta	aagactcgtt	60
ctctgaaatg	ttttatattt	tgtttatattg	tttaggttcc	agagtcttaa	atctggaaaag	120
atctacctcc	atagagacgt	acggctcctg	ttctctagaa	agtcaatgga	ggttgatagc	180
ggtgctgcat	atgaactcaa	atcttacact	gaatcaccaa	caaaccctca	gttttcaacca	240
agatggtgat	aaggagtgat	gatttaaagt	atttactcag	tacccaagtt	tgcaagtaaa	300
aattagcata	gaatggagtg	taccaaatta	acaatcagga	gagtggattc	tctcctgtta	360
tcctggacca	gtttttatga	aaggattcct	gaaatgaaat	ccatatattc	catgtagact	420
ggaaaaactc	atgtcctaata	cctttttgta	ctggtgaaac	cacttcattg	gacatgttgc	480
aatagcaaaa	ccccagttta	gatttagtgtt	tacacatttt	ctcagttatt	taatatattaa	540
tgtttttctt	aatactcaag	tgatgtttgt	ctctagtgtt	ctaagttagc	acaaatccta	600
tgtaaaatca	tactatgtat	ttttgacatt	aatgttgaaa	tcaaatatat	gcacaagtct	660
ttaattttgt	gtaatgtgtt	aagtgtgtgt	cattttaagtt	attgaaaatg	agaataaaaat	720
gttgagcttc	tttaaaagta	acacactatg	caagcatgtg	tactttttat	atctctcatg	780

tttagttttt	ataacacccat	atccagggtg	ctatctcaca	tagtagtcct	ttaacatatt	840
gtatttagcag	tgcaatgtgg	actaagctgc	ttcactttcc	ctttgcaagt	tcagatcatc	900
atgcccattc	atagccagga	ttccttatcc	ccaaaacagt	tctatttttc	cttaatcact	960
actatagagt	ctttacatta	aattactgtc	gtatgctaga	taattttctc	aaattgttaa	1020
aagaatatgt	actttggaaa	caaattagta	tttatattgt	aaatatattc	aaaaaaaaaa	1080
aaaaaaaaaa	aaaa					1094

&lt;210&gt; 15

&lt;211&gt; 1243

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

ggcacgagag	aacgaggaaa	aaaaggaagg	gagaggaaaa	gaaaaaaacc	taataaagga	60
gaaagtaaag	aagcaatacc	tgacagcaaa	agtctggaat	ccagcaaaga	aatcccagag	120
caacgagaaa	acaaacagca	gcagaagaag	cgaaaagtcc	aagataaaca	gaaatcggta	180
tcagtcagca	ctgtacacta	gagggttcca	tgagattatt	gtagactcat	gatgctgcta	240
tctcaaccag	atgcccagga	caggtgctct	agccattagg	accacaaatg	gacatgtcag	300
ttattgctct	gtctaaacaa	cattcccagt	agttgctata	ttcttcatac	aagcatagtt	360
aacaacaaag	agccaaaaga	tcaaagaagg	gatactttca	gatggttgtc	ttgtgtgctt	420
ctctgcattt	ttaaaagaca	agacattctt	gtacatatta	tcaataggct	ataagatgta	480
acaacgaaat	gatgacatct	ggagaagaac	atctttccta	taaaaatgtg	ttttcaagct	540
gttggttttaa	gaacaaaaga	tagttctgca	aattcaaaga	tacagtatcc	cttcaaaaaca	600
aataggagtt	cagggaagag	aaacatcctt	caaaggacag	tggtgttttg	accgggagat	660
ctagagagtg	ctcagaatta	gggcctggca	tttggaatca	caggatttat	catcacagaa	720
acaactgttt	taagattagt	tccatcactc	tcatcctgta	ttttttaaga	aacacaagag	780
tgcataccag	aattgaatat	accatatggg	attggagaaa	gacaatgtgg	aagaatcata	840
gagctggaga	ctacttttgt	gctttacaaa	actgtgaagg	attgtgggtc	acctggaaca	900
ggtctccaat	ctatgttagc	actatgtggc	tcagcctctg	ttaccccttg	gattatatat	960
caacctgtaa	acatgtgcct	gtaacttact	tccaaaaaca	aaatcatact	tattagaaga	1020
aaattctgat	tttatagaaa	aaaaatagag	caaggagaaat	ataacatgtt	tgcaaagtca	1080
tgtgtttttct	ttctcaatga	gggaaaaaca	attttattac	ctgcttaatg	gtccacctgg	1140
aactaaaagg	gatactattt	tctaacaagg	tatatctagt	aggggagaaa	gccaccacaa	1200
taaatatatt	tgtaatatgt	ttttcaaaaa	aaaaaaaaaa	aaa		1243

&lt;210&gt; 16

&lt;211&gt; 818

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

gattttctta	cctcaaagct	gaagtttttc	ttcctattat	aattccactc	tctgttataa	60
ctgactcttt	tctattttctg	acccaaagcc	taaaactcac	atcaaaatat	gctgtgaaga	120
aaagtgataa	gtttttcttaa	aaatcattaa	atgctacctc	ataaaataaa	tatatatatg	180
aaacttgaag	catttataga	tttctgttgt	tttctttagt	tgtaaacctg	gatacagggtg	240
gttttgatac	atcttttctg	tttgaagtag	caggcttccc	ctgcagtgtg	ggtgatggcc	300
acctggtggc	acatcacact	aatgcatgga	gcttctcatt	tgtcttcagg	attctgaaag	360
taaagccaat	gtttggatgt	tatgggagct	atgcatttca	ttgtattgtt	tcttgtaaat	420
agtttttagaa	aatactcttc	acattatagt	tcagccatca	aaaaaaattc	accagaaat	480
gaaggtagta	gtattctcca	tattgcagaa	atcgtagtta	gaatttgctt	taataagcag	540
tgagttatag	agaacaggat	tatgtagtta	gaaaagctaa	gaaaaatggg	aaacgaaaag	600
agttaaaggt	ggacttaatg	gacatatggt	atgggcataat	atatgtattt	ttaaatagta	660
ttgctgctgt	tttaactatt	ttaataaata	tattacaaaa	tttatggcaa	tttttaaatc	720

tttagaatta ttacctgcga atatctctta atgaatttcg ttagcttctc tttgaaaatg	780
gctttttgtg gtttttctta aaaaaaaaaa aaaaaaaa	818

&lt;210&gt; 17

&lt;211&gt; 847

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

tttttaacaa agtgggtccaa gggaaaaata atgtttttaag tttttatgtg cggattccag	60
acaagcattc tttctttttg ttcttctctgc actctcctac atttctatgt tggtactaga	120
cagtgatgtc tggttctgac ttattttatgt atttttttttc tatttgtttc tggcaatgca	180
cataataaga aatacacata gcaatattct tgctgttggt gttcataacc actgagaaaa	240
aaaagcagca acaagtagga ataaaagaag cctcagccac agaattggcaa gaaacgttat	300
atgtttttta gggccactct ttgcactcgt ttacagaca gggtactact tatcagctta	360
ctgaccttgg gaagacccaa cctctgagct amcamtityct catctgtaac ctraggatcc	420
agtgaagtca tgtaagacaa tgtagcattg tacgcttatg actagacccc agtcatgaca	480
tgagctgctt ctgtctagaa aatagagggc tggcagggca cgttggttca cgactgtaat	540
ctcagcactt tgggaggccg aggcaggcgg atcatgaggt caggagtcca agaccggcct	600
ggccgacatg gtgaaccctg tctctactaa aatacaaaaw ttagctgtgt gtgatgggtg	660
gcgcctgtaa tcccagctac tgaggaggct gagccaggag aatcgcttga acctaggagg	720
cggagattgc agtgagccaa gatcgcatca ctgcgctcca gcctggagac agagcgagac	780
tccatctaaa aaaaaaaaaa agaaaaaaga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	840
actcgag	847

&lt;210&gt; 18

&lt;211&gt; 662

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

ggcacgagaa agccagctag ctttaagata cagtgttggg catcacattt tgacacagag	60
caggcactct tttcttttgcc cccaggtggg actaagccac cacaagcctt tcctgggtgtg	120
tgcagtgggt gatgaatgct tgccctgctca gcacccactc cctgggtggac tgggtcacat	180
tactctgact cccctctgag cttcagtcga cgtctgggtc aagatgtatt gactcaaccc	240
gaggatccag aggtgggatg tagctctggc ctggccagag gactgaggat gctgaatgcc	300
atggctacag caactggtca gctttgggct catgtcctag tcagagccaa tgagatgtaa	360
tggtgggata tctgctgggc tggtgggaag gggaaaggct gcctgctca tccccattcc	420
tgatgctgag ggatctgaga aaattacttg taaaatttgg ggggtgtctg aaaaagggga	480
gactgatgtc tctttctctc tacagacatc tgatcagcta cagagcttga ctaacctacc	540
cagaggcaga atgatatggt gggttaaagt gggctctggg acgagttttt gccactgtac	600
tccagcctgg acgacagagt gagactccat ctcaaaaaaa aaaaaaaaaa aaaaaactcg	660
ag	662

&lt;210&gt; 19

&lt;211&gt; 2254

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

ctggttctta aaatgccctt tttctcctgc ttttaggagc tctgtgatca tatcgagctc	60
atctgggtta atttatttat attcacaata tccatggagt acatatggca tcttgtgggc	120

cttttagtctg	ccttctatga	atataagaca	ttatttcccc	agaatgaaaa	tgtatatgaa	180
gaaagagaaa	tatgccactg	gaaagtagtt	caaataatga	aagtcctttg	cttggttatag	240
ccttcctttt	tttctgttct	ttttcctttc	cttcataaaa	agatttatta	tatgctttta	300
ggaaaaataac	aaaagcagaa	tgtttgctaa	ccatctatgg	gtgaaagagt	agcaaatttg	360
gtttaaggag	atgttgtaaa	taatggtatg	acaattcctg	ggattctggg	agtcacttaa	420
taaggaaaaca	gaaatagatt	ccaatgccag	tgtgccccct	aatgagaaaa	gggaattcct	480
tttactgagg	cctataatct	tgggaatatg	ctttttaatt	aaaattataa	gtctaaaact	540
tttactatta	ctatgactgt	ttatagctcc	caatttaact	cacgtaacag	aggagtgagt	600
ggatgggtgg	gctagactta	agataataag	ggagtaagga	gtgaaagaaa	acccaccta	660
tctcctcacc	tgaaggccct	ggacacgtcc	agcctgcttt	gggagtaagg	caacaggaga	720
ttgaaatcaa	gttggcagtt	gagttttgaa	ctgtattaga	ctccacagtt	caattttcaa	780
gagttgtatg	aaaataaaaac	tagtaggata	aaagacagac	atccagtaca	ctcattcctt	840
cttcaatgtt	ctatttttatc	attatagcca	gtgcctgtgg	caggcacact	ctgagatggg	900
ccacagtgtt	ctgtctcccg	gaatccatac	ccttgtgtaa	tcgttaccac	ttcagtgtag	960
gctggatcta	gtgactcttt	actgaaaaat	acaaaatggc	agaaatgctg	ggatgccact	1020
gattttgtta	ggttatcaaa	acgctgtggc	ttttgccttg	aatttctctc	catgctcatt	1080
ttcttacttt	ctgtaatacc	ttacaaaagg	agaagcaagc	tggcataatg	agagcaactt	1140
tatggagaag	tccgtgtgag	aaggaattgg	tatctctagc	caacagctag	ggaggggtctg	1200
agcctgtcag	aaaccaatga	atgagtttgg	aagcagatct	gtcagcccct	ttcaagactt	1260
cagatggctg	aaactccaac	tgacaccttg	tctggaatct	cattaagcca	tgaagcacgt	1320
agctaaacac	attcttttacc	aatatgaact	attaagtaat	aataatctat	tgttttaatc	1380
catgttttca	ggttattata	gcaatagata	acaaatatgc	atagtgtcct	agtttaaaag	1440
ggattatatg	ccattaaatg	cacattgcaa	atatgattat	atggattcat	ttttccttat	1500
aggcaatatt	ataagcaaaa	gtcagattcc	tggacagcct	agcagaatgg	gattactttt	1560
ataaaaagat	gaccaaagtg	tttgcataaa	atattcttca	catttcataa	caccaaagca	1620
aaagagtact	tcacagctca	ggggagattg	taaaatactc	agatcgctta	cagaaccatg	1680
gggtcgtttc	atctctttga	cttgtggtat	ttatttccac	caagtctctt	ttaataaagt	1740
ctgcggaagt	ttcccgggta	agttcaagggt	gttgttcttt	ctgtagtgcg	tatttcaaaa	1800
aaaaaaaaaa	aaaaaaaaagc	tgacattcta	agcctttcaa	aagaattctt	cttctctttt	1860
tttgttttat	agagataaga	aaaatcaaat	attgtatttc	aattaaagat	acaaatcatt	1920
taaaagatca	gccattttatt	aatgaacaaa	tacgttgctc	aagaggtgac	aagcaatggt	1980
ttcagtgtat	ttatttttatt	ttcacttcaa	attgcttatt	ccagctctaa	ccacctacat	2040
tattattatg	agctgctgtt	aactgaacct	ctatttttatt	cctctttgtc	caagagctct	2100
acaaacactc	tctataatcc	ttaaaataac	tccacaaatg	gaagacatta	tttccatttc	2160
acagatgagg	aaacaaaactc	ggagttatca	ggtagttgtc	tgcctatgaa	atggaagaaa	2220
aggcatttga	attcaagttt	acctagtcct	aaaa			2254

&lt;210&gt; 20

&lt;211&gt; 1079

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

ggcacgagta	agtccttgca	caagtgggat	atatttttgaa	tattgaccat	ctagcttgat	60
gtctcataca	aaaagatat	ttttacacag	tttttaatat	ttccatagtt	tattttcaaa	120
gggactgtaa	tgcataattac	ttattaaaag	cattttttcc	cccaactttg	atggggaaaa	180
gcatatcaac	agatacctac	ttttgtagat	ggctataaaa	attgttacc	aaaaggcttt	240
cttagagcaa	attgtctctt	atgattatc	ctgcctgcct	ccatgggatc	tgtctttata	300
ttgttgagga	gaatagcaaa	catccataat	acattttttt	ttttttgtgg	atataacatt	360
caggatagaa	ttgtttgtca	tttattgctt	agattgatat	tttgtatact	tagggattat	420
gaaaaaagtt	tgtgttaata	ttgcattatt	aaagcacatg	tttaaagata	attattagat	480
tttaagttca	caatgcttca	tgtattgggt	tattctgggtg	attttatttt	tacttttagga	540
atttattctt	cattgcctgt	tgaagtcctt	ttgaatcaca	aacggtttgt	ttgtgatcta	600
actcaagctg	atcgtcttgc	cctctatgat	ttcgtagttg	aggagacaaa	gaaaaagcgc	660

tctgattctc	raattattga	aatgacagc	gatctctttg	tagacttggc	tgccaaaatc	720
aatcaagaca	atagtcgaaa	aagtccaaaa	tcctaccttg	aatccttggc	agaagtacga	780
gattataaaa	gaagacgcca	gtcctataga	gccagaatg	ttcacataac	caagaaatca	840
tatactgagg	tgattcgaga	tgtgataaat	gtgcacatgg	aagaactcag	caatcattgg	900
caagaagagc	aagagaaggc	agaggatgat	gccgaaaaga	atgaagaaag	gcgatcagct	960
tcagtagatt	cacggcagtc	tggtggaagc	tatttgatg	ctgagtgttc	acgacataga	1020
agggatagga	gtagaagccc	acataaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1079

&lt;210&gt; 21

&lt;211&gt; 632

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 21

ggcacgagtt	actcttcagt	tctcaccatt	tttacccttc	tgcaaagtct	cttgtaattc	60
ctaagtaatg	aaatgaaaag	tacaaatttc	ttaaaacaag	ctctgttctt	tttcttctgg	120
aaaacttggt	tagtttgctc	tgtgtatctg	tttctcatga	ggagaccggc	tttctgtggc	180
ccacgtgaac	actgagtaag	aaacaaaaga	ctgtggtctc	caggacacag	tgtgtgtttg	240
tcctctgcca	tggttattca	ccaagtggag	tccagcagtt	taggaatcgg	gaggtctccc	300
atgatgagtt	gtcattcttc	gaattgctgc	aagtgcacac	aaaggggccc	ccctaccagt	360
ttctcacttc	ccagtctcac	tactggatca	gctcttagga	gccaggagag	ttcactgctg	420
tggttaggat	agaaaagggc	agctagtgcc	ccagggtaga	tcttggaata	tatttttttg	480
gaaaaaatgt	attaaggcca	cccctaaaat	agatactgta	tctggctgta	ctataactaac	540
agtgatttgc	ctgcatgtgt	ttgatagaga	tttctaccat	gtactgcttg	gtgctggata	600
gtctatcaca	gcaaaaaaaaa	aaaaaaaaaa	aa			632

&lt;210&gt; 22

&lt;211&gt; 1492

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 22

agtttctaaa	actctgtcat	ttcaataatg	ttataaattg	aatcatcacag	tatgtaattt	60
tatgggatcc	gctttttttc	tggtgatctc	tttttttctg	agtcattctca	gctgttgtgt	120
gtatcaatag	ttcatttcct	tttactgttt	tggttggttt	aactcattga	cctattgaag	180
gatatctggg	ctcattccag	tctgtgaata	atgtgctgtg	ggacattcat	gtgcaagttt	240
ttgtgtgaac	attaagtttt	catttcttta	agataaatgc	ccaggagtga	aactgatggg	300
ttgtatgtag	ttacaaactg	ccaagctgtt	ttccagagtt	ggcgggtacca	ttctaatttc	360
ccatgacaat	tttttgaaga	gtccaagcta	gttgctttag	agtgcctcac	ctacattttt	420
taattgtctg	actttttcct	aatgacatca	cttaatttgt	tcctcaatct	cctgaatttc	480
ctatagactg	gagggttaggt	ctgggtggtt	gattagattt	tttaaaaaaa	tatgacttca	540
tctgtgatgc	tgtgtacttt	atgttgcata	acatgaaccc	taagaacaga	gtgagctgct	600
ggacagcaag	tttcatgggg	tgcagtaatt	aacacaccac	atagtataaa	tctgaaataa	660
tgacaaatgt	gttaagggcc	ttgggatatt	gggccatgta	ctctgaggag	acaaggtgag	720
gtgcagggtc	ctgcccctaa	agaactctat	cttttgagat	tagcaactaa	cagtgtgagc	780
ccactaatag	gatgtgaaag	ttgtcaaaat	caagttctgg	tcatttgtgt	aaaaatccta	840
acaaatagag	ctggggaagg	ccgtgaaagg	acgattttca	tgcacagatg	tctgataatg	900
aggactatca	ttaaaagact	gcacaaaacc	acaccttgca	caaaggccat	cacaacctga	960
cacacacaaa	aaataacttc	atgaggacat	ttgccagca	actccctgtc	caatgtccaa	1020
ctggcaacat	ccttgttatt	gataccttgta	gccaaaggata	attctctcaa	aacaatcatt	1080
tttgctttta	aaaccgttgt	cttcccttgac	ctccctgtat	atgcacatag	tttactgtgg	1140
cacttgtatt	cttattgcaa	tgcctactcc	tgaataaaca	tcattttctt	tcagagagtc	1200
tccctctctg	ttatttaggc	tgacaaggat	atgccaagaa	gtagcttggg	tatagcagtt	1260

aactctgcct	ttaggatatg	tgtatgggga	tataagagtt	aactaaaagc	tgacctttga	1320
gttggtcctt	gaataaaaga	agaatagatt	ccaaaaataa	gaggaataaa	taatgatcct	1380
gagctaggag	agtacatttt	gcagaccttt	ggtttccata	ttaagaaatt	cagattttta	1440
tgtacaataa	agaagttctg	gaattctcaa	aaaaaaaaaa	aaaaaactcg	ag	1492

<210> 23  
 <211> 638  
 <212> DNA  
 <213> Homo sapiens

<400> 23	
ggcacgaggg	accccaaaac ccaatgatcc tgcagcagcc cttgcagcga ggccccccagg 60
gaggggcccc	gcgcctcccc cgggccgcct tgggggtgac ttggggcctg gacgccaggg 120
agcctctgcg	caagcagttt ctgtctgagg agaacatggc caccacttc tctcaactca 180
gcctgcacaa	tgaccacccc tactgcagcc ccccatgac cttctcccca gccctgcccc 240
cactcaggag	cccttgctct gagctgcttc tctggcgcta tctggcagc ctcacctctg 300
aggccctccg	tctgctgagg ctgggggaca cccccagtc cccctaccct gcaaccccag 360
ctggggacat	aatggagctc tgagtgtctg tggacagtgc cctccccacc ttccttcttc 420
cccacaacag	aagagaccag cgactccgc aaagggacaa ggttcctccc tctcctgcag 480
agtaggcata	tgggcaccaa gaccttcctt caacagagga cactgagccc aacggagttc 540
tgggatggga	gggggtgggtg catgggaagg gaggcattccc acccccaaga agaactgaat 600
aaagattgct	gagcaaaaaa aaaaaaaaaa aaaaaaaaaa 638

<210> 24  
 <211> 944  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (864)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (939)  
 <223> n equals a,t,g, or c

<220>  
 <221> SITE  
 <222> (944)  
 <223> n equals a,t,g, or c

<400> 24	
ctttcgcgtg	ggcggacgcg tgggcggacg cgtgggcgga crsgtrgggg aaaggggact 60
tgtggtagtg	gaccatacct gggaaccaaa agagacccac tgtaattgat gcattgtggc 120
ccctgatctt	ccctgtctca cacttctttt ctcccatccc gggttgcaatc tctactcagac 180
atcacagtac	caccccaggg gtggcagtag acaacaaccc agaaatttag acagggatct 240
cttacctttg	gaaaataggg gttaggcatg aagggtggtg tgattaagaa gatggttttg 300
ttattaaata	gcattaaact ggaattgaca agagtgttga gcatccctgt ctaacctgct 360
ctttctcttt	gggtgccccctt atctcacccc ttccttgga ttttaataagt ctcaggcatt 420
tccaattgta	gactaaaacc actcttagca tctcctctag tattttccat gtatcaggam 480
agaggtgtct	tatgtaggga gggggcaagt atgaagtaag gtaattatat actactctca 540

ttcaggattc	ttgctcccat	gctgctgtcc	cttcaggctc	acatgcacag	gaatgctaca	600
tgatggccag	ctgcttcctt	ccttggttat	catccactgc	agctgctagt	tagaaaggtt	660
tggagggatg	acttttagta	aatcatgggg	attttattga	tttattttca	cttttgggat	720
tttgtggggt	gggagtgggg	agcaggaatt	gcactcagac	atgacatttc	aattcatctc	780
tgctaataa	aagggttctt	tctcttgggg	gaaatgtgtg	tgtcagttct	gtcagctgca	840
agttcttgta	taatgaagtc	aatnccatca	ggccaaggaa	ataaaaataat	tgcttacctt	900
aaaaaaaaaa	aaaaaaaaaa	aaaaaaactc	gaggggggnc	ccgn		944

&lt;210&gt; 25

&lt;211&gt; 408

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

ggcacgagtg	aatcttttct	tttttccccc	ctcctccaaa	tatataaatc	tgtgctctca	60
catgtaatgt	atttaatttt	tggaaaattt	aatgctatat	agtaaatcaa	gtgtgtgatt	120
taaagtaatc	agatcacttt	gattctttta	agttttcaga	atttcttttt	tccacaggta	180
attttggaga	aaataattac	caattaactg	ataataagca	cgctgtcctc	tgctgtaaaa	240
agtctgaata	gatactgtgt	atgtttttat	gtccatgaac	ttgagctact	cgtgtgcaat	300
tatgtgtatg	ggaatggagt	agtgttcatt	atgtgtatct	acatcatcat	atggatgtat	360
atttattaat	aaagtcatta	ctttaaacc	aaaaaaaaaa	aaaaaaaa		408

&lt;210&gt; 26

&lt;211&gt; 1687

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

ggcacgaggt	ttcatgggga	cagagtgggc	agatggagag	ttctgtggag	ggagggaggt	60
gatggtcccg	cattgctgtg	actgcttaat	gccagtgaac	tgcgcacctg	caatggtcaa	120
tctctgccct	ccagcgggtg	cccccatggg	gccaggcccc	cgatcatctt	gggtgcacag	180
cgcacgatgt	ctcaccagga	tgggatccgc	tgagggttgg	agcctcagct	caggccaggg	240
ctgcacctgc	ccccaccaca	ccatacacac	cttccccact	ctgttcccca	ttggccccct	300
cctcactcca	cccatgctgt	ctggccccct	ctccccccagg	ttctgcaaga	gcgggtgtgg	360
tgggccccact	tgcagtccca	gctcctgggg	aaattgcagg	ggctgctgga	accaggaaa	420
tcccagctcg	ctgggcatct	tgaactacca	aaaaacaccc	tcagcagaaa	atgagagggc	480
aggaaggatg	tcagcagagc	ctctgtggac	tcttcatggg	aggagactct	catgctgact	540
tgccaggagg	caccccgga	gccccagtc	caggagtga	atgggaaggg	tagttcagag	600
gcaggtttct	gacccctctg	gcctccatca	gagccaccgc	aaaaattcca	cctgttaaat	660
aaaaacggac	tgaggccgtg	gtgcactcct	gtaatcccag	ccctttggga	gccaaggtgg	720
gaggatcgct	tgaaccagag	agctcaagac	cagcttgggc	aacatagcaa	gactccacct	780
ctacaaaaaa	attagccggg	tgtggtggcg	tgtgcccttg	gtcccagcta	cccaggaggg	840
cgaggcgggg	ggattgcttt	ggccccactg	aatgttctgc	caggaggggg	tagtttagag	900
ccatgaacc	acggcaatga	gcactgcgag	cacggacaat	ggggggctgg	ggttgggggtg	960
ggctccacaa	cgctgaaagc	tgagctatcc	caggacctgg	gaggtgacct	tgtggctgcc	1020
gagttttctt	ctggatgcac	accatgtcta	tcaaccctaa	ggaagggtcc	ccactttgct	1080
gggagcctcg	caggctgggg	gcacatcccc	aatccctcgg	ggcaggctca	gatcctggca	1140
cactgacatt	gccagccact	cctgggcact	tgtgtgtgtc	cgacccagga	ctaggagctg	1200
cacctgggcc	ccaggatgtc	ccctagtgtg	caaagtgtga	aacttgagtc	tcagagactt	1260
agggacattg	gcctctgtga	atctcagggc	tgttctctct	gaggcctgaa	ccaggcctgg	1320
cactgtctgc	tgggtgcggg	cccagccctg	ggaggaccca	gcaggggtgc	tcagacctcc	1380
catcccatgg	ctaaccacc	gctggggagg	aaacctgtgg	acacttgcac	tgaataatgt	1440
cactctctac	aagtatgcaa	gggagtcgga	ccatctccca	tcagacctgg	acaccgcgcg	1500



gacaccacag	cccctaacac	tccggccagg	tcgctccagg	ctggcacagg	tggcaccag	1560
ccagtcacac	ttaaagcagg	gtttccctgg	ccttaccata	ttatcactcc	ccaaggaagg	1620
aaatgcgaat	ttttaattaa	gtgaattaaa	tgctaattta	attaagtga	aaaaaaaaaa	1680
aaaaaaa						1687

&lt;210&gt; 27

&lt;211&gt; 1189

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

ggcacgagat	tgtttttccc	accttgtcgc	tgaacctgtc	tcctcccagg	tttcttctct	60
ggagaagttt	ttgtaaacca	aacagacaag	caggcaggca	gcctgagagc	tggcccaggg	120
gtcccctggc	aggggaaact	ctgggtgccg	ggagggcacg	aggctctaga	aatgcccttc	180
actttctcct	ggtgtttttc	tctctggacc	cttctgaatc	atagaccgga	caagagcctg	240
cagcggaaag	gactctgggc	tgtgcctgaa	gctggctggg	ggcaggacaa	cacagctgct	300
tccccaggct	gccactctg	gggaccgctg	ggggctggca	aaggcatcgg	tcagcggggc	360
agcggggctg	gccatgaggg	tccaccttca	gccctttggc	ttcaaggatg	gagatggttt	420
tgccctccct	ctctgccctc	gggtggggct	ggtgggtctg	cagctgggtg	gggaacttcc	480
ccacggatgg	cgggtggagg	ggttcgcacc	gtgctgggct	ccccctgact	gtagcacgga	540
gtgttggggc	tgggcgccat	ctccaggagg	gcttgagagc	tcagcctgcc	tgggagagcc	600
cttgtggcga	ggcattaaaa	cttgggcacc	agcttctttc	tcggtggcag	aaattttgaa	660
gtcagagaag	aaacggctct	ttgttggctt	ctttgtcttc	tcgtgggtcc	tttggcaggc	720
ctcccttttg	ggagagggag	gggagagacc	acagccgggt	gtgtgtctgc	agcaccgtgg	780
gccctcaagc	tttctctgct	tcttctccct	cctcctcctt	tcccctttct	ctttcctcat	840
ttcctagacg	tacgtcaact	gtatgtacat	accggggctc	ctctcctaac	atatatgtat	900
atacacatcc	atatacatat	attgtgtggt	ttcccctttc	tttccctttt	ttaagcaaca	960
aaactatgga	aataataccc	caacagatga	gcgaaaatgt	attattgtaa	agttttattt	1020
ttttaatact	gttgtctata	atggggaaaa	aggacattgg	ccccgcagtg	ccctgccccca	1080
gtcagcctgg	ctgggctctg	gtgggggctc	ctgatccgca	tccaagctta	accaaggctc	1140
caataaacgt	gctaggaagc	aagcaaaaaa	aaaaaaaaaa	aaaaaaaaaa		1189

&lt;210&gt; 28

&lt;211&gt; 1820

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

ggcacgaggt	caatggcttg	gacagacaga	cgggctcagt	ggcatttggg	accctctttg	60
gtgccctccc	attctctctg	gaattgtttc	aagtctgctg	gttttcaaac	aagaaaagac	120
ctttctggcc	atagggagaa	tagcaggagg	tctatgtttt	ggtgggttaca	ttggaaacat	180
cttaagcaag	agagggaaag	ttgatttttag	gcacacatgt	accctccttg	acagcaggaa	240
ctcagacttc	aatcttgggg	gtctaagacc	agaatatatt	ccttctgcca	gaaaagaatc	300
ttgcacatat	actcctgagg	catgagtgtg	tgggtccatg	caagaaatag	ctaaaggctg	360
ctttccagg	cccaaagccc	catttaaatg	aagaaccaga	gaagtgttct	aggccattag	420
tggacaatgt	catgttttgg	gaaagataac	aacacaaata	atgtaacctt	tccttaaaag	480
gcagaactca	atccatttta	tttgatgctt	attctaacc	taaccctggg	tcacctggaa	540
tgaagaactc	tatgaataat	atttgatttt	acaacgtgtt	atgggttatgt	gaaaactaaa	600
catttgcctt	ttataaagac	tgacaaaata	taaatcttta	ttctaaccct	atccccaaaa	660
ctagccaggc	cacaccccag	atgttcttat	tgactattgg	gaagatagaa	aaggcggtgt	720
gttttttgtt	tttttgttgt	tggtgtcatt	gttgtttttt	tcagaagacc	agtgtctcag	780
ttctgtctta	gtagtaccac	acccgtaacc	gtgtttttaa	agtttgtttt	agcctagaga	840
cagatcatat	gagttcaaca	atgtacagtg	tgattgaaaa	gacaggttgg	tgtctatttt	900

tcttttttaa	atatctgaat	gtgtatttgt	aatacgtaaa	ggtaaaaaaa	aatagtgcca	960
aaaatgtgca	aggcatctca	ttacagctca	tgtacgtctg	tttttataag	atcaatatta	1020
aaacccattg	ggattaaata	tttttgaata	ggatacactc	ttgrgaaact	cgrgaatgga	1080
ctgagccttc	ctacaagcca	ctctttgttt	ttaaaacagt	ggggaaatac	gtttacagag	1140
attgtgagct	tcagagaatg	catgtgatgg	tgtgtattac	atgctaattc	atataagctg	1200
tatctgtcag	ctaccaccct	gtgcttttaa	aatgcacaca	ctcaaccctc	tttagcttgg	1260
agctcagctt	tttgcttttt	tttttttttt	tgtagaatta	tttagctaac	ataagtattc	1320
tgawtgctac	ctgatggcca	ttcttactta	gtttcataga	tgtgctttta	ctatgatcct	1380
ttgaagctca	ccccttggag	ascctacaga	acctcaggct	gatagctttg	aagactgcca	1440
aacagcccag	aaggaagcaa	agcatctgca	taatcaggag	ggttgtataa	caagtagtga	1500
tttgccaaat	atgtgggtag	ctttaggctg	aggcacgggc	ctcaggcaaa	aatgcccttc	1560
gagtgaatcc	gaagggcctg	atcttccctat	gtccttgact	aggcatgacg	agtcatttga	1620
ggtcagatat	tatttgagtt	gttcagcacc	cccaaaggta	ggcattctcc	tgggaaattt	1680
tcatttccat	tttatcgcca	aacaaaataa	aaagcaaaac	aaactttcta	agctagaata	1740
atgaaattaa	gtcatttttc	actttgtata	tattgatgct	aataaaaacag	atgaaaaara	1800
aaaaaaaaaa	aaaaactcga					1820

&lt;210&gt; 29

&lt;211&gt; 1573

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1549)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1550)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 29

gacaattcct	gcccctaaag	cctaggagat	ccctctccct	tgctagagag	ccacccccaa	60
atcaaaatgt	gaaaatccct	agaaagcaat	agccttcgag	gtaccttgca	ctgaatttcc	120
caccccagcc	cttccaccgg	atgggaggct	gtaacttggg	cactgggggtg	actttttcca	180
tgcccttgtc	atctccaggg	tgggaggcag	gccccacttc	ccctccccta	tccccactt	240
cccattgttg	ttgccccacc	cctaattctc	agactgaacc	cagatggaga	tctgagtgcc	300
aaaacaattc	ttgatgtaac	tttgtacata	tcttctacta	ccgttggggg	ctcttggggg	360
tagaggtggg	ggcggtctg	tgggccattg	ctcccttcca	cctctcaaaa	gaccttacag	420
tatttcacag	tatctctacc	cgcacgcgag	tattacagta	tctagctgga	atataccctt	480
acagcccccc	aggaccttat	gaggaaggga	aggagccagg	gagagtgaag	taaggtctgg	540
gactgggggag	gtgggatctg	aatgaactca	tttgcatatc	atttgcaccc	tccgtctggc	600
agccgctttc	tacaaactca	tttactggag	tctgggtccc	aatcagccgg	gtccaggact	660
cctctcacac	agacacatct	ccggaggctg	ggcctcctga	aaagtgtttg	cttgggggtg	720
ctgtgtaaca	acccctccct	attcatattt	cttgggggacc	ccctaccag	ccagccaggg	780
tgatctgaaa	ggtatacttt	gctagctcag	tgagctagtt	cactcaccat	gttgggtgagc	840
agagagccac	acctttcccc	attttaccyt	gggaaactca	ctccaccatc	tttgccatct	900
cttgaaagtc	ccttctgcaa	tctgacctca	atcttttgtg	ctgcagtttg	tccagagggg	960
acacagatgt	ggggtcaggg	atgaggatta	ttgraaaacc	catcatctct	ttttttttcc	1020
ccgtctccct	attagccaat	ccgatctcag	agtctctgag	tggcctcctt	gcacccttct	1080
cttcagcacc	cagtaggtgc	ttaataagtg	tttgctgcat	tgaattatct	ccctattcct	1140
tctcatttgc	cctctagctt	cccatacctt	ctccaagtgt	cttctctcct	ttctttgtct	1200
ggctccctat	gactttctat	ttttttttcc	tccgtgtggg	tcccattgtt	ttctgtcctg	1260

tctctatctt	agtctttgtc	tgtcttccctc	ctttccctcaa	atgtctcaac	tctctctccc	1320
caatttcccc	atttaaaaaa	aaaaaaaaagt	gccaaacttc	cttggaactg	agccgctctg	1380
gggggagagg	accttggata	gaggggagga	aatgggacca	tttctctttg	aggaggtccc	1440
taagaggcat	tgcaaaagtg	tgacatgga	gctaaattgg	gtcccccttc	cacagccctc	1500
ccaccctgag	tttttcttag	aatctttgta	aaaaaaaaaa	aaaaaaaaacnn	gagggggggg	1560
ccggtaacca	att					1573

&lt;210&gt; 30

&lt;211&gt; 1328

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

tcgagttttt	tttttttttt	tttgcttttt	aaatttat	tttaactgtt	ctcatatgta	60
gcaacccctc	ctccccctct	gggcatgttt	acacaggctc	tgctctgggg	gctggcctgg	120
ctgtgaggtt	tctggggagg	cagagaggca	gggacttttg	ggccttagtc	accatccatg	180
gtatcacctc	atctcacttc	ctgtgaggga	cagggcctgg	ctgatgtgat	cccagctccc	240
cccagttcag	gactgtcttt	cagctccttt	gccccctggag	gtgggggctg	ctggctgagg	300
aggggtcaag	gtgagttcaa	gaaagctacc	tgtggaaaat	ggaccaggtt	gggggggtga	360
ttgcaaagtc	tccccaaagc	ctggctcctc	atgctcagtg	ccaggggcag	aacactgggg	420
agccaggtat	agagagcctt	cctgtcataa	ctgccagtc	tcttctctca	aggcctctgc	480
atattctcat	gttccccctc	cccatcatgc	cagccacccc	tatccctctt	ctagcagggc	540
caagatgggg	acagcagcag	ccctctggcc	tgggatgtg	atgataaagc	aagcctaggg	600
ccagggtttg	gggagcagag	agagccaaga	agttgaccac	gtgtgatttc	cagcccttcc	660
cactgggact	tgacttccca	ggtcaaggag	tccgtctcat	tctggctggg	cgagtgacca	720
gaggcctgtg	tgaatgtgtg	cacctgcttt	tccctgctgg	aatgttttct	ggctcagctg	780
cagcaacatc	tgtgagccca	gtgtctgccc	tgtgtccctg	ggctcgtctc	aagtgcagga	840
acatacatgc	agggcccaac	atgatgatgg	tgtgaagggc	aggaaacagt	cctctgaagg	900
agtggggagg	tgggcagctc	gcccccgcca	ggtaccatcg	cctcctgcca	gcttccttag	960
accaggcagg	gctgccatgg	tgttagctgc	aagtccatca	gtattgaccg	tctcgtctca	1020
tcttggtcct	ccggagtccc	aagtttcctt	ttcatcaaat	ctgacaagag	agaagaaaca	1080
tgggtgtgct	tggcccacag	ggcctggtgg	tgatggacct	ccccgctccc	tcaagctctg	1140
gatggctgca	gtgttgtact	agactttgtt	caggctgttc	tcatctcagt	attgcccttt	1200
cctttcactt	tcacacttca	tctcatctct	gttgtcactt	tccccgaaac	gaataaagtc	1260
tccccagctc	ctgctgtgta	ggctgggcag	aaaccacaaa	aaaaaaaaaa	aaaaaaaaaa	1320
aaaaaaaa						1328

&lt;210&gt; 31

&lt;211&gt; 590

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (27)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (37)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (58)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 31

cctaagggga	ccaaaagctg	gagctcnacc	gcgttgncgg	ccgctctaga	actagtgnac	60
cccccgggct	gcggaattcg	gcacgagtc	aggtcttatt	cagatttcac	cagttcgtgt	120
gtgtgtgtgt	gtatgtgtgt	agttgtgcac	tttaaccacc	accataatat	acagaattgt	180
tccgttgcca	caaaggaacc	cttgactac	ccacttattg	tcacacctta	tcacccatcc	240
tttcaccacc	ttccctaacc	cctggaaatc	actaatcttc	actgtaattt	tgatcattca	300
agaagttagt	aatggcataa	tatatgtgat	acaaacatag	catatgttga	ctttttgaga	360
atgggtttcc	cctccacagt	atgatgctct	tgaattccat	ccaagtttta	ttatatgtac	420
cgatagtttt	tcgtttttta	acagattgct	gtggatgtac	cagtttgtct	gactgctcac	480
tcactctattg	taggacattg	ggttggttct	agttttgggc	tattaaaaat	aaagctgcta	540
tgaacatttg	tgtataaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaactcgag		590

&lt;210&gt; 32

&lt;211&gt; 752

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 32

ccggccgctc	tagaactagt	ggatcccccc	gggctgcagg	aattcggcac	agcttcttcc	60
atggagctag	aaagatggca	tttggcctcc	tctgcaaate	ccagctcaca	attctacctc	120
cagggaaaaga	ctgaatgtac	tgcttcagct	ccattcaaaa	acagttctgg	gaaagattcg	180
gggtggcctg	ctttgattat	atgtgcaccc	tcttctgcac	ccccatctcc	ttatttctct	240
gctggcttcc	tggaccatcg	cttgagccat	gggtggagag	actcctgaaa	gaccatcaca	300
ggcagcccta	actagactac	ctgattgaaa	taagcagcag	cttttcttgg	ccgcaaaggg	360
gaatgctgct	cacagaataa	gagagcaggg	cagagcttcc	aagtgtctac	acagcctttc	420
tttcccttat	gtaaataaat	taattcatta	ttaacatate	tcaacracac	gaatctgacc	480
tcttttaaat	ttgggtgggtg	gaaagagact	tagatgcaaa	ttgtgggttt	attcattctg	540
tgactttggg	caaataattt	accttctctg	gaagcaaata	tgtaatttcc	ccttttctat	600
aaaatgacaa	tggaaactagc	atcttaataa	ttttattgtg	aattattgat	tatgtagcat	660
atttcaatgc	ccagctcagg	gtcagacaca	agatagacac	tcaataaatg	ctgattttcc	720
tttcaaaaaa	aaaaaaaaaa	aaaaaactcg	ag			752

&lt;210&gt; 33

&lt;211&gt; 1867

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

ggcagcagct	aaccacacca	tccatctgca	gatagaacat	catcagcaac	attgaaatac	60
cccatgtgct	ctctgcttcc	taatctcagc	gccctcctgt	cctccatgat	gaatgtttct	120
cttgaaattt	actttgttgt	ctttctttca	cttttctgtg	tcgttttacc	attacatgca	180
ttattcctaa	agtcgttctt	ttttttggat	tttgcttgct	ttcaaatttt	atgaaaatgg	240
cagcaaaatg	tttatattct	tctgccactt	gctttttttc	atttaatat	gttttgaaga	300
tctatgcata	ctgatgcaca	ctgatcatat	tttcagtgtt	gtactaatta	cagtttatga	360
ctattccaca	atttgttcat	ccattttatt	cctcattttt	gtgtatttgg	tttggttgga	420
gatttcctgg	cttttttttt	ctattatcaa	ctgggctgcc	acatatgtct	ttatgcctgt	480
ttcatgataa	atacgtgcaa	gaggccaggc	tatgcaaagt	gccagcttta	caagatcatg	540
agcagctgtt	ttgctacgtg	gtattgccaa	tttagactca	cacaagtggg	aaagaagcat	600
ccccagtgct	ctaaatcctt	gccaacactt	agtgtcacca	ggcttcttaa	tttttaccac	660

tctggccagt	ggataatagc	attgaaaagt	tatTTTTtatt	gtgatttggga	gtccaatatt	720
TTTTctgtaa	gaaatgcatg	tacaatgtta	taagaacatg	caaaaatcaa	aatactttat	780
aaatgttcac	tttataaaaa	atgaatgaga	aaaacataac	gcattttaca	taaagaaaaat	840
ctgtaatgct	tttgttattg	gctgggtatt	ttcacacctt	ccaagtgggtg	gctgttgata	900
aattcagcaa	taataccttt	atgtttatta	atattaaatg	ataatgtcga	attaattaaa	960
tgaagttaa	tttattgata	gttattcatt	atatacctgct	gttctaactc	atcctggaag	1020
agaatagagt	gaaagagaaa	ttgcctttat	atataaatga	ctcatagaat	ttcatatact	1080
gacctaaata	gatttcattg	caaagggtatt	atagaggtaa	taacacagta	actcttagga	1140
ctgttttgag	attttcacaa	tttgaaaaat	ccttttagat	ccttggttga	caaatgccct	1200
ggctgtgcta	attatatgac	atttcctgac	actagtgcg	tggcatggcc	tctcccggct	1260
tacattatag	attgttttct	gcccccatgg	gatctgattt	gttaaggctc	atTTTctatt	1320
ttaatgtggt	ggaagaattt	tagaaacctt	agaacccac	tttcacctc	cactgaaaca	1380
aatagaagca	gtgtattagt	cagttttctc	actgttatga	agaaatacct	gagactgggt	1440
aattttatgaa	gaaaagaggt	tgaattggct	catggttcca	cagctgtaca	ggaagcatgg	1500
caacatctgc	ctctggagag	gcctcaggaa	acttttactc	atgggtgggag	gcaaagtggg	1560
agcaggtgtc	ttatgtgaca	gaagcaggac	caagagagag	acgggggagg	tgctacacac	1620
TTTTTTTaaa	acagagtctc	actctgtccc	tcaggctgga	atgcagtgat	gtgatctcgg	1680
ttcgctgcaa	ccttcgcctc	ccaggttcaa	gcattttctca	tgcctgtagt	cccagctact	1740
cgtgaggctg	agacaggaga	atcgcttcaa	cccagagaggt	ggagtttgtg	gtgagccgag	1800
atcgcgccat	tgcactccag	cctgagcaat	cagaacgaaa	ctctgtctca	aaaaaaaaaa	1860
aaaaaaa						1867

&lt;210&gt; 34

&lt;211&gt; 419

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

cggcacgagc	ctccaagccc	ctgtacataa	cctggagcgt	gtgaccttca	gagcttttca	60
ctttatgcaa	aatggctcct	gtgagggctg	caagctggag	ggtggtgcag	gccttgggcc	120
acagggaggc	gcctgtggaa	tagggggagt	tcatgcaccc	ctttttccc	cagaggggct	180
ggactcaggt	tagtttgggg	gtgggggctc	ctgcactttg	ccacaggcac	ggggagggtt	240
ttctcctcac	cccctctgcc	ctcccaactt	gggttgtact	ttctaagaag	gtgattcccc	300
ctgcccttgc	ccccttcccc	agaacaaaac	atgttgatca	tgtgcaatat	ttcttactgt	360
gccgagaagc	cgcaatgagc	gagattaaag	ctgtttaaca	caaaaaaaaa	aaaaaaaaa	419

&lt;210&gt; 35

&lt;211&gt; 949

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (940)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (947)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 35

cnacaagtac	aggttatcaa	gtttgcactt	aactatgcca	aaaaaagttt	gaagcgctct	60
attctcagac	atgctgtatt	attacttctc	attcaagatt	gaaaaatata	aaggatatcca	120
aactctgtct	taatgtaa	gtaactat	ttccttcaag	tgttgactag	ggagtcgggt	180
tctctcttaa	agacactcac	tgtacaactg	aaagcagctg	tcataatttct	ggcaaaatgt	240
gtttacgtat	ctgacaagtt	gtacatttgt	gtatgaactg	acataaaatg	tgaaagcctg	300
taagtgtaca	tgtagtgggtg	tggtgttctg	tctagaggat	acaactgaat	gtttttaatt	360
tgctgactta	cagacacagg	ctgtttacaa	aatgctagct	ggaaagtctg	taatgttcat	420
gtcataactt	ttagttaatt	gccattgagc	acctgttctg	aggaggtgag	atgtggactt	480
gtgcttataa	actggagagt	ttagtcataa	tcctctctgg	ctttgtgtga	atagcttgct	540
cactttgctg	gcctttgaaa	tgtgttctcc	gtgataagct	atccatgtgt	ttgtgataag	600
agtgttgtc	aaccatgacc	atctttgagc	cttcttagtc	ctccacctgg	cacagtattt	660
gaaatggcaa	aggatgtgct	tcatcctcta	acagtgtaca	ctcccagagc	tgatattctg	720
gattttgact	gtgcacattt	cctctagtgc	atgtctgtag	tccttataga	atgatctgta	780
ataaaatagt	atactggact	gtgcatcaaa	gggatgtaaa	attacagtat	tccaaagggt	840
gaagttctgc	tgttttgtta	taatgcctga	tacacatctt	gaataaagtc	ttaacatttt	900
tcttttaaaa	aaaaaaaaaa	aaaaaaacyc	gagggggggg	cccgtancc		949

&lt;210&gt; 36

&lt;211&gt; 1635

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (6)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (24)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (34)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1615)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1630)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 36

gtggangcgc	tctagaacta	tggntcccc	gggntgcaga	ttcggcacga	gctcgtgccs	60
cttctgtgtc	tcagggtaat	actattcaga	gtcgcccctt	tgctcatttt	ctcccgtatt	120

tgttaccttc	ctgaggcctc	agtattagtc	gtgagcacia	agttttgaga	cctttggcgt	180
tgtttcttga	tgtgggaggg	gaggtgttag	tgcatgcaag	ggttgaacta	gatagaccct	240
gccttagtag	agggtagggac	tataacctta	gaggccagaa	cttgatccag	aagttgctgt	300
ccacagaagt	gctttctatt	tcatcatttt	tgtttctagg	gctctttttc	tgtagccagg	360
tcttcccaag	gatttttagta	tttgcatggg	agttgaggtt	tactctaata	atggtagccc	420
agctgtgccc	agaggacagc	caggcaggcc	cygggagggg	gtttagaaag	acagtcctgg	480
tgaatgggct	tcaagtgggt	acaaagaggg	tggctgtgag	gtgaccccag	acactgcaga	540
acgatgtgca	ccctctgcgt	tttgatgtgc	cttggaatgt	gggagcctag	aaataaccct	600
gtggatggaa	ttggggcagc	ggctgctgga	gatctgtgtg	ccttgccctc	cttcagcagg	660
accgtctagg	tgccgagcca	cctatggatg	cgtcccagcc	agccccgtcg	ctctcgtcca	720
tcctcagaga	caaagaagag	ggcagggagt	ttgggcttgg	ttttgaactt	tcctttcaat	780
gtagcaaagc	attcctagtt	aaccagagcc	ttggaatcta	ctgcctgctg	gccaggcttt	840
aaaatgaaaa	gtgttttaat	gctgccataa	aagggagggc	ggggggagga	agggaaaata	900
aaggcatctt	tccaagtact	catctaattt	aattgtcaaa	agattgatag	gccatgaatt	960
acttctccat	ctcactaagg	gttaaaggcg	tgcaaccccc	cactggctgt	gtcccctgcc	1020
accgaagtga	gtgacctgcc	ctacaaccag	gtgggaccac	ctgtgctgca	gtccggaggg	1080
gcttctgcag	gaagcactca	ccccccacac	sttccccggc	ctgagcttcc	cctacctttc	1140
gtcaccacct	gagggcatga	gcacaggcca	tggggcgctg	ctggtgagtc	tgccctgtgt	1200
tcaggcttag	cctgtgggtc	cctgtgtgct	gctgcccgcg	tgggatgcgc	aggggagggc	1260
tggggatccg	caggaggggt	gttgggatac	accggatacc	tctgctctca	ttgcttggtt	1320
scaaattgct	tatggacatt	tgtgtgctaa	atcctattaa	ataaaaaaga	cgggttaaaa	1380
cccagatgct	gtatatctcat	ttgtaattat	gtataaagt	aagcagtttt	aaactgtaaa	1440
gatttttttc	agtgtgtttt	ctcgaatttt	gccacaacat	actggcttcg	tattttattt	1500
atctttcttt	ctagttacca	gcttcagacc	cttgtaaagt	ctccctcagc	cctttcaaaa	1560
aataataaat	ttcctgtgaa	gttaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaanaaaaa	1620
aaaaaaaaan	aaaaaa					1635

&lt;210&gt; 37

&lt;211&gt; 2414

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 37

ggcacgaggt	aaactggttt	ttggtgaaat	tgggctcggc	ctcatgcgaa	cctatctagg	60
tccgtctctg	gagtggcctg	tcacgatcgc	cgccctccat	cgctgagcga	cacaagtggg	120
accccccagg	actgcattct	gcgaaactac	cctgcttttc	ccccagtttt	ggctctgtga	180
cacatttggt	aaattcttct	gacccatccg	ctactctctt	gaggtacagt	aaattgtctt	240
tgacttccat	caccttttgt	aatattacaa	ttgccatcta	ctgagttcct	atgatctgcc	300
tggcattgta	ctaagtgcct	tatatcttct	tgttcatttt	tttatgccaa	cataaatgag	360
tttgactcaa	ttcttattat	aacctgccaa	aataagtatt	cctcttccct	tttacagaga	420
gagaaacaag	ggaaaaaaa	gttaattgcc	tgaggatata	ctgctagtgt	tgccgggaca	480
atagagacgg	gagaggccaa	acagaattca	ggacaaccct	ttattattaa	ggtagtact	540
ggctcagtcg	gactgggggt	cagaaagtct	gagcccggaa	caaagggatc	aatcaccttt	600
taagcaattt	gtggcgggag	tttggtgcta	cagggaagcct	acttgacaga	gcaagaacaa	660
aggcagttaa	ttattctttt	acatttggtt	caccacatgt	cttacatcct	tgggaatgca	720
tgcttttgtt	agcaattgct	tatcaacctt	gtgactttac	agcgcgctag	ggagggaagc	780
aggaactcgt	ggtagcctca	ggaacgtgaa	acgggtggaac	acagataagc	ctctccgtgc	840
gtagaggaga	aataagcggg	taatatcttc	tcttaaccca	ggcctccgga	gggtgggggg	900
ctatatattca	ttctacctct	aaggaaaaga	ataaacttct	tgactaatat	acttataaaa	960
ttcatttaatt	cccccttcac	tagtaaataa	taaatacctg	acttaggatt	ccaaaatagg	1020
tcttgaattt	ttctgcagga	gtcctcctca	gtctctgaat	ccccacaacc	ttttattaga	1080
tcacttatcc	cattgtatta	taattatctg	tttaacaatt	tttcctcacc	cctaaaggac	1140
cctgagcttc	tgggaaggtag	ggactgtatc	ttaaaatctt	ggcattttct	atgtctagaa	1200
tatttcatct	aagagaaggg	cttagtaact	gttgtgtggg	tgggttgatg	agttaactgt	1260

caggcaggtg	aactaatagg	ctgaagataa	agtttttcac	tgttggaagt	gaatttgga	1320
ttagtcatcc	taatccccctc	acttttagag	atggggcaac	tgagtcatgc	gcggtggctc	1380
acgcctgtaa	ttctagcact	ttgggaggcc	aagggtgagcg	gatcacctga	agtcagggat	1440
ttgagaccag	cctggccaac	atggcgaaac	cccatctcta	ctaaaaatac	aaaaattagc	1500
caggcatggt	agcaggagcc	tgtaatccca	gctgttcggg	aggctgaggc	aggagaatca	1560
cttgaacctg	ggaggcggag	gttgcaagtga	gttgagatcg	cgcaattgcg	ctccagcctg	1620
ggcgacaaga	gcaaaactcc	acctcaaaga	aaaaacaaaa	caagacatgg	ggcagctgag	1680
gccttttgta	taccaggcat	ttatcgagtt	attcatgtgg	gccagtttct	gtgataaatg	1740
at ttgcaaat	aatacccaac	ttaatctgta	ttgttat ttt	tatttttaca	gatgaagaaa	1800
cagctcagat	ttgaccaaga	tcacacaagt	aggagttagag	gaattggaac	ccactatgac	1860
tgactccaaa	ttaatatagt	gtttatagcc	acttacataa	taatggtaaa	ggttttat ttt	1920
ttttcactat	acaatatgta	gtagataaat	ttactgagga	aatgagacag	cacgtgacaa	1980
atccttaaaag	gatacacaca	tgacaggaag	gattagaaga	ggaaacagaa	acaggtaaca	2040
gagaagcaga	aaaagtaaaa	tctcccaaaa	taaaaataga	aggaattaaa	aataaaaagca	2100
gactccgggt	gcagtggctc	atgcccagca	ctttgggagg	ccgaggcagg	tgaatcactt	2160
gaggtcagga	gtttgagacc	agcctcgcca	acatgatgaa	accctgtttc	tactagaaat	2220
aaaaaaatta	gccaggtgta	gtggcgtgca	cctgtaattc	tggttacttg	ggaagctaag	2280
gcagaaagaat	cacttgaacc	cgagaggcag	aggttgagtg	gagccaagat	catgccattg	2340
cactccagcc	tgggtgacag	agcaagatta	taaagtcttc	tcaataaaaat	cgtctcaaaa	2400
aaaaaaaaaa	aaaa					2414

&lt;210&gt; 38

&lt;211&gt; 929

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

gggacagctg	gcctctgtgc	agtgggtggc	gggggggtgc	tggacggctg	gcctctatgt	60
agtcggggag	gctggatgac	tggcctctgt	gcagtggggg	caggcgctgg	atgactggcc	120
tctatgcagt	gggtcggggg	gcctggacag	ctggcctctg	tgcggtgggt	tgggggggcc	180
tggacgtctg	gcttctgtgc	ggtgcggggg	ccggcctcgg	gccgktgggc	aggctggaag	240
gtcagccagg	ccctgtggtg	tctttagtga	gaaacaggaa	ataaatcaag	gggatgttgt	300
agaagtttta	ccgctgctgg	tttagaagtg	tttgttttta	ttatttgtat	ataatgtgtt	360
gttttcagtc	ttgcaactgg	gttttttaact	atcgtgtaaa	gatgcaaaca	gcaggccaga	420
aaactgcctt	gatatatcac	attggcaaaag	accaagcgca	ccccggatcc	tagtgggtga	480
gccggaccga	ttggcagtg	cctttgtggt	gcctccagg	agcctcacga	tgctcacgcc	540
tctgaccagt	ggttctgcgc	tcagggcagt	accggagatg	tggctctggga	gttggttgtg	600
tggatgtttg	cgttgccagc	gtttacaaca	acagagggga	acagcccaca	gccagcagc	660
acaatgagga	tggctcctgt	gataaatggg	atgctgtgat	tagattgttt	ctgaataagc	720
taaaatgctc	aaaatgtgat	ctttctttta	taaaaaggcc	aggatgggtg	atacgtgcct	780
gtaaccccag	cactttggga	ggctgaggtg	ggaggatccc	ttgagcccag	gagttggaga	840
atgcagtgag	ctgtgatcac	accctctagc	ctaggcagca	gagcgagacc	ttgttctaag	900
aaataaaaaa	aaaaaaaaaa	aaactcgag				929

&lt;210&gt; 39

&lt;211&gt; 1125

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

gcttaaacct	gattgtagct	atctaagttg	tagccaatga	ggtgtcagaa	aatgtaccat	60
gcagtgcAAC	ttcttaagta	gcctagaaag	atggggtcac	atccccctgg	gttggttctc	120
tctttgtgtc	at tttgatct	ctctcgctgg	aaaacagaaa	tgatgtctgg	agtgtaaagca	180



gccatcttgg	gctaggagac	tacatgctga	catggtgaca	caccaccag	aaagaatctg	240
agtagtcgat	gatacacagag	ctaccatata	agcactagac	tctctacctc	cacattttat	300
ttgtgtgaca	tagaaataaaa	gtttaagctt	tgttttttct	ggggtgtatg	tgtgtacaca	360
tgtgctctc	tgtctgtctc	acatgcagtg	agacctaat	ctaactgata	tgccggtagg	420
atgggggggac	cttgccacaa	ggtataattg	tgggtgggatg	gctgtgtctt	gctcccattc	480
aggatccatg	gaaagctcct	cctccttctt	gttcccctgg	atgccaaagcg	atgggcctga	540
gaccctggct	ttgccatgtg	gatgctcccc	ctcaggactt	tggatgtgca	gtgattgcta	600
caaaaagggc	aatgaagatg	cctggaccac	agccacacag	tctccatcac	agcagtggta	660
ttctgtggaa	tgagtactaa	ttttctggcc	acattttctg	ttaaaaactg	gttgttgagc	720
tgcccacttg	tcttgggtcaa	gggatgctgg	agtggcctgg	gaaggcagtg	ttgtttgctt	780
ttcttaggat	ggaacacact	taggcataatt	tttgggtcaaa	gcaaaaagaag	aattcaaagg	840
tgagagaagt	tgaagataaaa	aaggggaaaa	tgagggtagc	agatgaagca	aagtcaacaca	900
gaaagctggc	aggcgagtag	tagaatcaga	gtacaggctg	agcatcccaa	atctgaaaat	960
ccaaaactcg	aagtgctcca	aaacctgaaa	cttttggagc	accaacacaa	tgctcaaagt	1020
aaatgctcag	tgaagcatatt	tggatttcat	attttcagat	ttgggtttgc	ccaaccggtg	1080
taatacaaat	attccaaaat	caaaaaaaaa	aaaaaaaaaa	aactc		1125

```
<210> 40
<211> 1121
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> SITE
<222> (286)
<223> n equals a,t,g, or c
```

```
<220>
<221> SITE
<222> (1102)
<223> n equals a,t,g, or c
```

```
<220>
<221> SITE
<222> (1105)
<223> n equals a,t,g, or c
```

```
<220>
<221> SITE
<222> (1106)
<223> n equals a,t,g, or c
```

<400> 40							
gtttcttttat	tcctaattgca	gttagaaaaga	cctttctctct	tgagctctttt	gactcccaga		60
aggtaccccca	gtccccagtg	tacttagaaa	ggatctcgaa	cattgctgga	cgctctcata		120
gtactcacia	agggctagcc	ttgaatgtca	ctcgccagtg	cttcagctctc	ctgacttaga		180
gatacaatca	cgtcacaggt	ctcttggcct	caatctgaaa	actgctgccg	ccgcgccgag		240
gagactcgca	tgccgccacc	acctcactgg	gagggcgccg	agccanccgt	cgccccctag		300
accctgacag	ctgcagctgc	cttgcccttg	cgccgcctcc	ctgcaggggc	cctgttccaa		360
tgaaaaacag	aacacaaaag	agcagagcac	ctaagcctgt	ctctgcctcc	ctgtctaccg		420
gactggccag	ggccaagacc	cccgctgctc	cactgcgggg	ctgggcgggc	tgactccctg		480
cttcctccar	gctgtctcct	ccccctgagc	cagggctctg	gcagggtgca	gccggtctct		540
gggcgcacgc	agcttccttc	aagtacactg	tgtgtgcttc	ccggacctgc	ggcgatgccca		600
cgggcctgcc	ttttctatgc	gcctcactag	cttaccaccc	tgtgcaggta	atgcaactga		660

ctttgtctca	tcagtccttt	tctttccctg	ccacccttta	tttatcaagc	gtaatgttac	720
actttaaaag	acagcaaata	agaactttgt	agaatccac	caggactttg	ctaacaataa	780
tgtrtggaag	taaagaagt	ctctgaaaaa	atatcagcca	ccaaaatagt	tatgttggca	840
ctgtgttcac	acgcatgggc	cccacacccc	cagggtgggt	gggttttttt	gttttttggg	900
tttttttggg	gggggggctt	tttcatgtta	catccatata	tgtatttata	tcttatttgt	960
ttcactttca	agtgtatcat	ggcaaagtga	cagatttttt	tgttaataat	gtgctaggat	1020
ttgctaataa	agaaaaaaa	aaaacccttt	tgagtttgcc	ctagaataaa	tgagacttaa	1080
ttcaaaaaaa	aagggaataa	gnagnnggaa	aaaaaaaagg	g		1121

&lt;210&gt; 41

&lt;211&gt; 2168

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (676)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (735)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1551)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 41

ggcacgagcc	aggtagcccc	atccccacaca	ctttccacta	gtgcctatat	tgggatgaat	60
cttctggcaa	tcttccttaa	aacttcaaga	gacctcttta	agattctgta	gttgggtgag	120
cctgtctgtg	cccagggtag	cctgaagtta	aggtgagggc	ttaacagtgg	taccaggcag	180
tttggtatccc	aactagctag	ctgtccttgg	gcaaatagact	actttccctac	tactgtttact	240
tcttggtttct	attttctcct	ctgtaaaactg	tgggtaaatga	aaagtgtgcgc	aaagttaggg	300
gggtgagtga	cagctacagg	ctcactgttc	ctgaaaactg	caattcgaaa	ttccacagcc	360
ccctgaaaaat	aaaaagtctt	tcagtgaatg	ttcagcaaac	tcccttggtg	ggcaaaaactg	420
acttgaatta	acatgaggct	atattatgata	ttttgtaaat	ctcagttagt	gagactagtt	480
gtacattttta	ctgtagaaat	atccatgtgt	ttgatttcag	agtactacce	tgaaaaatacc	540
cagggtatttg	gaagcataga	atatatgaat	taaatgactt	ttttaaaaaa	ttgtgggtaaa	600
atacatataa	cacaacattt	actatcttga	ccacttttta	gggtacagtt	caatagtgtt	660
tgtatatatt	caattgntgtg	caaccaatct	ccagaattct	ttttacttat	taaactgaaa	720
ctctgtacct	atcanacaac	tccccgtttt	cccctttccc	cagccccctgg	caaccacccat	780
ttgacttttcc	aaatttcttt	tagaattgca	aaaattccaa	attccaaaac	acatctggcc	840
tcacaagcat	ttcagataag	ggattgggaa	acttaattag	ggaaggggaa	tgcctamcat	900
ttgatagagt	caccaccttt	cgagatgata	ctggaagagg	gagttcttcc	aatcttacag	960
gggcctgtca	accacagcac	actgttctcc	aaywtcgatg	ccctttttgt	aaagatggca	1020
cagkrsgcaag	aacaagggcc	attagattct	gaccagagca	aacagaacgc	aggtagttgc	1080
atgggatgtt	tgataccttc	cccttctttt	ctgtcattcc	cttgatctca	gttttttytca	1140
gtaagtagaa	gtgcacatgt	tggcagctgg	ttctggcatc	tactttttat	ttccattcca	1200
gggaccatga	tggaggatgc	ttttacagtg	aaaaggcatg	aaaaccttta	tgcagactga	1260
actcatgggg	aagatgctga	cagttctgtt	agaacttgct	ggaagccatt	gaaggccaga	1320
tttatccatt	ttgcagaact	ctctaggaat	cttcagaaaa	gcagtagggc	ttactctgct	1380
gttccccagc	agatcacagt	atggagaccg	gttctgagtc	atgctcccta	taactggaat	1440

aacacagggga	attcttcaca	tgtttcataa	tgtgtgtgag	tgaaaggaca	accagactt	1500
gktattgaaa	aaccgmcag	tggtcaggca	tcattattgg	aatgtcttct	nccacactgc	1560
ccattctgta	aacatcctgg	ggaaatgtcg	aggttacttt	ctgtgtgagg	sttgtgcktt	1620
cttwtcctgk	ttgtaaatac	cagggatgaa	agtggatgcc	tyaraattg	gagccctgaa	1680
cmcaaaattc	tgcagaatac	aaaaccctct	gatggaccac	tcctgataaa	tataaaataa	1740
ccttagtacc	agaacttcta	cttttggggt	atggaaaata	tgccaagaat	tttatgtttt	1800
aaaaacaaac	tacaggctgg	gcgtgggtggc	tcacgcctgt	aatctcagca	ttttgggagg	1860
ccaagccagg	tggatcaact	gaggtcagga	gttcgagacc	agcctggcca	atatggtgaa	1920
accccatcac	tactaaaaat	acaaaaatta	gccaggcatg	gtggtatatg	catgtgttcc	1980
cagctacttg	ggaggctgag	gcaggagaat	tgcttgaacc	cgggaggcag	agtttgcagt	2040
gagccgagat	cgcaccgttg	caccacagcc	tgggtgacaa	gagtgagact	ctgtgtcaaa	2100
aaaaaaaaaa	aaaaaaaaac	tcgtaggggg	ggaccsgtac	ccaatccgac	cctgtgagtg	2160
tattcgta						2168

&lt;210&gt; 42

&lt;211&gt; 2105

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 42

ggcagagtca	gtacagagca	gcgttttctaa	agcacaaatg	tgatcattcc	ctcttgctta	60
aaattacaca	atggctttct	cttagttatt	ggccaaaatc	taaactcggt	ggctaaactg	120
ttcatgatct	gactatgctt	gccctttcta	ccagtcctcc	ttatcacact	tactttctag	180
aaaactgaac	tgtttgctgt	tttctgagta	agtcatgaaa	tttgtgcttt	tctacctttg	240
cacaggttat	cccctttttt	gggaatgttt	ctttattttg	taactgtgat	tcattcttaa	300
acatgtcatc	tccagatata	tctgactctc	catcttttta	ccctagagtt	tggcccatca	360
ttttgtgctc	aaataagaca	ttaattgcat	ccttcatagt	atttaggaca	aagtattata	420
tttgtgtgtc	caccatgcca	aatagatgac	agctcctttc	tttagagtaa	tattttctata	480
ggagtagaat	acagtgttaa	agagtacaaa	ctttttactc	agttctgatt	ttggctctac	540
tttttaaagt	agactatact	agagatataa	ttgcctcata	ctaaaaatsa	gatgataata	600
ctatctgagg	atgacttgta	ttaagtgaga	taatgcatct	aaactagtgt	tgtctggcac	660
atggaaatct	cttagtagat	ggaagtcaca	ttagtatctt	cagcacagat	tcattattca	720
gcaaataatct	gttgaataaa	tgatcacata	tcaggctact	aggatatatg	cactgttaac	780
tggtagtatc	taactcggta	attggttatc	tacttagtat	ctggtaagga	tatataaaag	840
aatcttatt	tctgtagaag	cctycacacc	actaatamca	gtaataggaa	atggctaaga	900
aacatacagg	atattatgtg	gttacacact	tttataaaag	gggagaattc	actgtaaatt	960
ctcaaatgat	cagtcttaca	tgatcaaaaag	ttctaaattc	aacatttgga	ttttattctg	1020
taggttgagt	aatggcaatt	catggagggt	ataaagcaga	ggacaagcat	gatcgaattg	1080
gtcctttaag	aaggataaac	aggatgatgt	ttagacagtg	atagggttg	agggaggag	1140
tagggtagtg	acacctgaag	gagggtgata	agtgggaatg	gagttggggc	tatgaatgaa	1200
attcatttga	agaaagacat	atgggtttga	gtatagagaa	taagagacag	ataggagtca	1260
aagattatct	tggccatgag	cctggatgac	tgagagaatg	ataatatcat	tggaggaaat	1320
agagatgtgg	aagagggatc	ttgtttgggg	aaaagtgggt	tagggaaaat	gagtttactt	1380
ggggcttatt	atttgagttg	ttcagtgtaa	aaattcttgt	aggctcttgt	agatacagaa	1440
ttggagtaaa	gatgagagac	tgggtgctgaa	tgtgagtact	agtactgttt	actaagggaa	1500
tacttttraa	tgtttgtttt	tttaatgtgt	tgtttttttc	tttaaaatca	agaccagttg	1560
ttacaaagaa	rtaacatgtt	tkartgtcat	tgtcgtttta	aaaatatatt	tacttcttaa	1620
agtagactag	ccaaaaggtc	ccgaacttgg	tatctaaaat	gaatatgggg	aataaggatt	1680
ttgaaaaata	gctgttgctc	tcattctttt	tttttttttt	caatatgaga	cagggtcttg	1740
ctgtgtcacc	cagggttgat	tgcagtgggt	tgatcatggc	ttactgcagc	ctcaacctcc	1800
caggactcaa	gcaatcctca	cagctcagcc	tcctgagtag	ctgggactat	agatctgcat	1860
caccacatct	ggcaaatttt	ttgtattttt	tgttgagaca	ggatctcact	ttgttgtcca	1920
ggctgggtctt	gaactcctgg	gctcaagcga	tcctgtcacc	ttcgctccc	aaagtgtgta	1980
gattacaggt	gtcagccata	gcagccggcc	tattctctca	ttcttgaaga	atgaaagctc	2040

```
catgttatgt ttgaagtttt acgagttgac tgctttgtaa gaaaaaaaaa aaaaaaaaaac 2100
tcgag 2105
```

&lt;210&gt; 43

&lt;211&gt; 1491

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

```
ggcagagggg aggtcgtcta gggaatcgag gtgccggctg ctcttcctc acaatttggg 60
ttgtgctgca aggggagggg ccccatcatc tggccccagt ggtgtaagga gctgactggg 120
attcagtcac tgacttggag ccgctcgggg gaagtcccg gcaggagcgc ttcacctcta 180
tgacacgatt gtattatcgg gatgcctctg cctgtgttat tatgtttgac gttaccaatg 240
ccactacctt cagcaacagc cagaggtgga aacaggacct agacagcaag ctcacactac 300
ccaatggaga gccggtgcc tgcctgctct tggccaacaa gtgtgatctg tccccttggg 360
cagtgaagccg ggaccagatt gaccggttca gtaaagagaa cggtttcaca gggtggacag 420
aaacatcagt caaggagaac aaaaatatta atgaggctat gagagtcctc attgaaaaga 480
tgatgagaaa ttccacagaa gatatcatgt ctttgtccac ccaaggggac tacatcaatc 540
tacaaaccaa gtctccagc tggtcctgct gctagtagtg tttggettat tttccatccc 600
agttctggga ggtcttttaa gtctcttccc tttggttgcc cacctgacca ttttattaag 660
tacatttgaa ttgtctcctg actactgtcc agtaaggagg ccattgttca cttagaaaag 720
acacctggaa cccatgtgca tttctgcac tcctggatta gcctttcaca tgttgctgac 780
tcacattagt gccagttagt gccttcggtg taagatcttc tcatcagccc tcaatttgtg 840
atccggaatt ttgtgagaag gattagaaat cagcacctgc gttttagaga tcataattct 900
cacctacttc tgagcttatt tttccatttg atattcattg atatcatgac ttccaattga 960
gaggaaaatg agatcaaatt tcatttccca aatttcttgt aggcggttgt ttcagattct 1020
ttctgtcttg gaatgtaaac atctgattct ggaatgcaga aggaggggtc tgggcatctg 1080
tggatttttg gctactagaa gtgtcccaga agtcaactgta tttttgaaac ttctaacgtc 1140
ataattaagt ttctcttgtc ttggcatcaa gaatagtcaa gttttttggc cgggcatggg 1200
ggctcatgcc tgtaatccca gcacttgggg aggcccaaggc aggcggatca catgaggcca 1260
ggaattcgag accaacctgg tcagcatggc aaaacccgt ctctactaaa agtacaaaaa 1320
ttagccaggc gtgatggcac gtgtctgtaa tcccagctac tctggagact gaggtgggag 1380
aatcgcttga gactgggagg cagaggttgc agtgaaccga gatcatgcca ccgcacttca 1440
gcctgggtga cagagaaggc tccgtctcaa aaaaaaaaaa aaaaaactcg a 1491
```

&lt;210&gt; 44

&lt;211&gt; 2902

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1059)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1064)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2902)

<223> n equals a,t,g, or c

<400> 44

caggcttcta	atggtgatgc	ttctacttta	ggggaggaga	caaaaagtgc	ttctaattgtg	60
ccatcagggg	agagcactga	tgaagaagag	gaggcacaga	ccccacaggc	tcctcggaca	120
ctgggtccat	cacctcctgc	cccatcatcc	actccaacac	caacagcccc	tattgccact	180
ctgaaccagc	ctccaccact	tcttcgtsca	acactgcctg	ctgccccggc	tcttcaccgg	240
cagcctcctc	cactccagca	gcaggctcgg	ttcatccagc	cccggccaac	tttaaatacag	300
cctccaccac	ctcttatctg	ccctgctaata	tccatgccac	cccgtctaaa	cccaagaccg	360
gtgttggtcca	cgggttggtg	tcaacagcca	ccatcactta	ttggaattca	gacagattca	420
cagtctctac	tgcactaaaa	attaaattgg	acacagctgc	agtaactttt	caccccatca	480
ttataccagt	gctcatctga	ctgatgaaaa	agaggaaaga	ataatcattt	ctagatactg	540
aggctgcgaa	ctagttctgt	ggcagtggtg	tagcataagt	ggatgtctaa	gaaatttttc	600
agttcactag	actaaaatgt	tttacaacaa	aaagcctcca	gttagcctcc	tttctagagt	660
atatgttcag	caatgtgatc	tcataaaaagg	aaaaacaaaa	gatttaagta	ttctatatac	720
caagtttttg	ttttgttttt	actgtatttta	ttttattgag	gttctttata	ttcctgcctc	780
ttcatagtca	aggctcttag	tacaggaata	ttgacttagg	aattgtgaaa	actccttaag	840
tttcttaagt	taaggatgtt	tggctttttt	ctttaatttt	ttaaaaacca	tttycctatg	900
ttaggagtgc	aagaatagcc	agcatttccg	attttgacat	atgttcattt	tatgcatatt	960
taagaaatta	tagctgcata	tcccttcttt	caaaaaatgt	gctttttttt	ttaaaggaat	1020
tttaatatat	ycctttaaar	gaaagcaatt	taatcaatng	caangcaatt	atataaaaacc	1080
acaaagaatg	tactgaacct	actaaccttc	taacatacag	tttaggggtcc	tagcgcagag	1140
tccttggtta	aaggctcattg	actcatcatc	tgctagtaat	gagaggattg	gaagaataat	1200
tttgcataca	aatgaggact	taatttggtg	aaatataatc	tctttaagtt	ccttgaaaaat	1260
ggagttgggt	ttttttgttt	ctaaatgcta	tctgctttta	actagtagtt	gcctacatct	1320
ggggacttca	gagaagaatt	atattttgtt	agttaagtag	acacagtggg	tatggaagca	1380
tttctttaca	gtacctttta	cgtgttttgt	ttctgaactt	aaaattgccc	tcatacttaa	1440
taatatgggtc	tgcattttaat	atgaaagggtg	ttttattgat	aaatctattg	tactattttg	1500
atacatttgt	gtattccttg	cagccaacct	gtattcgtgg	gattgggtgta	gggttaaatac	1560
atcaacatta	tttcataaaa	taagaatttg	ttctgtgtta	tctaaagatg	tatcagtata	1620
ttgtcacagt	tgtgctgtta	actaaaaatg	ctgagacccc	tttttataga	aaaacaaaaa	1680
gacatcaagt	cttcttaatt	caaccataaa	tcattaagta	cttaacaaaag	aatatttttac	1740
aagtgatagt	atttcaacaa	tgtgtaatta	atatttttga	tacagtgtat	tcataattgga	1800
atcattattt	gtgcaaagg	acagacagat	cacttagatt	gctatactag	tggacatagg	1860
ctaaatgttt	gcacattcac	attcttatca	cgtgtagaat	acttcacaaa	atagtcaaca	1920
tctaaggccc	taatttatgt	tttgaaagat	catgtgttcc	caaagtattc	cctattgttg	1980
gctccacagc	cttaaagtgc	tatagattta	aattcattga	ttagttttaa	tttttaattt	2040
tagactgtgt	atttccataa	ataccctacg	tactggcata	tttgaaactc	tttttccagg	2100
ttaggtcctt	ttctttctca	ttgaatcatc	ttaaatagtt	cttggccctg	aatttagctg	2160
atttaaaatt	cttaatatc	aagaatttat	acttattttt	tccttaaaag	ccacagggga	2220
cagttaaata	tcttaaaata	tctaaaacat	tttttaaagc	acttagattg	tcttacgtat	2280
gtgcatacta	tacctttaca	gcgtttattg	tcttgctctc	tgtcagtaga	ccttcagtag	2340
acagtatgtg	ggatatgtca	gtcaagttgg	tcagcaccag	catctgtcca	gctgttcagt	2400
atattgtgat	tcattaaaaa	atctcttcta	tcacagacat	gggccaagg	gctgtatctg	2460
agggatgtgc	tgtaatgtga	tttacatgca	ttagagcaca	cagtagaaaa	acgttagctt	2520
cattagtaat	atgacacatg	tatatagtga	gatgtcttta	ttgtgtgctt	tgcataattt	2580
gtaaatatatt	tgcacgtcat	tatttttctt	ttttgtttta	gcagtgtttg	gcctggaaga	2640
gtgatatgct	tgctgcttaa	tcaaaggatt	aaagatttaa	agatgtctat	gtcttctatt	2700
tttatataat	ttcatgttct	atgaggaatt	tagtacctct	tcactgtgaa	attcgaaata	2760
atgattttta	taaaagcaaa	actagaaatc	ttttaatgac	aattttcatt	aatttcagg	2820
ttatcatttt	tgagaaatct	acaccaaagt	ggttttttta	aattacmtaa	ctaaaaataa	2880
accacaccgt	gggggttacct	cn				2902

<210> 45

&lt;211&gt; 1460

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 45

atgttcgtat	gttctttccac	ttcagtga	ttctggatgg	gaaccagctc	catattgcag	60
atgaagtaga	gtttactgtg	gttcctgata	tgctctctgc	tcaaagaaat	catgctatta	120
ggattaaaaa	acttcccaag	ggcacggttt	catttcattc	ccattcagat	caccgttttc	180
tgggcacggt	agaaaaagaa	gccacttttt	ccaatcctaa	aaccactagc	ccaaataaag	240
gcacagagaa	ggaggctgag	gatggcatta	ttgcttatga	tgactgtggg	gtgaaactga	300
ctattgcttt	tcaagccaag	gatgtggaag	gatctacttc	tcctcaaata	ggagataaag	360
ttgaatttag	tattagtgtg	aaacagaggc	ctggacagca	ggttgcaact	tgtgtgcgac	420
ttttagggtcg	taattctaac	tccaagaggc	tcttgggtta	tgtggcaact	ctgaaggata	480
attttggtatt	tattgaaaca	gccaatcatg	ataaggaaat	ctttttccat	tacagtgtgt	540
tctctgggtga	tgttgatagc	ctggaactgg	gggacatggt	cgagtatagc	ttgtccaaag	600
gcaaaggcaa	caaagtcagt	gcagaaaaag	tgaacaaaac	acactcagtg	aatggcatta	660
ctgaggaagc	tgatcccacc	atctactctg	gcaaagtaat	tcgccccctg	aggagtgttg	720
atccaacaca	gactgagtac	caaggaatga	ttgagattgt	ggaggagggc	gatatgaaag	780
gtgaggtcta	tccatttggc	atcgttggga	tggccaacaa	aggggattgc	ctgcagaaag	840
gggagagcgt	caagttccaa	ttgtgtgtcc	tggggcaaaa	tgacaaaact	atggcttaca	900
acatcacacc	cctgcgcagg	gccacagtgg	aatgtgtgaa	agatcagttt	ggcttcatta	960
actatgaagt	aggagatagc	agaagctctt	ttttccatgt	gaaagaagtt	caggatggca	1020
ttgagctaca	ggcaggagat	gaggtggagt	tctcagtgtg	tcttaatcag	cgcactggca	1080
agtgcagcgc	ctgtaatgtt	tggcagtgct	gtgagggccc	caaggctgtt	gcagctcctc	1140
gacctgtcgc	gttggtcaat	cgcttgaaga	atatcactct	ggatgatgcc	agtgtcctct	1200
gcctaattggt	tcttcgtcag	ccaaggggac	cagataactc	aatgggggtt	gggtgcagaaa	1260
gaaagatccg	tcaagctggt	gtcattgact	aaccacatcc	acaaagcaca	ccattaatcc	1320
actatgatca	agttgggggg	aatctggtga	agggttctga	atatctccct	cttcatccct	1380
cccgaatctt	ggaatactta	ttctattgag	ctattacacc	agttttaaca	ccttctctgt	1440
gttatgttta	aaaaaataaa					1460

&lt;210&gt; 46

&lt;211&gt; 896

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 46

ggcacgagct	aatttgggggt	atcactgagc	tgaagacaaa	gagaaggggg	agaaaaccta	60
gcagaccacc	atgtgctatg	ggaagtgtgc	acgatgcac	ggacattctc	tgggtggggct	120
cgccctcctg	tgcacgcggg	ctaataattt	gctttacttt	cccaatgggg	aaacaaagta	180
tgcctccgaa	aaccacctca	gccgcttcgt	gtgggtcttt	tctggcatcg	taggaggtgg	240
cctgctgatg	ctcctgccag	catttgtctt	cattgggctg	gaacaggatg	actgctgtgg	300
ctgctgtggc	catgaaaact	gtggcaaacg	atgtgcgatg	ctttcttctg	tattggctgc	360
tctcatttga	ttgcaggatc	tggctactgt	gtcattgtgg	cagcccttgg	cttagcagaa	420
ggaccactat	gtcttgattc	cctcggccag	tggaaactaca	cctttgccc	gcaccgaggg	480
ccagtaacct	ctggatacct	ccacatggtc	cgagtgcact	gaacccaagc	acattgttga	540
atggaatgta	tctctgtttt	ctatcctctt	ggctcttggt	ggaattgaat	tcattctgtg	600
tcttattcaa	gtaataaatg	gagtgccttg	aggcataatg	ggcttttgct	gctctcacca	660
acagcaatat	gactgctaaa	agaaccaacc	caggacagag	ccacaatctt	cctctatttc	720
attgtaattt	atatatttca	cttgatttca	tttgtaaaac	tttgtagtag	tgtaacatac	780
tccccacagt	ctacttttac	aaacgcctgt	aaagactggc	atcttcacag	gatgtcagtg	840
tttaaattta	gtaaaccttct	tttttgtttg	tttaaaaaaa	aaaaaaaaaa	aaaaaa	896

<210> 47  
 <211> 2070  
 <212> DNA  
 <213> Homo sapiens

<400> 47  
 agaagactaa gggggcgctct ccccggtctc ctgcggcgctc ggtggcgagc tgaggtggag 60  
 gcaggctgcg gcagacggcg acagtggcgg cggcgcatgg cagggttgc aggatecctg 120  
 ctgccttggg gatcccgggc tgacagccag agagcacagc ggctcagctc ctggagagtg 180  
 aggggtgaag aaagcggagg gcagccgctt gcgcccgtg gctccatta ggtcgggttc 240  
 tgcagcggtg cccggcagcc ttggtgaagg ccctgcccgg cagagatcat gtattgcctc 300  
 cagtggctgc tgcccgtcct cctcatcccc aagcccctca accccgcctt gtggttcagc 360  
 cactccatgt tcatgggctt ctacctgctc agcttcctcc tggaacggaa gccttgcaca 420  
 atttgtgcct tgggttttcct ggcagccctg ttccttatct gctatagctg ctggggaaac 480  
 tgtttcctgt accactgctc cgattccccg ctccagaat cggcgcatga tcccggcgtt 540  
 gtgggcacct aacggcctgc cctgttagct ttccaaggaa gcagaagacg ggaggggagg 600  
 cattgacata ggtcataaag cattggagtt tcaaattccc cagcctcgcg ggtgtcacat 660  
 tccctgacgg gccttttttg cctgtgatgt tttatcctta caatgtgaat aatggcactg 720  
 accggtgctt tcatattgtaa gtccctatagt cgtgggtggg cttgtggttg tgtgtgttct 780  
 gtccccatct aggtcctggc tggcgcgatg accaccctc tcgctcatt actgtgagga 840  
 gtctgggtcc atcctggtca gctgccccaa tgtgacctgg ggcagataaa atgccagtct 900  
 cattgtcacc tctgtgacct ctccttgcca gggctcctt ccttcccaga atgttactga 960  
 ctctcagtc cctcttctgg tttcccttta tttctcttct acccttttcc ttttttgggg 1020  
 agtacctgtc caagacaggg ctcatTTTTG cacttatctc gaatttgaag agattgctga 1080  
 cgcccgagag cctcgctttt tcatccttct ttccttgctc agcaggctag acagaaacat 1140  
 gtcttgactg ttagttgtcc acaaatcttc agtattttct ccacttcatt ttttaagaaag 1200  
 gaagcaacag atagatgttg ctctttcacc tgggtgtctg ggctcaagct ttcccgccca 1260  
 gcctcacttc ctttgccctt ctcctgctt tttcactg tccaaggag ggggcctcat 1320  
 tgtgtctccc gtgcatgctc tgcagcattg aagtatgggt tgttcacgta gttctagcag 1380  
 tccccagctg agtgagtggg agagtacctg tgtgtttcgt aacggccttg atccccctga 1440  
 tagatgtttg gatatttttt ggtgtgccct gtgtgtgtgt gtgtacaaat acatgtgtat 1500  
 attcctttta aagaagcttt atcgaacgtg gtctgatttt gaggttttagc aatagctagc 1560  
 tatatatggg aggtgcccgt acagttttta tttagcatgg ggattgcaga gtgaccagca 1620  
 cactggactc cgagggtggt cagacaagac agaggggagc agtggccatc atcctccccg 1680  
 caggagcttc ttcgttctct cgcatataga ctgtacgtta tgaagaatac ccaggaagac 1740  
 tttgtgactg tcacttgctg ctttttctgc gcttcagtaa caagtgttg caaacgagac 1800  
 tttctcctgg cccctgctg ctggagatca gcatgcctgt ctttccagtc tgatccatcc 1860  
 atctctctct tgcctgaggg gaaagagaga tgggccaggc agagaacaga actggaggca 1920  
 gtccatctag ggaatgggac tgtgaggcca tacttgtgaa acgtctggac tgctattcta 1980  
 gagcttttat ttggtgtgtt cgttgcacag ctgtttgaaa tgtttaataa agctttataa 2040  
 acttttaaaaa aaaaaaaaaa aaaaaaaaaa 2070

<210> 48  
 <211> 569  
 <212> DNA  
 <213> Homo sapiens

<400> 48  
 cgagaagaaa atagtgaatc actgtagcaa tggctttgat tcagacctta aaatcacata 60  
 agaagaatta caacatgtta tggattttta agtggcaggt attgtaactg ttttttgtgt 120  
 gcaaaatact gagtaaccac tgggaaaata tttcagatga aagggatgac aaaagcatgt 180  
 tgcgctttgc atcagcaagg cattgacttc tgaaaaaatg atctgaaaaa agtttccaccg 240  
 tttgtcttct tacctcattt taagaagcat gtgaaaatgg gatactatag actactgaga 300  
 atttcagaaa ttgagaacaa tttcataata aaacggctat atttgaagag agaatacatt 360

ttatataaac	aggaaaatac	at ttgacact	ttatggaatt	ttatgagact	ttttgtggga	420
acagaaggtc	ttcaaattgt	aaaatgtaaa	gattgctctt	tttattaagt	ctttaacagg	480
gatgtatttc	attgtatgtt	ttgggtatgg	ctttggaata	aatcatttta	tatttttaaaa	540
aaaaaaaaaa	aaaaactcga	gggggggggc				569

&lt;210&gt; 49

&lt;211&gt; 1255

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

gagatgaaat	gtcttttctc	caggacccaa	gtttcttcac	catgggggatg	tgggtccattg	60
gtgcaggagc	cctgggggct	gctgccttgg	cattgctgct	tgccaacaca	gacgtgtttc	120
tgtccaagcc	ccagaaagcg	gccctggagt	acctggagga	tatagacctg	aaaacactgg	180
agaaggaacc	aaggactttc	aaagcaaagg	agctatggga	aaaaaatgga	gctgtgatta	240
tggccgtgcg	gaggccaggc	tgtttctctt	gtcgagagga	agctgcggat	ctgtcctccc	300
tgaaaagcat	gttggaccag	ctgggcgtcc	ccctctatgc	agtggtaaag	gagcacatca	360
ggactgaagt	gaaggatttc	cagccttatt	tcaaaggaga	aatcttctctg	gatgaaaaga	420
aaaagtctta	tgggtccacaa	aggcgggaaga	tgatgtttat	gggattttatc	cgtctgggag	480
tgtggtacaa	cttcttccga	gcctggaacg	gaggcttctc	tggaaacctg	gaaggagaag	540
gctcatcctt	gggggagttt	tcgtggtggg	atcaggaaag	cagggcattc	ttcttgagca	600
ccgagaaaaa	gaatttgggg	acaaagtaaa	cctactttct	gttctggaag	ctgctaagat	660
gatcaaacca	cagactttgg	cctcagagta	aaaatgattg	tgtgaaactg	cccagctcag	720
ggataaccag	ggacattcac	ctgtgttcat	gggatgtatt	gtttccactc	gtgtccctaa	780
ggagtgaaga	accattttat	actctactct	cagtatggat	tattaatgtt	ttttaatatt	840
ctgttttaggc	ccactaaggc	aaaatagccc	caaaacaaga	ctgacaaaaa	tctgaaaaaac	900
taatgaggat	tattaagcta	aaacctggga	aataggaggc	ttaaaattga	ctgccaggct	960
gggtgcagtg	gctcacacct	gtaatcccag	cactttggga	ggccaagggtg	agcaagtcac	1020
ttgaggtcgg	gagttcgaga	ccagcctgag	caacatggcg	aaaccccgtc	tctactaaaa	1080
atacaaaaat	cacccggtg	tgggtggcagg	cacctgtagt	cccagctacc	cgggagggtg	1140
aggcaggaga	atcacttgag	cctgggaggt	ggaggttgcg	gtgagctgag	atcacaccac	1200
tgtattccag	cctgggtgac	tgagactctt	aactaaaaaa	aaaaaaaaaa	aaaaa	1255

&lt;210&gt; 50

&lt;211&gt; 2354

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

aaaaagtcac	cttgtaactc	atctcttttt	attgtaagtt	tattaaaaat	gaagaggaca	60
acaatgagaa	ggaacataaa	gggttagcta	gcactgtctc	ctgggtgcatg	gggctgtgca	120
gatgtcccg	ccacttcttc	cttcatactt	cccttagaga	acttgctctg	ctacaagcag	180
tgggcttggga	ctaaaagtga	ttaaaatacc	acaggcataa	ggagaaaagg	agtatatgta	240
gtagtaataa	ttactagtat	aaattatttt	cttcacatgc	tatgagtaat	aatattaaaa	300
aactcatttt	accattaaga	ttccttatgc	tgaagctctt	ccatttagaa	tactgtcaat	360
gtcattttact	ggtatgaact	aaagggtccc	cttcttttcc	acttcaactgg	gaaccttagt	420
aaaacaccag	catatcttac	ctctctttct	gactggccga	tgcttccaga	gactgaatgt	480
tgggaaaacc	tagtagccaa	acaattytag	gacagaataa	cattttttata	tttggttcca	540
ccatcttatt	acatttagtt	atagttttta	aaaagaaatt	caagcccat	aaaatatgtc	600
tgggtcaatga	aatgcttcct	tttattgtgt	tgtgctattg	tactttgttt	ttcaaaacat	660
tgtaaaaaata	gtatcttttg	tttagtattt	tggattatat	attataatct	gaggagtgtt	720
ttgcttatgt	agaatccaga	tatatttctg	ttacctagga	gatgttactt	acatatgtaa	780
tactgtatcc	tgcacgtgga	aatattcaga	attgtagata	gcataactct	ccctgctcct	840



attcttttga	gcctaggtat	aatttttttt	tttttttttt	agaaaaagac	atattttagct	900
ttaatttcta	tttatgctaa	acatattttat	aagtagtctg	tcaatataat	accaactatt	960
tttattttta	cataattcaa	ttatttcatt	tgacatgtct	ggcagactca	agacatttaag	1020
taaaaaattg	gaactatgat	ttttctttgt	catttttttaa	aaaagaatta	ttttattaac	1080
ctgctggcat	ataatctgga	gttcttttca	caaccttact	ttttctgatt	tgcttttattg	1140
aatgattgaa	tactcatttc	tttctaaaaa	tatgttgtaa	attctccctt	ggcaagattt	1200
ctccctatga	gggtagttat	tatttgagtc	tgccaagtgg	ttaccatggg	gcaagggtgcc	1260
atgatgtatt	cttgggtgca	ttggtttttt	gcgcatgtga	aatttaagac	acttatagta	1320
agtggactca	ttcatagatg	agtttcagaa	cctttttacgt	tctcggtaga	ggcttctgtc	1380
ggacaggcag	aagagtgtat	tcctcacttt	tttttttgtc	ttcaaattcc	agtaaggcat	1440
agcactttta	agaaattaga	atttttctat	catctatgca	aatgatattt	atgttaatat	1500
taaatatctt	atgttacact	gggagtaatt	tgagggtgcaa	ttatttttat	tactactttg	1560
aatagaggac	cattatcctt	ctttcttcag	aaaactaaga	agtaagtgtg	actttttaaag	1620
taagtatata	tcagtggagag	taggcttggt	ttacaactat	ttctagccag	tgagttgtgt	1680
tttcatgtct	catcaaaaaga	caataccaca	ttgcatcatt	ttacaaaata	tgttgtcatt	1740
ttcatttcag	ttgtaacata	ggaaaataga	tatttcctag	atgatttctg	agtttcttac	1800
tgcaaaagaac	agttataaat	tggtatacat	gtgtctctgt	aatagggata	atattgatat	1860
atctgttgct	acatatttaa	gaatcattct	atcttatgtt	gtcttgaggc	caagattttac	1920
cacgtttgcc	cagtgtattg	aattgggtgt	agaaggtagt	tccatgttcc	atttgtagat	1980
ctttaagatt	ttatctttga	taactttaat	agaatgtggc	tcagttctgg	tccttcaagc	2040
ctgtatggtt	tggattttca	gtaggggaca	gttgatgtgg	agtcaatctc	tttgggtacac	2100
aggaagcttt	ataaaatttc	attcacgaat	ctcttatttt	gggaagctgt	tttgcatacg	2160
agaagaacac	tgttgaaata	aggaactaaa	gctttatata	ttgatcaagg	tgattctgaa	2220
agttttaaat	tttaatgttg	taatgttatg	ttattgttaa	ttgtacttta	ttatgtattc	2280
aatagaaaat	catgatttat	taataaaaagc	ttaaattctc	aaaaaaaaaa	aaaaaaaaaa	2340
aaaaaaaaact	cgag					2354

&lt;210&gt; 51

&lt;211&gt; 2267

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

gggtcgaccc	acgcgtccgc	ccacgcgtcc	ggaaggcttt	gggcacagac	cacacaagga	60
tctatgggca	agcaaaaatg	aaaacgaaga	gatttttgag	agaccagccc	agcttgcaaa	120
tgcaaggagg	acccctcaca	gcccaggggt	agaagatgcc	cctattgcta	agggtgggtgt	180
cctggctgca	agtatggaag	ccaaggcctc	ctctcagcag	gagaaggaag	acaagccagc	240
tgaaccaag	aagctgagga	tcgcctggcc	acccccact	gaacttgga	gttcaggaag	300
tgcttgagg	gaagggatca	aaatgtcaaa	gccc aaatgg	cctcctgaag	acgaaatcag	360
caagcccga	gttcctgagg	atgtcgatct	agatctgaag	aagctaagac	gatcttcttc	420
actgaaggaa	agaagccgcc	cattcactgt	agcagcttca	tttcaaagca	cctctgtcaa	480
gagcccaaaa	actgtgtccc	cacctatcag	gaaaggctgg	agcatgtcag	agcagagtga	540
agagtctgtg	ggtggaagag	ttgcagaaaag	gaaacaagtg	gaaaatgcca	aggcttctaa	600
gaagaatggg	aatgtgggaa	aaacaacctg	gcaaaaacaa	gaatctaaag	gagagacagg	660
gaagagaagt	aaggaaggtc	atagtttgga	gatggagaat	gagaatcttg	tagaaaatgg	720
tgcaactcc	gatgaagatg	ataacagctt	cctcaaacaa	caatctccac	aagaacccaa	780
gtctctgaat	tggtcgagtt	ttgtagacaa	cacctttgct	gaagaattca	ctactcagaa	840
tcagaaatcc	caggatgtgg	aactctggga	gggagaagtg	gtcaaagagc	tctctgtgga	900
agaacagata	aagagaaatc	ggtattatga	tgaggatgag	gatgaagagt	gacaaattgc	960
aatgatgctg	ggccttaaat	tcatgttagt	gttagcgagc	cactgccctt	tgtcaaaatg	1020
tgatgcacat	aagcagggtat	cccagcatga	aatgtaattt	acttggaagt	aactttggaa	1080
agaatttctt	tcttaaaatc	aaaaacaaaa	caaaaaaaca	caaaaaaacac	attctaaata	1140
ctagagataa	ctttacttaa	attcttcatt	ttagcagtga	tgatatgcat	aagtgtgtga	1200
aggcttgtaa	ctggggaaat	attccacctg	ataatagccc	agattctact	gtattcccaa	1260

aaggcaatat	taaggtagay	agatgattag	tagtatattg	ttacacacta	ttttggaatt	1320
agagaacata	cagaaggaat	ttaggggctt	aaacattacg	actgaatgca	cttttagtata	1380
aagggcacag	tttgtatatt	tttaaataaa	taccaattta	atTTTTtagt	atttacctgt	1440
taagagatta	tttagtcttt	aaatTTTTta	ggTTaatttt	cttgctgtga	tatatatgag	1500
gaatttacta	ctttatgtcc	tgctctctaa	actacatcct	gaactcgacg	tcctgaggta	1560
taatacaaca	gagcactttt	tgaggcaatt	gaaaaaccaa	cctacactct	tcggtgctta	1620
gagagatctg	ctgtctccca	aataagcttt	tgtatctgcc	agtgaattta	cygtactcca	1680
aatgattgct	ttcttttctg	gtgatattctg	tgcttctcat	aattactgaa	agctgcaata	1740
ttttagtaat	accttcggga	tactgtccc	ccatcttccg	tgttagagca	aagtgaagag	1800
tttaaaggag	gaagaagaaa	gaactgtctt	acaccacttg	agctcagacc	tctaaaccct	1860
gtatttccct	tatgatgtcc	cctttttgag	acactaattt	ttaaatactt	actagctctg	1920
aaatatattg	atTTTTatca	cagtattctc	agggTgaaat	taaaccaact	ataggccttt	1980
ttcttgggat	gattttctag	tcttaagggt	tggggacatt	ataaacttga	gtacatttgt	2040
tgacacacagt	tgatattcca	aattgtatgg	atgggaggga	gaggtgtctt	aagctgtagg	2100
cttttctttg	tactgcattt	atagagattt	agctttaata	tttttttagag	atgtaaaaca	2160
ttctgctttc	ttagtcttac	ctagtctgaa	acatttttat	tcaataaaga	ttttaattaa	2220
aattttraaaa	aaaaaaaaaa	aaactcgagg	gggggcccgt	acccaac		2267

&lt;210&gt; 52

&lt;211&gt; 2298

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

caggaactcc	aggttctgct	ggccgtggca	tcctctctcc	aagtctgctc	ccttaccgga	60
gctacgataa	cgtagcatga	atgacacctg	agattagagg	ctggggctca	ctgcaggctg	120
tggagagtca	tgctggtcca	catgaacact	tggcagtgtc	ctcgtagacc	cctcggtgat	180
gtggaatgga	cagggtgcctc	gcaagagaag	caagcacgtt	cataacaaaa	cagcaacaca	240
aagacatgtt	aagcatgttt	atTTatttgg	ctgtttttgt	ttatttactc	gagctgtggt	300
cacagctgcc	aggTacctaa	gcaagtcagt	tgggtacagc	aggacacgcc	accattccag	360
ggtactggta	ccgccagaaa	caggagtggg	tcttgtctctg	ttgcaggcac	actgcagtgg	420
ttttttcttc	ggcagctctc	caacaaacgc	ctgagtcaca	ggccagagct	gccttggtat	480
gttggttaagt	ccaaaacttc	ttctctgggc	tacctatctt	ccttcatgaa	gcagggtgctc	540
aggacccgga	agaatcatct	acctcccagc	tttgtgagac	agaaccaagt	aaaaggaaac	600
atgctagaaa	acgtgcctag	agaagacact	tcaacctttg	ccttatccaa	ccccctctca	660
gagaaagggtg	tcccatggcc	ccaaaaagaa	ctgccaaagt	ttggtgagga	gtaacaccct	720
ggcatgacat	tccttctctt	tcctggccct	caaccacttc	cttcctttgg	ctcttaagac	780
ctagcagggt	ctgtgaactc	tcaggccttg	gccagcacta	gttaggggag	gtcagggtgt	840
caatgtcctg	gtgattttat	gagactgccc	cactgagaaa	acttacttac	ttcaggcatc	900
cagtgccttc	acccagggtt	caggccctgt	ctaagggtgt	gcttaaagac	aaaaaggcaa	960
catgtgcctc	actggtgggtg	tgccactgtt	ctcatgtctc	ctcctaagtg	actccgattt	1020
tcagccctgg	tagaataagg	aagacagctg	atgctctctt	agcccttag	cacatgttcc	1080
taagggtgtgt	tgtcaagcca	acctgaagtt	ctgcctccct	gttatagtcc	ctgtctcccc	1140
cacagagacc	tgtgggtgct	cccagcagag	ttgagactgg	ctccgttgag	ttaatgacta	1200
gaatatagtg	ctttcactac	ttgattgtta	acctgttttc	ttctgatgcc	atcaagtacc	1260
agcagtcaga	ctattccact	ggTTaagtgt	ttactaccat	taaagcgagg	catgaagcaa	1320
agagctgagt	gagtcctctg	ctctccagag	gaccaagaaa	tacctgtgtg	acacagaccc	1380
acttcagtgt	gtacagcaaa	ttctatagtg	cttctgagcc	cagcagggtc	ttacctgccc	1440
ctggagagtt	ttagccgtct	tgtgtttctt	gtttacttca	caaccaaatt	tgtccctctc	1500
tctctctgtt	aaggggagaga	agtcacttta	gctggataat	acctatgtaa	caaactgagc	1560
agctgttatt	tgggcaaaat	caaagggaaga	aagagactat	ggtcttctat	ttattgtggg	1620
aaggaaaaca	gggtggggcg	ggtgagtga	aagggtgaaa	tccttggtac	cttgccctgt	1680
ggttacacag	tttaaccata	ggccaatttt	aggggcctct	gaagtatctt	tctacaaacg	1740
cagacaagct	ccactacccc	taacctgcca	ggatgtctaa	gtccactgtc	acaatccctt	1800



tcatacctcc	atggatttta	ctattgtgga	ggatatagca	agtgtccttg	gtctggctta	720
gtttgggagc	acagatggag	gaaacttagc	ctgcccatta	tacttcattg	tttggtagac	780
taagaaagta	actggtattg	ctgttttctg	agtaaccagt	tttctcaggc	tatagtattc	840
ttcctgggca	tcaacagtgg	gcactaagac	tttcacatgc	tatggccgaa	agtgccttaa	900
catttttctt	ggggaaattt	tatattagag	ctctcaactt	tttaaattag	gaagtgattt	960
tccctaaaac	agggaaataga	cctggcatga	tagccatctt	tgaaacttgc	cacagctctc	1020
ctcagcatac	ttggttgga	cgctgtacca	agatgctact	ggatactcaa	aaaatccctt	1080
ttctttttaa	ctctaaggaa	gggtattaaa	aaaacatccc	aacagtattt	aaaaacaaat	1140
taaagcttgg	gatcagtttt	aaaaagtgtc	atggctgggt	gtgatggctt	gcacttgtaa	1200
ttccagcact	tcgggaggcc	aagggtggaag	gactgcttga	gctcagctctg	ggctaaacat	1260
aacaagacac	tgtatcttaa	aaaaaaaaaa	aaaaaa			1296

&lt;210&gt; 55

&lt;211&gt; 1432

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (1399)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 55

gaattcggca	cgaggaaaga	ttgaagatgt	gggctagtca	aaagtttact	gataatttct	60
cactcttcta	cagattgata	atattgtcct	gatgaaggac	tcagatacag	gccgctctaa	120
agggttatgg	ttcatcacgg	tgagtctcta	gtctttttaa	ttttttattt	tactttgggt	180
ttgtccactt	gtctgatttt	gcagcttagc	ttcatcctcc	tccttttagga	actgtctaaa	240
tgaccataaa	accctgggtgc	ctgaagcttg	gactagcctt	ctacccttg	agatgagtgc	300
attgcttgag	atcttggtct	tgctcctaca	ctctgtcagt	ggcctgggta	tgggggtggt	360
caggagttca	cccagcttcc	ctggtgctgc	aacttggctc	tttgggtaat	agtaaccagg	420
ctgcagctaa	aaggwtgggg	gtgtgagggg	sgttaggtat	gggcttttaa	agacatgctt	480
tatagaattg	atgtttctca	taacagggat	gggaatagga	aattataact	ccctctgggt	540
ctaccccat	tgaagcaatt	tctgcaccga	gaaggatcag	ttattaacgt	agcactatgg	600
ggagaatagt	gaggccacct	aattatgggc	aagcttcacc	ttttcctgac	attccaacaa	660
aatgggttgc	aattcctata	actggttctt	ccagctccat	gtgactccag	gctgagaact	720
ggctgccagc	cacaaagtct	gatagaagct	tgtattttct	gggcttaaac	caggcagcat	780
acactcccac	agtgaccac	agggcagagg	gcagtagggt	gtattctgtc	attggaattg	840
ctcacctcaa	aaatatccag	taaaggcaag	ccatgtataa	cacctgccta	ggaactgtgc	900
agtaccacat	gccaggccct	aaggcaggta	atgctgctag	ctagctaaac	aagctagctg	960
tggtgttgac	aattctgtgg	tggcaagtaa	cttttgtaac	cttttctcgc	tctctgtgtg	1020
actgagatat	ggaaaggctt	ctgtggggca	tttttgccct	tgcattgttg	ccttttgggt	1080
caacaacctt	gacacttaaa	caaacagcag	actgggaatc	ctctttgtac	cagtgtgttg	1140
ctgggtgctg	ctgataaaaag	ggactagaga	gtaaaggcct	tctggtcagc	aggtcactta	1200
gtcaacagct	cttgtgtgta	tgtgggggtg	tgggttctgc	cttgctgtca	gcactagggg	1260
gtgttccctc	ttaagcctgg	aattaggagt	tccagattcc	tagkacttaa	ctaaaatttg	1320
gccaggcgcg	gtggctcaca	cctgtaatcc	cagcagtttg	ggaggccaag	gtgggtgaat	1380
cgcctgaggt	caggagttnng	agaccagcct	gaccaacgtg	gtgaaacccc	at	1432

&lt;210&gt; 56

&lt;211&gt; 979

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 56

ggcagcagga	gattgggtttc	atagtaagga	ttctttcttt	aaggtgatat	ggagcaactg	60
ctttcacagg	ggcaccaggt	tggtaaaaac	tcaaagctaa	caccattcag	aacttaccta	120
atgatgtttg	aaaaagaggc	ggtaggctga	taactgcttc	ctaggaagac	ctaccaactt	180
caccagcatg	ctcattactc	tcatcattgt	tgttggaaca	ttttgttctt	gggtacaagc	240
ttactttaca	cacatctggt	tactatgttt	gttctggaag	acggcagcct	catggagtga	300
ctagagaaca	cttgatactg	agctctggaa	atcgtttttt	aacagcttta	ttgatattgt	360
ataattttaca	gatcatcaag	ataagccctt	taaagtatac	aattcattgg	ttttttgaat	420
atctacagag	ttgtgtgacc	ataactatag	tccatttttt	aaacatttca	tcactccaaa	480
aggagacttt	gtacccatcc	tccctacctc	tctctttttc	tcccctagct	ctaggcagct	540
ttctttcttt	ttctttttct	ttttctcttt	ctttctttct	ttctttcttt	cttttctttc	600
ttttctttct	ttctttctct	ccttccttcc	ttccttctct	tccttccttc	cttccttctt	660
tccttccttc	cttccttctt	tcctttcttt	ctttctttct	ttctttcttt	cttttctttt	720
ttctttttct	ttctttcttt	cttgatttgc	ccattttgaa	catctcatat	gaatggaatc	780
atgcaatatg	tgagaatcct	gaaaattttg	atgttgatttt	caaagaccgt	gtagacaaac	840
atgacctggt	tactctgcag	gccccatctt	cccaacagtg	tcctgagtgt	gagattaaga	900
ccaaatgaac	tctgtcagtg	ctgccaactg	ttttattgga	ttaaatgact	tgtttttggc	960
caaaaaaaaa	aaaaaaaaaa					979

&lt;210&gt; 57

&lt;211&gt; 2667

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2652)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2659)

&lt;223&gt; n equals a,t,g, or c

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2662)

&lt;223&gt; n equals a,t,g, or c

&lt;400&gt; 57

gcaaagactt	ctgggtactct	gtgggtcaacg	gggtcatctt	taactgcttg	gccgtgcttg	60
ccctgtcatc	ccacctgaga	accatgctca	ccgaccctgg	ggcagtaccc	aaaggaaacg	120
ctacgaaaga	atacatggga	gagcttgacg	ctggaagccc	ggggaagtca	tctacaagtg	180
ccccaaagtgc	tgtgttatta	aacccgagcg	cgcccaccac	tgcagtattt	gcaaaagatg	240
tatttcggaaa	atggatcatc	actgcccgtg	gggtgaacaat	tgtgtaggag	aaaagaatca	300
aagatttttt	gtgtctcttca	ctatgtatat	agctctgtct	tcagtccatg	ctctgatcct	360
ttgtggattt	cagttcatct	cctgtgtccg	agggcagtg	actgaatgca	gtgatttttc	420
acctccgata	actgtaatcc	tgttgatctt	cctgtgcctt	gagggctctc	tgtttttcac	480
tttcaactggc	agttatgttt	ggcacccaaa	tccactccat	atgcaacgac	gagacggaga	540
tcgagcgatt	gaaaagtgag	aagcccacat	gggagcggag	gctgcgatgg	gaagggatga	600
agtcctgtct	tggggggccc	ccctcamtcc	tctggatgaa	tccctttgtg	ggcttccgat	660
ttagggcgact	gcccacgaga	cccagaaaag	gtggcccggga	gttctcagtg	tgaggcggtg	720
ctcatcgagc	tgaaacttgc	tcacagactt	ccagttattt	atttggggtc	tgaaggatat	780
caacagctca	tctgtgacca	acagggcaac	tggaacctac	acaaaccaat	tgcttgcagc	840

aagcagagtt	ttatatattt	atagtcacag	atggcagagg	aagaggctct	cagtcacccac	900
ctgtacaaca	acggaaaggt	gtgtggccac	acgaagaagc	caaacgccgt	ggcctcctgc	960
agagctgggg	cttctgtgga	raatacttcg	ggttattaca	tgggttattc	aaatcctggg	1020
tcctgagctg	ctgtttccaa	tcatagaaga	aaacagtga	tccagtgaac	agggattctc	1080
caagcagtca	tttcaggggg	ctcctgctga	ccccgccact	cagcagtga	ctccccggat	1140
cacagcaggg	cgtttacata	gaaagacgtt	ttggtctcga	ttagctccga	tgctttgcac	1200
tgaagttgca	aaagatctgt	gcactgaaca	gtgaagggtg	cttccggcac	actccccgct	1260
gccccggaag	agacatcctt	tgaccctctc	agcaagtctg	tgtgtgtgcg	tgtctgtgcg	1320
tgtgctgctg	tgtgtgcatg	tgtgtcaaaa	ttgccagtgt	tgcttaggca	atgtaacatt	1380
taccggctgt	gtacagcaaa	caagctatatt	tttagaaacc	gacgtttcag	ggaagagggg	1440
agagagccgc	ggggctcctgc	ccgtgggttac	tatgaatgta	ttgctgttgg	aggacatctc	1500
gatccaaaga	acagccgttc	ctgtgcggcc	cttcgttgcc	ctcctgcttt	catttttttaa	1560
agaaatcttg	agtgccttgag	ggccttgga	ctgatttttt	ttttttgttc	cagccaaatt	1620
agcagtgtat	aaatggcacc	taggtaagag	cagagctgcg	gctcggtagc	ttgatacttg	1680
gggcagcccg	atgctgtgtg	tggggcaggg	gaggcatcct	tactggagag	gcagggccca	1740
gccattgggc	acctctggga	aggggagggg	accatgaggc	agccagcccc	tggcaggggc	1800
gactgtgcca	ccgcaggcag	cgctccagtc	gggaatggcc	aggatggcgc	cctcttgttg	1860
gagtttttgg	ttagctttta	cgttttcttc	tccacccacg	gcacaggtga	taaaatagga	1920
tccttggtgc	ggagcttaaa	attatgccag	aaagccaaca	gctcccctcg	tggggccttg	1980
ccttaaaactt	gcctgggttg	tacatttttt	gccggacgca	tcaagaagca	atctgtgaca	2040
aagtctgagg	gtcttccttt	atgcttgccc	tccacactaa	gagaagttag	cgtctccctc	2100
ctgggaattg	ttttgccttt	ctgttcacat	gtgaactgtt	ttttgttttt	aattactctg	2160
taccccatcc	gaatcagggc	ttctaccact	gctgatgcaa	aaccacaaa	gggacctacc	2220
tgagccaccg	tcctagccaa	gcgagcaaac	ctgcaggggg	tttggaagtg	gacttggtca	2280
ccgcagaagc	gtgtgcgccg	ttgggggaag	agctgcgtca	cagccagagg	gacaaagtgt	2340
gggtgatcct	ggagacgcca	gtttccgaga	ttgttctgca	tattcatttg	cacattgttg	2400
tctgggttgg	acatgcgtgt	gggcttcagt	gtgaggcttt	taatatgtat	atcctgttat	2460
caataaaaca	attatccaag	tggttgaatc	ctgtgagact	tggcaagtgt	gtgcaaatca	2520
agtatacttg	acttttcaac	ctcttctttc	aatgtaaactt	ttatatgaaa	taaagtaatc	2580
aattaacagt	tctcaaaaaa	aaaaaaaaaa	aaaaaaaaaa	accccggggg	gggggccccg	2640
ggaacccaat	tnccccccna	angggggg				2667

&lt;210&gt; 58

&lt;211&gt; 1356

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 58

ggcacagagat	cccactcaat	cttacctgta	ctatttgcca	tctcctctca	atcaagttca	60
ttccataatg	atgttttccct	gagtaaacca	gaaaaaaaaa	aatccttata	aataggcaaa	120
gattacttag	taaaataggc	aaaggctctc	agagcctttt	ttctcatgtg	acatgcatgt	180
atctgaatag	gaacttcttg	tttctggctg	ttgaattatc	caaaaggatg	gctttagtgg	240
taagttttgt	gcataaagga	aagggaaggca	tgttagccag	tggagaatac	atttaggccca	300
cagggacctt	gtggtgcttc	ttggaattac	gctgggcttt	atgcatggat	agcagatgag	360
gccagggggag	gaaaaactct	agagatggaa	ggatggatgg	acaaaggaag	aagagaagaa	420
agagagaata	aatagcatat	gcaatatgtt	gaatttgac	tgctagcaag	caagaactta	480
agtcacacac	ataccagag	ccaggccctc	aggaatattt	ctcaaccttt	atgctttact	540
cttctccaga	tagtaaaaca	cctacaaaat	atcatcaggc	cttggttcttc	agcatgacaa	600
aatgtcctgc	ttcttgcttc	tgtgaacaag	ggggactgag	attgtatctc	ctttccacaa	660
gagcccttgg	gatcaacacc	tatggaaggg	aagggaggga	tggaaaaaag	gggagctgag	720
ctgggatgca	gatctgctag	atgcctcaga	caaccctgcg	gggagttatg	gagatggaat	780
gcccttttaga	attgtccagg	gttcaggaga	caggggaggc	ctttttatac	ctgcatcacc	840
tagtcactgg	gtatgcttga	tggggaagaa	tcatgacgtt	aggagagggg	ggtggctctc	900
tgaagctgta	gcaatcccca	aagggggcag	acctttgcag	atgaggtctt	tcttctggca	960

```

gcactccag aaactaaggt aaagtctctg ttattcctga gggggatccg gatggcttat 1020
catgtccatc acactccaca atgctatata ataccataaa actttgcaaa tagaatgaag 1080
gagaaaaataa atattgcaaa agcatactgt caaatttcta cataacagaa tattagggtt 1140
gaaaaatatta cccattagtt aaatttctcc tccttttaca atgatgtttg cctttaaagg 1200
ggtcaccaga gaataatttg ttccacagag cagccatgct attaatacta ctgttttagag 1260
atatagaaac gactttgcaa ctgtgaattg ctaagaattg tagaaataaa gggaaaaataa 1320
aattttcttt tctaaaaaaa aaaaaaaaaa aaaaaa 1356

```

<210> 59  
 <211> 850  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (18)  
 <223> n equals a,t,g, or c

```

<400> 59
gctgacctca tgatctgngc gcctcggcct cccaaagtgc taggwttaca ggcgtgagca 60
ctgcgtgctt atttaaagta tgtstgtgtc ctctccttat gaatgttttt ttttttcctt 120
gttggtcact cattttcatg atcgttttcc aattctttaga tctgacttca tgcattgagaa 180
gtacagaaag tacacaagaa taaagaataa aagtagggga gtcaggaatg ttctatttta 240
aacgcaacta ttagattcct cactgtctcc ctcaattat tcctgatctt tactctttga 300
gtcctccctg aaataacatg ctattagatt gatgcaaaag taattgtcgt ttttgccatt 360
actttttttt ttttcttttc tttttttaga cggagtttca ctcttattgc ccaggctgga 420
gtgcaatggt acaatctcgg ctactacaa cctctgcctc ctgggttcaa gcgattctcc 480
tgcctcagcc tcccagtag ctgggattat aggcattgcg caccacacc agctaatttt 540
gtatttttag tagagatggg gtttctccat gttggtcagg ctgctctcga actcctgatc 600
tcaggtggtc ctcccgctt ggctcccaa agtgctggga ttacaggcat gagccaccac 660
atccttgcca ttactttcaa tagcaaaaac cacaattact ttttcaccaa tctatataat 720
gattagaaat atatcccta ctctggaggg tgaggcaagt gatectatcg cgtgaacca 780
acagtttgag gccagcctgg gcaacatggg gagattttgc ctcaagaaaa aaaaaaaaaa 840
aaaactcgag 850

```

<210> 60  
 <211> 110  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (110)  
 <223> Xaa equals stop translation

```

<400> 60
Met Val Phe Leu Pro Ala His Ser Gln Pro Phe Leu Phe Pro Phe Leu
  1                      5                      10                      15

Phe Phe Phe Ser Pro His Thr Phe Leu Gly Leu Pro Pro Trp Lys Glu
                20                      25                      30

Glu Gly Leu Thr Trp Val Leu Ser Ser Pro Gln Val Arg Arg Val Thr

```

35                                      40                                      45  
 Arg Ala Pro Ser Leu Trp Thr Trp Ala Leu Pro Glu Pro Leu Pro Gln  
     50                                      55                                      60  
 Leu Pro Gln Leu Arg Pro Ala Gly Pro Gly His Gly Leu Arg Lys Ser  
     65                                      70                                      75                                      80  
 Pro Ala Pro Asn Pro Gly Arg Asn Trp Pro Leu Leu Gly Ser Leu Phe  
                                     85                                      90                                      95  
 Asp Phe Phe Arg Lys Met Ile Ser Ile Ser Phe Gln Pro Xaa  
                                     100                                      105                                      110

<210> 61  
 <211> 27  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (27)  
 <223> Xaa equals stop translation

<400> 61  
 Met His Pro Asp Leu Trp Pro Asp Cys Gly Leu Trp Leu Pro Gln Ala  
     1                                      5                                      10                                      15  
 Phe Ser Ser Leu His Trp Val Phe Leu His Xaa  
                                     20                                      25

<210> 62  
 <211> 37  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (37)  
 <223> Xaa equals stop translation

<400> 62  
 Met Ile Arg Tyr Trp Gly Phe Gly Gly Thr Gln Thr Leu Ala Ile Leu  
     1                                      5                                      10                                      15  
 Cys Val Pro Leu Asp Gln Ser Pro Lys Arg Arg Gly Ala Gly Arg Lys  
                                     20                                      25                                      30  
 Glu Trp Gly Ser Xaa  
                                     35

<210> 63



<211> 64  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (64)  
 <223> Xaa equals stop translation

<400> 63

Met	Phe	Tyr	Ile	Leu	Phe	Ile	Cys	Leu	Gly	Ser	Arg	Val	Leu	Asn	Leu
1				5					10					15	
Glu	Arg	Ser	Thr	Ser	Ile	Glu	Thr	Tyr	Gly	Ser	Cys	Ser	Leu	Glu	Ser
			20					25					30		
Gln	Trp	Arg	Leu	Ile	Ala	Val	Leu	His	Met	Asn	Ser	Asn	Leu	Thr	Leu
			35				40					45			
Asn	His	Gln	Gln	Thr	Leu	Ser	Phe	His	Gln	Asp	Val	Asp	Lys	Glu	Xaa
	50					55					60				

<210> 64  
 <211> 6  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (6)  
 <223> Xaa equals stop translation

<400> 64

Ser	Tyr	Cys	Ser	Val	Xaa
1				5	

<210> 65  
 <211> 31  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (31)  
 <223> Xaa equals stop translation

<400> 65

Met	Lys	Leu	Glu	Ala	Phe	Ile	Asp	Phe	Cys	Cys	Phe	Leu	Val	Val	Leu
1				5					10				15		

Thr Trp Ile Gln Val Val Leu Ile His Leu Phe Cys Leu Lys Xaa  
                   20                  25                  30

<210> 66  
 <211> 11  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (11)  
 <223> Xaa equals stop translation

<400> 66  
 Met Leu Leu Leu Asp Ser Asp Val Trp Phe Xaa  
       1                  5                  10

<210> 67  
 <211> 15  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (15)  
 <223> Xaa equals stop translation

<400> 67  
 Met Pro Trp Leu Gln Gln Leu Val Ser Phe Gly Leu Met Ser Xaa  
       1                  5                  10                  15

<210> 68  
 <211> 42  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (42)  
 <223> Xaa equals stop translation

<400> 68  
 Glu Ser Leu Leu Leu Val Ile Ala Phe Leu Phe Phe Cys Ser Phe Ser  
       1                  5                  10                  15

Phe Pro Ser Ile Lys Asp Leu Leu Tyr Ala Phe Arg Lys Ile Thr Lys  
                   20                  25                  30

Ala Glu Cys Leu Leu Thr Ile Tyr Gly Xaa  
           35                  40

<210> 69  
 <211> 16  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (16)  
 <223> Xaa equals stop translation

<400> 69  
 Lys Asn Cys Tyr Pro Lys Gly Phe Leu Arg Ala Asn Cys Leu Leu Xaa  
           1                          5                          10                          15

<210> 70  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (41)  
 <223> Xaa equals stop translation

<400> 70  
 Met Lys Ser Thr Asn Phe Leu Lys Gln Ala Leu Phe Phe Phe Phe Trp  
           1                          5                          10                          15

Lys Thr Cys Val Val Cys Pro Val Tyr Leu Phe Leu Met Arg Arg Pro  
                           20                          25                          30

Ala Phe Cys Gly Pro Arg Glu His Xaa  
                           35                          40

<210> 71  
 <211> 38  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (38)  
 <223> Xaa equals stop translation

<400> 71  
 Asn Ser Leu Lys Thr Ile Ile Phe Ala Leu Lys Thr Val Val Phe Leu  
           1                          5                          10                          15

Asp Leu Pro Val Tyr Ala His Ser Leu Leu Trp His Leu Tyr Ser Tyr  
                           20                          25                          30

Cys Asn Ala Tyr Ser Xaa  
35

<210> 72  
<211> 56  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (56)  
<223> Xaa equals stop translation

<400> 72  
Met Thr Phe Ser Pro Ala Leu Pro Pro Leu Arg Ser Pro Cys Ser Glu  
1 5 10 15  
Leu Leu Leu Trp Arg Tyr Pro Gly Ser Leu Ile Pro Glu Ala Leu Arg  
20 25 30  
Leu Leu Arg Leu Gly Asp Thr Pro Ser Pro Pro Tyr Pro Ala Thr Pro  
35 40 45  
Ala Gly Asp Ile Met Glu Leu Xaa  
50 55

<210> 73  
<211> 20  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (20)  
<223> Xaa equals stop translation

<400> 73  
Gly Met Leu His Asp Gly Gln Leu Leu Pro Ser Leu Val Ile Ile His  
1 5 10 15  
Cys Ser Cys Xaa  
20

<210> 74  
<211> 30  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (30)

<223> Xaa equals stop translation

<400> 74

Leu Glu Leu Leu Val Cys Asn Tyr Val Tyr Gly Asn Gly Val Val Phe  
1 5 10 15

Met Ile Cys Ile Tyr Ile Ile Ile Trp Met Tyr Ile Tyr Xaa  
20 25 30

<210> 75

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 75

Trp Pro Leu Leu Pro Gln Val Leu Gln Glu Arg Val Trp Trp Ala His  
1 5 10 15

Leu Gln Ser Gln Leu Leu Gly Lys Leu Gln Gly Leu Leu Glu Pro Arg  
20 25 30

Lys Ser Gln Leu Ala Gly His Leu Glu Leu Pro Lys Asn Thr Leu Ser  
35 40 45

Arg Lys Xaa  
50

<210> 76

<211> 119

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (119)

<223> Xaa equals stop translation

<400> 76

Met Pro Phe Thr Phe Ser Trp Cys Phe Ser Leu Trp Thr Leu Leu Asn  
1 5 10 15

His Arg Pro Asp Lys Ser Leu Gln Arg Lys Gly Leu Trp Ala Val Pro  
20 25 30

Glu Ala Gly Trp Gly Gln Asp Asn Thr Ala Ala Ser Pro Gly Cys Pro  
35 40 45

Leu Trp Gly Pro Leu Gly Ala Gly Lys Gly Ile Gly Gln Arg Gly Ser

50					55					60					
Gly 65	Ala	Gly	His	Glu	Gly 70	Pro	Pro	Ser	Ala	Leu 75	Trp	Leu	Gln	Gly	Trp 80
Arg	Trp	Phe	Cys	Pro 85	Pro	Ser	Leu	Pro	Ser 90	Gly	Gly	Ala	Gly	Gly 95	Ser
Ala	Ala	Gly	Val 100	Gly	Thr	Ser	Pro	Arg	Met 105	Ala	Val	Glu	Gly 110	Val	Arg
Thr	Val	Leu	Gly	Ser	Pro	Xaa									
115															

```
<210> 77
<211> 37
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (37)  
<223> Xaa equals stop translation
```

```

<400> 77
Met Phe Leu Leu Thr Ile Gly Lys Ile Glu Lys Ala Leu Cys Phe Leu
 1             5             10             15
Phe Phe Cys Cys Cys Cys His Cys Cys Phe Phe Gln Lys Thr Ser Val
          20             25             30
Ser Val Leu Ser Xaa
      35

```

```
<210> 78
<211> 42
<212> PRT
<213> Homo sapiens
```

```
<220>
<221> SITE
<222> (42)
<223> Xaa equals stop translation
```

```

<400> 78
Asp Phe Leu Phe Phe Phe Pro Pro Cys Gly Ser His Cys Phe Leu Ser
 1             5             10             15
Cys Leu Tyr Leu Ser Leu Cys Leu Ser Ser Ser Phe Pro Gln Met Ser
      20             25             30
Gln Leu Ser Leu Pro Asn Phe Pro Ile Xaa
      35             40

```

<210> 79  
 <211> 143  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (143)  
 <223> Xaa equals stop translation

<400> 79

Met	Cys	Ala	Pro	Ala	Phe	Pro	Ala	Trp	Asn	Val	Phe	Trp	Leu	Ser	Cys
1				5				10						15	
Ser	Asn	Ile	Cys	Glu	Pro	Ser	Val	Cys	Pro	Val	Ser	Leu	Gly	Ser	Leu
			20					25					30		
Gln	Val	Gln	Glu	His	Thr	Cys	Arg	Ala	Gln	His	Asp	Asp	Gly	Val	Lys
			35				40					45			
Gly	Arg	Lys	Gln	Ser	Ser	Glu	Gly	Val	Gly	Arg	Trp	Ala	Val	Cys	Pro
	50					55					60				
Arg	Gln	Val	Pro	Ser	Pro	Pro	Ala	Ser	Phe	Leu	Arg	Pro	Gly	Arg	Ala
65					70					75					80
Ala	Met	Val	Leu	Ala	Ala	Ser	Pro	Ser	Val	Leu	Thr	Val	Ser	Leu	His
				85					90					95	
Leu	Gly	Pro	Pro	Glu	Ser	Gln	Val	Ser	Phe	Ser	Ser	Asn	Leu	Thr	Arg
			100					105					110		
Glu	Lys	Lys	His	Gly	Cys	Ala	Trp	Pro	Thr	Gly	Pro	Gly	Gly	Asp	Gly
		115					120					125			
Pro	Pro	Arg	Ser	Leu	Lys	Leu	Trp	Met	Ala	Ala	Val	Leu	Tyr	Xaa	
	130					135						140			

<210> 80  
 <211> 44  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (44)  
 <223> Xaa equals stop translation

<400> 80

Met	Met	Leu	Leu	Asn	Ser	Ile	Gln	Val	Leu	Leu	Tyr	Val	Pro	Ile	Val
1				5					10					15	

Phe Arg Phe Leu Thr Asp Cys Cys Gly Cys Thr Ser Leu Ser Asp Cys  
                   20                                  25                                  30

Ser Leu Ile Tyr Cys Arg Thr Leu Gly Cys Phe Xaa  
                   35                                  40

<210> 81  
 <211> 37  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (37)  
 <223> Xaa equals stop translation

<400> 81  
 Met Cys Ile Leu Phe Cys Thr Pro Ile Ser Leu Phe Leu Cys Trp Leu  
       1                                  5                                  10                                  15

Pro Gly Pro Ser Leu Glu Pro Trp Val Glu Arg Leu Leu Lys Asp His  
                   20                                  25                                  30

His Arg Gln Pro Xaa  
                   35

<210> 82  
 <211> 43  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (43)  
 <223> Xaa equals stop translation

<400> 82  
 Met Met Asn Val Ser Leu Glu Ile Tyr Phe Val Val Phe Leu Ser Leu  
       1                                  5                                  10                                  15

Phe Cys Val Val Leu Pro Leu His Ala Leu Phe Leu Lys Ser Phe Phe  
                   20                                  25                                  30

Phe Leu Asp Phe Ala Cys Phe Gln Ile Leu Xaa  
                   35                                  40

<210> 83  
 <211> 82  
 <212> PRT  
 <213> Homo sapiens

<220>



&lt;221&gt; SITE

&lt;222&gt; (82)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 83

Met His Pro Phe Phe Pro Gln Arg Gly Trp Thr Gln Val Ser Leu Gly  
 1 5 10 15

Val Gly Ala Pro Ala Leu Cys His Arg His Gly Glu Gly Phe Leu Leu  
 20 25 30

Thr Pro Ser Ala Leu Pro Thr Trp Val Val Leu Ser Lys Lys Val Ile  
 35 40 45

Pro Pro Ala Leu Ala Pro Phe Pro Arg Thr Lys His Val Asp His Val  
 50 55 60

Gln Tyr Phe Leu Leu Cys Arg Glu Ala Ala Met Ser Glu Ile Lys Ala  
 65 70 75 80

Val Xaa

&lt;210&gt; 84

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (8)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 84

Met Thr Ile Phe Glu Pro Ser Xaa  
 1 5

&lt;210&gt; 85

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (18)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 85

Met Ser Thr Gly His Gly Ala Cys Leu Val Ser Leu Pro Val Val Gln  
 1 5 10 15

Ala Xaa

<210> 86  
 <211> 8  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (8)  
 <223> Xaa equals stop translation

<400> 86  
 Met Ile Cys Leu Ala Leu Tyr Xaa  
       1                              5

<210> 87  
 <211> 58  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (39)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (58)  
 <223> Xaa equals stop translation

<400> 87  
 Met Thr Gly Leu Tyr Ala Val Gly Arg Gly Ala Trp Thr Ala Gly Leu  
       1                              5                              10                              15

Cys Ala Val Gly Trp Gly Gly Leu Asp Val Trp Leu Leu Cys Gly Ala  
                               20                              25                              30

Gly Ala Gly Leu Gly Pro Xaa Gly Arg Leu Glu Gly Gln Pro Gly Pro  
                               35                              40                              45

Val Val Ser Leu Val Arg Asn Arg Lys Xaa  
       50                              55

<210> 88  
 <211> 35  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (35)  
 <223> Xaa equals stop translation

&lt;400&gt; 88

Met Gln Cys Asn Phe Leu Ser Ser Leu Glu Arg Trp Gly His Ile Pro  
 1 5 10 15

Leu Gly Cys Phe Leu Leu Cys Val Ile Leu Ile Ser Leu Ala Gly Lys  
 20 25 30

Gln Lys Xaa  
 35

&lt;210&gt; 89

&lt;211&gt; 70

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (70)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 89

Met Val Pro Thr Pro Pro Gly Trp Val Gly Phe Phe Val Phe Trp Val  
 1 5 10 15

Phe Leu Gly Gly Gly Leu Phe His Val Thr Ser Ile Ser Val Phe Ile  
 20 25 30

Ser Tyr Leu Phe His Phe Gln Val Tyr His Gly Lys Cys Thr Asp Phe  
 35 40 45

Phe Val Asn Asn Val Leu Gly Phe Ala Lys Lys Glu Lys Lys Lys Thr  
 50 55 60

Leu Leu Ser Leu Pro Xaa  
 65 70

&lt;210&gt; 90

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (21)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 90

Met Thr Thr Phe Leu Leu Leu Leu Leu Val Ser Ile Phe Ser Ser  
 1 5 10 15

Val Asn Cys Gly Xaa  
 20

<210> 91  
 <211> 23  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (23)  
 <223> Xaa equals stop translation

<400> 91  
 Met Lys Phe Val Leu Phe Tyr Leu Cys Thr Gly Tyr Pro Leu Phe Trp  
           1                          5                          10                          15  
 Glu Cys Phe Phe Ile Leu Xaa  
                           20

<210> 92  
 <211> 16  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (16)  
 <223> Xaa equals stop translation

<400> 92  
 Asp Thr Ile Val Leu Ser Gly Cys Leu Cys Leu Cys Tyr Tyr Val Xaa  
           1                          5                          10                          15

<210> 93  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (41)  
 <223> Xaa equals stop translation

<400> 93  
 Met Lys Gly Val Leu Leu Ile Asn Leu Leu Tyr Tyr Leu Asp Thr Phe  
           1                          5                          10                          15  
 Val Tyr Ser Leu Gln Pro Thr Cys Ile Arg Gly Ile Gly Val Gly Leu  
                           20                          25                          30

Asn His Gln His Tyr Phe Ile Lys Xaa  
           35                          40

<210> 94  
 <211> 11  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (11)  
 <223> Xaa equals stop translation

<400> 94  
 Met Gly Thr Ser Ser Ile Leu Gln Met Lys Xaa  
       1                          5                          10

<210> 95  
 <211> 69  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (69)  
 <223> Xaa equals stop translation

<400> 95  
 Lys Leu Trp Gln Thr Met Cys Asp Ala Phe Phe Cys Ile Gly Cys Ser  
       1                          5                          10                          15

His Trp Ile Ala Gly Ser Gly Tyr Cys Val Ile Val Ala Ala Leu Gly  
                           20                          25                          30

Leu Ala Glu Gly Pro Leu Cys Leu Asp Ser Leu Gly Gln Trp Asn Tyr  
           35                          40                          45

Thr Phe Ala Gln His Arg Gly Pro Val Pro Ser Gly Tyr Leu His Met  
       50                          55                          60

Val Arg Val His Xaa  
       65

<210> 96  
 <211> 27  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (27)  
 <223> Xaa equals stop translation

&lt;400&gt; 96

Met Ala Leu Thr Gly Ala Phe Ile Val Lys Ser Tyr Ser Arg Gly Trp  
1 5 10 15

Ser Cys Gly Cys Val Cys Ser Val Pro Ile Xaa  
20 25

&lt;210&gt; 97

&lt;211&gt; 11

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (11)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 97

Met Leu Trp Ile Phe Lys Trp Gln Val Leu Xaa  
1 5 10

&lt;210&gt; 98

&lt;211&gt; 196

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (196)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 98

Met Gly Met Trp Ser Ile Gly Ala Gly Ala Leu Gly Ala Ala Ala Leu  
1 5 10 15

Ala Leu Leu Leu Ala Asn Thr Asp Val Phe Leu Ser Lys Pro Gln Lys  
20 25 30

Ala Ala Leu Glu Tyr Leu Glu Asp Ile Asp Leu Lys Thr Leu Glu Lys  
35 40 45

Glu Pro Arg Thr Phe Lys Ala Lys Glu Leu Trp Glu Lys Asn Gly Ala  
50 55 60

Val Ile Met Ala Val Arg Arg Pro Gly Cys Phe Leu Cys Arg Glu Glu  
65 70 75 80

Ala Ala Asp Leu Ser Ser Leu Lys Ser Met Leu Asp Gln Leu Gly Val  
85 90 95

Pro Leu Tyr Ala Val Val Lys Glu His Ile Arg Thr Glu Val Lys Asp  
100 105 110

Phe Gln Pro Tyr Phe Lys Gly Glu Ile Phe Leu Asp Glu Lys Lys Lys  
           115                    120                    125  
 Phe Tyr Gly Pro Gln Arg Arg Lys Met Met Phe Met Gly Phe Ile Arg  
           130                    135                    140  
 Leu Gly Val Trp Tyr Asn Phe Phe Arg Ala Trp Asn Gly Gly Phe Ser  
 145                    150                    155                    160  
 Gly Asn Leu Glu Gly Glu Gly Ser Ser Leu Gly Glu Phe Ser Trp Trp  
                     165                    170                    175  
 Asp Gln Glu Ser Arg Ala Phe Phe Leu Ser Thr Glu Lys Lys Asn Leu  
                     180                    185                    190  
 Gly Thr Lys Xaa  
           195

<210> 99  
 <211> 42  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (42)  
 <223> Xaa equals stop translation

<400> 99  
 Met Lys Cys Phe Leu Leu Leu Cys Cys Ala Ile Val Leu Cys Phe Ser  
   1                    5                    10                    15  
 Lys His Cys Lys Asn Ser Ile Phe Gly Leu Val Phe Trp Ile Ile Tyr  
                     20                    25                    30  
 Tyr Asn Leu Arg Ser Val Leu Leu Met Xaa  
           35                    40

<210> 100  
 <211> 34  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (34)  
 <223> Xaa equals stop translation

<400> 100  
 Met Pro Leu Leu Leu Arg Trp Val Ser Trp Leu Gln Val Trp Lys Pro  
   1                    5                    10                    15

Arg Pro Pro Leu Ser Arg Arg Arg Lys Thr Ser Gln Leu Lys Pro Arg  
                   20                  25                  30

Ser Xaa

<210> 101  
 <211> 22  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (22)  
 <223> Xaa equals stop translation

<400> 101  
 Met Phe Ile Tyr Leu Ala Val Phe Val Tyr Leu Leu Glu Leu Trp Ser  
      1                  5                  10                  15

Gln Leu Pro Gly Thr Xaa  
                   20

<210> 102  
 <211> 23  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (23)  
 <223> Xaa equals stop translation

<400> 102  
 Met Val Ile Phe Ile Ile Leu Leu Thr Cys Phe Gly Phe Ser Asn Gly  
      1                  5                  10                  15

Ser Phe Ser Phe Ser Leu Xaa  
                   20

<210> 103  
 <211> 30  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (30)  
 <223> Xaa equals stop translation

<400> 103  
 Met Ser Lys Pro Lys Lys Tyr Thr Ile Val Leu Val Leu Leu Pro Tyr



1 5 10 15

Arg Glu Ala Ile Gln Thr Val Gly Met Asn Leu Ser Tyr Xaa  
20 25 30

<210> 104  
<211> 39  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (39)  
<223> Xaa equals stop translation

<400> 104  
Met Ser Ala Leu Leu Glu Ile Leu Ala Leu Leu Leu His Ser Val Ser  
1 5 10 15

Gly Pro Gly Met Gly Val Phe Arg Ser Ser Pro Ser Phe Pro Gly Ala  
20 25 30

Ala Thr Trp Leu Phe Gly Xaa  
35

<210> 105  
<211> 39  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (39)  
<223> Xaa equals stop translation

<400> 105  
Met Leu Ile Thr Leu Ile Ile Val Val Gly Thr Phe Cys Ser Trp Val  
1 5 10 15

Gln Ala Tyr Phe Thr His Ile Trp Leu Leu Cys Leu Phe Trp Lys Thr  
20 25 30

Ala Ala Ser Trp Ser Asp Xaa  
35

<210> 106  
<211> 79  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE



&lt;210&gt; 109

&lt;211&gt; 49

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 109

Pro Gly Arg Pro Gly Ala Pro Gly Ser Pro Gly Leu Pro Gly Gln Ile  
 1 5 10 15

Gly Arg Glu Gly Arg Gln Gly Leu Pro Gly Val Arg Gly Leu Pro Gly  
 20 25 30

Thr Lys Gly Glu Lys Gly Asp Ile Gly Ile Gly Ile Ala Gly Glu Asn  
 35 40 45

Gly

&lt;210&gt; 110

&lt;211&gt; 49

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

Pro Ser Gln Pro Gly Trp Leu Pro Ser Pro Lys Arg Pro Gly Arg Ala  
 1 5 10 15

Gly Arg Gln Gly Leu Gln Val Leu Pro Ala Leu Arg Pro Arg Pro Gln  
 20 25 30

Ile Ser Asn Asn Glu Gln Thr Leu Gly Pro Ala Trp Ala Gly Asp Leu  
 35 40 45

Gly

&lt;210&gt; 111

&lt;211&gt; 93

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 111

Pro Gly Gln Pro Gly Pro Lys Gly Glu Gln Gly Pro Glu Gly Val Gly  
 1 5 10 15

Arg Pro Gly Lys Pro Gly Gln Pro Gly Leu Pro Gly Val Gln Gly Pro  
 20 25 30

Pro Gly Leu Lys Gly Val Gln Gly Glu Pro Gly Pro Pro Gly Arg Gly  
 35 40 45

Val Gln Gly Pro Gln Gly Glu Pro Gly Ala Pro Gly Leu Pro Gly Ile

50                      55                      60

Gln Gly Leu Pro Gly Pro Arg Gly Pro Pro Gly Pro Thr Gly Glu Lys  
65                      70                      75                      80

Gly Ala Gln Gly Ser Pro Gly Val Lys Gly Ala Thr Gly  
85                      90

```
<210> 112
<211> 100
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (36)  
<223> Xaa equals stop translation
```

```
<220>  
<221> SITE  
<222> (38)  
<223> Xaa equals stop translation
```

```

<400> 112
Pro His Leu Pro Arg Pro His Arg Ala Gln Asn Arg Arg Gly Leu Pro
  1                               10                      15

Gln Pro Ala Glu His Arg Val Ser Gly Ala Pro Ile His Arg Cys Arg
      20                               25                      30

Arg Gly Tyr Xaa Trp Xaa Pro Pro Cys Gly Phe Glu Gly Ser Ser Pro
      35                               40                      45

Trp Pro His Pro Ser Gln Pro Gly Trp Leu Pro Ser Pro Lys Arg Pro
      50                               55                      60

Gly Arg Ala Gly Arg Gln Gly Leu Gln Val Leu Pro Ala Leu Arg Pro
      65                               70                      75

Arg Pro Gln Ile Ser Asn Asn Glu Gln Thr Leu Gly Pro Ala Trp Ala
      85                               90                      95

Gly Asp Leu Gly
      100

```

```
<210> 113
<211> 49
<212> PRT
<213> Homo sapiens
```

<400> 113  
Gly Thr Asp Glu Gln Leu Leu Lys Ala Val Lys Ile Ile Lys Ile Leu  
1 5 10 15

Tyr Gln Asn Pro Tyr Pro Lys Pro Ala Gly Ser Arg Ala Ala Arg Arg  
                   20                  25                  30

Asn Lys Arg Arg Arg Trp Lys Lys Arg Gln Asp Gln Val Glu Gly Leu  
                   35                  40                  45

Ala

<210> 114

<211> 47

<212> PRT

<213> Homo sapiens

<400> 114

Gly Thr Lys Trp Ser Leu Leu Pro Glu Gln Lys Leu Ala Leu Ile Ala  
       1                  5                  10                  15

Ser Gln Asn Pro His Pro His Pro His Gly Ser Val Ala Gly Cys Arg  
                   20                  25                  30

Pro Ser Lys Trp Thr Pro Arg Asp Arg Ser Gly Asp Cys Leu Ala  
                   35                  40                  45

<210> 115

<211> 47

<212> PRT

<213> Homo sapiens

<400> 115

Arg Ser Ser Arg Ser Gly Arg Leu Ser Leu His Gly Asp Ile Arg Leu  
       1                  5                  10                  15

Leu Phe Ser Arg Arg Ser Leu Glu Leu Asp Thr Gly Leu Pro Tyr Glu  
                   20                  25                  30

Leu Gln Ala Val Thr Glu Ala Pro His Asn Pro Arg Tyr Ser Pro  
                   35                  40                  45

<210> 116

<211> 47

<212> PRT

<213> Homo sapiens

<400> 116

Gln Ser Ser Lys Ser Gly Lys Ile Tyr Leu His Arg Asp Val Arg Leu  
       1                  5                  10                  15

Leu Phe Ser Arg Lys Ser Met Glu Val Asp Ser Gly Ala Ala Tyr Glu  
                   20                  25                  30

Leu Lys Ser Tyr Thr Glu Ser Pro Thr Asn Pro Gln Phe Ser Pro  
                   35                                  40                                  45

<210> 117  
 <211> 41  
 <212> PRT  
 <213> Homo sapiens

<400> 117  
 Leu Tyr Phe Tyr Phe Tyr Phe Thr Ile Ile Phe Phe Ile Phe Cys Leu  
   1                                  5                                  10                                  15  
 Asn Ser Tyr Phe Leu Met Cys Asn Ile Phe Ile Tyr Ile Ile Leu Asn  
                                   20                                  25                                  30  
 Thr Ile Ala Leu Leu Thr Lys Val Phe  
                                   35                                  40

<210> 118  
 <211> 45  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (23)  
 <223> Xaa equals stop translation

<400> 118  
 Ile His Phe Phe Phe Phe Val Asp Ile Thr Phe Arg Ile Glu Leu Phe  
   1                                  5                                  10                                  15  
 Val Ile Tyr Cys Leu Asp Xaa Tyr Phe Val Tyr Leu Gly Ile Met Lys  
                                   20                                  25                                  30  
 Lys Val Cys Val Asn Ile Ala Leu Leu Lys His Met Phe  
                                   35                                  40                                  45

<210> 119  
 <211> 34  
 <212> PRT  
 <213> Homo sapiens

<400> 119  
 Thr Trp Phe Phe Gly Gln Ser Leu Cys Lys Val Ile Pro Tyr Leu Gln  
   1                                  5                                  10                                  15  
 Thr Val Ser Val Ser Val Ser Val Leu Thr Leu Ser Cys Ile Ala Leu  
                                   20                                  25                                  30

Asp Arg

<210> 120  
 <211> 36  
 <212> PRT  
 <213> Homo sapiens

<400> 120  
 Ser Trp Lys Ile Phe Phe Gly Lys Asn Val Ile Lys Ala Thr Pro Lys  
     1                    5                    10                    15  
 Ile Asp Thr Val Ser Gly Cys Thr Ile Leu Thr Val Ile Cys Leu His  
                     20                    25                    30  
 Val Phe Asp Arg  
                     35

<210> 121  
 <211> 50  
 <212> PRT  
 <213> Homo sapiens

<400> 121  
 His Cys Trp Trp Asp Cys Lys Leu Val Gln Pro Leu Trp Lys Thr Val  
     1                    5                    10                    15  
 Trp Arg Phe Leu Arg Asp Leu Gln Leu Glu Ile Pro Phe Asp Pro Ala  
                     20                    25                    30  
 Ile Pro Leu Leu Gly Ile Tyr Pro Lys Asp Tyr Lys Ser Cys Cys Tyr  
                     35                    40                    45  
 Lys Asp  
           50

<210> 122  
 <211> 45  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (40)  
 <223> Xaa equals stop translation

<400> 122  
 Asn Cys His Gly Lys Leu Glu Trp Tyr Arg Gln Leu Trp Lys Thr Ala  
     1                    5                    10                    15  
 Trp Gln Phe Val Thr Thr Tyr Asn Pro Ser Val Ser Leu Leu Gly Ile  
                     20                    25                    30  
 Tyr Leu Lys Glu Met Lys Thr Xaa Cys Ser His Lys Asn

35

40

45

&lt;210&gt; 123

&lt;211&gt; 49

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 123

Pro Gln Gly Pro Pro Gln Gln Glu Gly Asn Asn Pro Gln Gly Pro Pro  
 1 5 10 15

Pro Pro Ala Gly Gly Asn Pro Gln Gln Pro Gln Ala Pro Pro Ala Gly  
 20 25 30

Gln Pro Gln Gly Pro Pro Arg Pro Pro Gln Gly Asp Lys Pro Ser Arg  
 35 40 45

Pro

&lt;210&gt; 124

&lt;211&gt; 51

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (2)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 124

Pro Xaa Gly Pro Pro Ser Ala Glu Ala Gly Gly His Pro Gln Ser Pro  
 1 5 10 15

Leu Pro Cys Asn Pro Ser Trp Gly His Asn Gly Ala Leu Ser Ala Gly  
 20 25 30

Gly Gln Cys Pro Ser His Leu Pro Ser Ser Pro Gln Gln Lys Arg Pro  
 35 40 45

Ala Thr Pro  
 50

&lt;210&gt; 125

&lt;211&gt; 27

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 125

Thr Ile Val Pro His Ile His Pro Gln His Thr Thr Asn Val Asn His  
 1 5 10 15



Gln His Phe Gln His Val His Tyr Phe Pro His  
                   20                  25

<210> 126

<211> 28

<212> PRT

<213> Homo sapiens

<400> 126

Thr Leu Leu Ile Asn Ile His Pro Tyr Asp Asp Val Asp Thr Asn His  
   1                  5                  10                  15

Glu His Tyr Ser Ile Pro Ile His Ile Ile Ala His  
                   20                  25

<210> 127

<211> 57

<212> PRT

<213> Homo sapiens

<400> 127

Gly Phe Lys Arg Phe Ser Cys Leu Ser Leu Pro Ser Ser Trp Asp Tyr  
   1                  5                  10                  15

Arg His Val Pro Pro Arg Gln Val His Phe Val Phe Ser Val Glu Thr  
                   20                  25                  30

Gly Phe His Arg Ala Gly Gln Ala Gly Leu Glu Leu Leu Thr Ser Ser  
                   35                  40                  45

Val Pro Pro Thr Ser Ala Phe Pro Lys  
                   50                  55

<210> 128

<211> 54

<212> PRT

<213> Homo sapiens

<400> 128

Gly Pro Lys Gln Ser Ser Arg Leu Gly Leu Leu Gly Ser Trp Asp Gln  
   1                  5                  10                  15

Gly His Thr Pro Pro His Pro Ala Asn Phe Phe Val Glu Val Glu Ser  
                   20                  25                  30

Cys Tyr Val Ala Gln Ala Gly Leu Glu Leu Leu Gly Ser Ser Asp Pro  
                   35                  40                  45

Pro Thr Leu Ala Pro Lys  
                   50

<210> 129  
 <211> 91  
 <212> PRT  
 <213> Homo sapiens

<400> 129  
 Ser Asp Glu Glu Asp Ser Asp Gln Phe Ile Asp Glu Ala His Gln Gly  
   1                          5                          10                          15  
 Pro Pro Leu Gly Gly Gln Leu Ser Lys His Pro Ala Gly Asp Gly Asn  
                           20                          25                          30  
 Lys Asp Asp Gly Pro Gln Gln Lys Pro Pro His His Gly Gly His His  
                           35                          40                          45  
 His Arg Pro Pro Pro Pro Thr Gly Glu Pro Gln Asn Thr Gln Lys Pro  
                           50                          55                          60  
 Gly Gly His His His His Arg Pro Thr Pro Pro Pro Gly Lys Pro Glu  
   65                          70                          75                          80  
 Gly Ser Pro Pro Pro Ser Gln Gly Ser Arg Pro  
                           85                          90

<210> 130  
 <211> 83  
 <212> PRT  
 <213> Homo sapiens

<400> 130  
 Thr Glu Lys Glu Ala Gly Ala Gln Val Leu Met Pro Arg His Lys Gly  
   1                          5                          10                          15  
 Ser Pro Arg Gln Ala Glu Leu Ser Ser Pro Pro Gly Asp Gly Ala Gln  
                           20                          25                          30  
 Pro Gln His Ser Val Leu Gln Ser Gly Gly Ala Gln His Gly Ala Asn  
                           35                          40                          45  
 Pro Leu His Arg His Pro Trp Gly Ser Ser His Thr Ser Cys Arg Pro  
   50                          55                          60  
 Thr Ser Pro Thr Arg Gly Gln Arg Gly Arg Ala Lys Pro Ser Pro Ser  
   65                          70                          75                          80  
 Leu Lys Pro

<210> 131  
 <211> 46  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 131

Gly Asp Arg Gly Pro Arg Gly Glu Arg Gly Pro Pro Gly Pro Pro Gly  
 1 5 10 15

Arg Asp Gly Gly Asp Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Pro  
 20 25 30

Pro Gly Pro Pro Gly Leu Gly Gly Asn Phe Ala Ala Gln Tyr  
 35 40 45

&lt;210&gt; 132

&lt;211&gt; 45

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 132

Gly Asp Arg Gly Gly Glu Val Gly Pro Ala Ser His Pro Gly Asp Asp  
 1 5 10 15

Lys Gly Met Glu Lys Val Thr Pro Val Pro Lys Leu Gln Pro Pro Ile  
 20 25 30

Gly Trp Lys Gly Trp Gly Gly Lys Phe Ser Ala Arg Tyr  
 35 40 45

&lt;210&gt; 133

&lt;211&gt; 48

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 133

Leu Gly Gly Gly Gly Ala Gly Gly Cys Cys Pro Pro Ala Gln Pro  
 1 5 10 15

Gly Cys Gly Asn Pro Cys Gln Gly Ala Gly Pro Gln Tyr Ala Ala Ala  
 20 25 30

Pro Pro Pro Pro Pro Pro Gln Tyr Ala Ala Ala Pro Gln Gly Gly Ala  
 35 40 45

&lt;210&gt; 134

&lt;211&gt; 50

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (19)

&lt;223&gt; Xaa equals stop translation

<220>  
 <221> SITE  
 <222> (27)  
 <223> Xaa equals stop translation

<400> 134  
 Leu Gly Gly Ala Gly Ile Thr Ser Ala Arg Pro Cys Pro Ser Gln Glu  
           1                  5                  10                  15  
 Val Arg Xaa Gly Asp Thr Met Asp Gly Asp Xaa Gly Pro Lys Val Pro  
                   20                  25                  30  
 Ala Ser Leu Pro Pro Gln Lys Pro His Ser Gln Ala Ser Pro Gln Ser  
           35                  40                  45  
 Arg Ala  
       50

<210> 135  
 <211> 18  
 <212> PRT  
 <213> Homo sapiens

<400> 135  
 Gln Pro Gly Gly Ser Thr Ser Ser Arg Ala Ala Ala Thr Ala Val Glu  
           1                  5                  10                  15  
 Leu Gln

<210> 136  
 <211> 18  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (5)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>  
 <221> SITE  
 <222> (12)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 136  
 Gln Pro Gly Gly Xaa Thr Ser Ser Arg Ala Ala Xaa Asn Ala Val Glu  
           1                  5                  10                  15  
 Leu Gln

<210> 137  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 137  
Gly Asp Pro Leu Val Leu Glu Arg Pro  
1 5

<210> 138  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 138  
Gly Asp Pro Leu Val Leu Glu Arg Pro  
1 5

<210> 139  
<211> 18  
<212> PRT  
<213> Homo sapiens

<400> 139  
Tyr Ala Ser Ser Thr Val Ser Ile Ser Leu Ile Ser Asn Ser Cys Ser  
1 5 10 15

Pro Gly

<210> 140  
<211> 17  
<212> PRT  
<213> Homo sapiens

<400> 140  
His Leu Ser Ser Ser Met Glu Glu Ala Val Pro Asn Ser Cys Ser Pro  
1 5 10 15

Gly

<210> 141  
<211> 34  
<212> PRT  
<213> Homo sapiens

<400> 141  
Phe Asn Cys Glu Ser Cys Ser Val Thr Leu Ala Gly Val Gln Trp Arg  
1 5 10 15

Asp Leu Gly Leu Leu Gln Pro Leu Pro Pro Lys Phe Lys Arg Phe Ser  
                   20                                  25                                  30

Cys Leu

<210> 142  
 <211> 34  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (15)  
 <223> Xaa equals stop translation

<400> 142  
 Phe Lys Thr Glu Ser His Ser Val Pro Gln Ala Gly Met Gln Xaa Cys  
       1                                  5                                  10                                  15

Asp Leu Gly Ser Leu Gln Pro Ser Pro Pro Arg Phe Lys His Phe Ser  
                   20                                  25                                  30

Cys Leu

<210> 143  
 <211> 36  
 <212> PRT  
 <213> Homo sapiens

<400> 143  
 Val Thr Leu Ala Gly Val Gln Trp Arg Asp Leu Gly Leu Leu Gln Pro  
       1                                  5                                  10                                  15

Leu Pro Pro Lys Phe Lys Arg Phe Ser Cys Leu Ser Phe Pro Ser Ser  
                   20                                  25                                  30

Trp Asp Tyr Arg  
                   35

<210> 144  
 <211> 36  
 <212> PRT  
 <213> Homo sapiens

<400> 144  
 Ile Ala Gln Ala Gly Val Gln Trp Arg Asp Leu Gly Ser Pro Gln Thr  
       1                                  5                                  10                                  15

Pro Pro Leu Gly Leu Lys Arg Phe Ser Cys Leu Ser Leu Thr Ser Ser

20 25 30

Trp Asp Tyr Arg  
35

<210> 145  
<211> 36  
<212> PRT  
<213> Homo sapiens

<400> 145  
Cys Arg Met Arg Glu Ala Ser Ala Gly Asp Gln Gly His Met Val Cys  
1 5 10 15  
Pro Cys Gly Glu His Cys Gly Cys Asn Pro Cys Asn Cys Pro Lys Thr  
20 25 30  
Gln Thr Gln Thr  
35

<210> 146  
<211> 40  
<212> PRT  
<213> Homo sapiens

<220>  
<221> SITE  
<222> (25)  
<223> Xaa equals stop translation

<400> 146  
Cys Thr Leu Pro Gln Ala Arg Gly Gly Phe Ser Pro His Pro Leu Cys  
1 5 10 15  
Pro Pro Asn Leu Gly Cys Thr Phe Xaa Glu Gly Asp Ser Pro Cys Pro  
20 25 30  
Cys Pro Leu Pro Gln Asn Lys Thr  
35 40

<210> 147  
<211> 98  
<212> PRT  
<213> Homo sapiens

<400> 147  
Met Asp Gly Pro Pro Glu Trp Asn Gln Gly Gly Thr Leu Gly Arg Gly  
1 5 10 15  
His Gly Ser Val Ala Ser Val Ser Asp Val Arg Asn Gln Asn Gln Asp  
20 25 30

Pro Arg Arg Tyr Gly Lys Val Pro Arg Thr Ala Ser Thr Pro Asn Ala  
                   35                                  40                                  45

Ala Ala Leu Leu Arg Gln Ser Leu Asn Gly Ser Ala Ser Gly Pro Pro  
           50                                  55                                  60

Thr Asn His Pro Ser Pro Ser Thr Pro Pro Glu Ser Gly Leu Ser Ser  
   65                                  70                                  75                                  80

Ala Val Pro Ser Arg Pro Gly Ser Pro Gly Gly Ser Lys Asn Gly Asp  
                                   85                                  90                                  95

Pro Asn

<210> 148  
 <211> 87  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (57)  
 <223> Xaa equals stop translation

<220>  
 <221> SITE  
 <222> (69)  
 <223> Xaa equals stop translation

<220>  
 <221> SITE  
 <222> (73)  
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 148  
 Leu Asp Gly Pro Ala Glu Gly Arg Gln Gly Thr Gln Ile Ser Ser Ser  
   1                                  5                                  10                                  15

Arg Cys Pro Asn Ser Ile His Arg Val Ile Ser Arg Leu Pro His Ser  
                                   20                                  25                                  30

Lys Asp Ile Gln Asn Ala Glu Gly Ala His Arg Ser Ala Val Ser Gly  
           35                                  40                                  45

Val Thr Ser Gln Pro Pro Ser Leu Xaa Pro Leu Glu Ala His Ser Pro  
           50                                  55                                  60

Gly Leu Ser Phe Xaa Thr Pro Ser Xaa Gly Leu Pro Gly Cys Pro Leu  
   65                                  70                                  75                                  80

Gly Thr Ala Gly Pro Pro Ser  
                                   85



<210> 149  
 <211> 82  
 <212> PRT  
 <213> Homo sapiens

<400> 149  
 His Leu Arg Ser Gly Val Gln Asp Tyr Pro Gly Gln His Gly Lys Ile  
   1                  5                  10                  15  
 Pro Ser Leu Leu Lys Ile Gln Glu Leu Ala Gly His Gly Gly Arg Cys  
                   20                  25                  30  
 Leu Gln Ser Gln Leu Leu Arg Arg Leu Arg Gln Glu Asn His Leu Asn  
                   35                  40                  45  
 Ser Gly Gly Arg Gly Cys Ser Glu Pro Lys Ser His Leu Cys Ile Pro  
                   50                  55                  60  
 Ala Trp Val Thr Glu Gly Asp Ser Val Ser Lys Gln Asn Lys Thr Lys  
   65                  70                  75                  80  
 Asn Glu

<210> 150  
 <211> 81  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> SITE  
 <222> (7)  
 <223> Xaa equals stop translation

<220>  
 <221> SITE  
 <222> (34)  
 <223> Xaa equals stop translation

<400> 150  
 His Leu Lys Ser Gly Ile Xaa Asp Gln Pro Gly Gln His Gly Glu Thr  
   1                  5                  10                  15  
 Pro Ser Leu Leu Lys Ile Gln Lys Leu Ala Arg His Gly Ser Arg Ser  
                   20                  25                  30  
 Leu Xaa Ser Gln Leu Phe Gly Arg Leu Arg Gln Glu Asn His Leu Asn  
                   35                  40                  45  
 Leu Gly Gly Gly Gly Cys Ser Glu Leu Arg Ser Arg Asn Cys Ala Pro  
   50                  55                  60  
 Ala Trp Ala Thr Arg Ala Lys Leu His Leu Lys Glu Lys Thr Lys Gln

65                      70                      75                      80

Asp

<210> 151

<211> 46

<212> PRT

<213> Homo sapiens

<400> 151

Asp Gln Ser Leu Ala Leu Ser Pro Lys Leu Glu Cys Ser Gly Thr Ile  
1 5 10 15

Leu Ala His Ser Asn Leu Arg Leu Leu Gly Ser Ser Asp Ser Pro Ala  
20 25 30

Ser Ala Ser Arg Val Ala Gly Ile Thr Gly Val Cys His His  
35 40 45

<210> 152

<211> 44

<212> PRT

<213> Homo sapiens

<400> 152

Glu Gln Gly Leu Ala Leu Leu Pro Arg Leu Glu Gly Val Ile Thr Ala  
1 5 10 15

His Cys Ile Leu Gln Leu Leu Gly Ser Arg Asp Pro Pro Thr Ser Ala  
20 25 30

Ser Gln Ser Ala Gly Val Thr Gly Thr Tyr His His  
35 40

<210> 153

<211> 41

<212> PRT

<213> Homo sapiens

<400> 153

Pro Ser Gln Ser Leu Gln Tyr Gln Lys Glu Ser Val Glu Arg Ala Leu  
1 5 10 15

Thr Cys Ala Asn Cys Gly Gln Lys Leu His Val Leu Glu Val His Val  
20 25 30

Cys Glu Gly Gly Cys Ala Glu Leu Met  
35 40

<210> 154

&lt;211&gt; 47

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 154

Pro Ser Gln Ala Thr Pro Ala Ser Leu Asp Gln Asp Lys Trp Ala Ala  
 1 5 10 15

Gln Gln Pro Val Phe Asn Arg Lys Cys Gly Gln Lys Ile Ser Thr His  
 20 25 30

Ser Thr Glu Tyr His Cys Cys Asp Gly Asp Cys Val Ala Val Val  
 35 40 45

&lt;210&gt; 155

&lt;211&gt; 25

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 155

Phe Ser Phe Arg Leu Leu Pro Pro Leu Ala Gly Arg Leu Thr Ser Ser  
 1 5 10 15

Val Lys Ser Ala Phe Gln Phe Glu Cys  
 20 25

&lt;210&gt; 156

&lt;211&gt; 27

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 156

Phe Ser Met Arg Leu Thr Ser Leu Pro Pro Cys Ala Gly Asn Ala Thr  
 1 5 10 15

Asp Phe Val Ser Ser Val Phe Phe Phe Pro Cys  
 20 25

&lt;210&gt; 157

&lt;211&gt; 67

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 157

Leu Arg Ser Gly Val Gln Asp Tyr Pro Gly Gln His Gly Lys Ile Pro  
 1 5 10 15

Ser Leu Leu Lys Ile Gln Glu Leu Ala Gly His Gly Gly Arg Cys Leu  
 20 25 30

Gln Ser Gln Leu Leu Arg Arg Leu Arg Gln Glu Asn His Leu Asn Ser  
 35 40 45

Gly Gly Arg Gly Cys Ser Glu Pro Lys Ser His Leu Cys Ile Pro Ala  
 50 55 60

Trp Val Thr  
 65

<210> 158  
 <211> 67  
 <212> PRT  
 <213> Homo sapiens

<400> 158  
 Leu Arg Ser Gly Val Arg Asp Gln Pro Gly Gln Tyr Gly Glu Thr Pro  
 1 5 10 15

Ser Leu Leu Lys Ile Gln Lys Leu Ala Arg His Gly Gly Ile Cys Met  
 20 25 30

Cys Ser Gln Leu Leu Gly Arg Leu Arg Gln Glu Asn Cys Leu Asn Pro  
 35 40 45

Gly Gly Arg Val Cys Ser Glu Pro Arg Ser His Arg Cys Thr Thr Ala  
 50 55 60

Trp Val Thr  
 65

<210> 159  
 <211> 131  
 <212> PRT  
 <213> Homo sapiens

<400> 159  
 Gly Gln Glu Arg Phe Thr Ser Met Thr Arg Leu Tyr Tyr Arg Asp Ala  
 1 5 10 15

Ser Ala Cys Val Ile Met Phe Asp Val Thr Asn Ala Thr Thr Phe Ser  
 20 25 30

Asn Ser Gln Arg Trp Lys Gln Asp Leu Asp Ser Lys Leu Thr Leu Pro  
 35 40 45

Asn Gly Glu Pro Val Pro Cys Leu Leu Leu Ala Asn Lys Cys Asp Leu  
 50 55 60

Ser Pro Trp Ala Val Ser Arg Asp Gln Ile Asp Arg Phe Ser Lys Glu  
 65 70 75 80

Asn Gly Phe Thr Gly Trp Thr Glu Thr Ser Val Lys Glu Asn Lys Asn  
 85 90 95

Ile Asn Glu Ala Met Arg Val Leu Ile Glu Lys Met Met Arg Asn Ser

100							105				110				
Thr	Glu	Asp	Ile	Met	Ser	Leu	Ser	Thr	Gln	Gly	Asp	Tyr	Ile	Asn	Leu
115							120				125				

Gln Thr Lys  
130

```
<210> 160
<211> 131
<212> PRT
<213> Homo sapiens
```

```
<400> 160
Gly Gln Glu Arg Phe Thr Ser Met Thr Arg Leu Tyr Tyr Arg Asp Ala
   1                   5               10             15
```

Ser Ala Cys Val Ile Met Phe Asp Val Thr Asn Ala Thr Thr Phe Ser  
20 25 30

Asn Ser Gln Arg Trp Lys Gln Asp Leu Asp Ser Lys Leu Thr Leu Pro  
35 40 45

Asn Gly Glu Pro Val Pro Cys Leu Leu Leu Ala Asn Lys Cys Asp Leu  
50 55 60

Ser Pro Trp Ala Val Ser Arg Asp Gln Ile Asp Arg Phe Ser Lys Glu  
65 70 75 80

Asn Gly Phe Thr Gly Trp Thr Glu Thr Ser Val Lys Glu Asn Lys Asn  
85 90 95

Ile Asn Glu Ala Met Arg Val Leu Ile Glu Lys Met Met Arg Asn Ser  
100 105 110

Thr Glu Asp Ile Met Ser Leu Ser Thr Gln Gly Asp Tyr Ile Asn Leu  
115 120 125

Gln Thr Lys  
130

```
<210> 161
<211> 35
<212> PRT
<213> Homo sapiens
```

```
<400> 161
Ile Tyr Phe Ser Trp Trp Ser Lys Phe Leu Thr Ile Tyr Phe Ser Ser
  1             5             10             15
```

Arg Phe Tyr Ile Ile Ile Cys Val Ser Val Thr Asn Ser Thr Phe Lys  
20 25 30

Leu Thr Phe  
35

<210> 162  
<211> 36  
<212> PRT  
<213> Homo sapiens

<400> 162  
Val Val His Gly Trp Trp Ser Thr Ala Thr Ile Thr Tyr Trp Asn Ser  
1 5 10 15  
Asp Arg Phe Thr Val Leu Thr Ala Leu Lys Ile Lys Leu Asp Thr Ala  
20 25 30

Ala Val Thr Phe  
35

<210> 163  
<211> 429  
<212> PRT  
<213> Homo sapiens

<400> 163  
Val Arg Met Phe Phe His Phe Ser Glu Ile Leu Asp Gly Asn Gln Leu  
1 5 10 15  
His Ile Ala Asp Glu Val Glu Phe Thr Val Val Pro Asp Met Leu Ser  
20 25 30  
Ala Gln Arg Asn His Ala Ile Arg Ile Lys Lys Leu Pro Lys Gly Thr  
35 40 45  
Val Ser Phe His Ser His Ser Asp His Arg Phe Leu Gly Thr Val Glu  
50 55 60  
Lys Glu Ala Thr Phe Ser Asn Pro Lys Thr Thr Ser Pro Asn Lys Gly  
65 70 75 80  
Lys Glu Lys Glu Ala Glu Asp Gly Ile Ile Ala Tyr Asp Asp Cys Gly  
85 90 95  
Val Lys Leu Thr Ile Ala Phe Gln Ala Lys Asp Val Glu Gly Ser Thr  
100 105 110  
Ser Pro Gln Ile Gly Asp Lys Val Glu Phe Ser Ile Ser Asp Lys Gln  
115 120 125  
Arg Pro Gly Gln Gln Val Ala Thr Cys Val Arg Leu Leu Gly Arg Asn  
130 135 140  
Ser Asn Ser Lys Arg Leu Leu Gly Tyr Val Ala Thr Leu Lys Asp Asn  
145 150 155 160

Phe Gly Phe Ile Glu Thr Ala Asn His Asp Lys Glu Ile Phe Phe His  
 165 170 175  
 Tyr Ser Glu Phe Ser Gly Asp Val Asp Ser Leu Glu Leu Gly Asp Met  
 180 185 190  
 Val Glu Tyr Ser Leu Ser Lys Gly Lys Gly Asn Lys Val Ser Ala Glu  
 195 200 205  
 Lys Val Asn Lys Thr His Ser Val Asn Gly Ile Thr Glu Glu Ala Asp  
 210 215 220  
 Pro Thr Ile Tyr Ser Gly Lys Val Ile Arg Pro Leu Arg Ser Val Asp  
 225 230 235 240  
 Pro Thr Gln Thr Glu Tyr Gln Gly Met Ile Glu Ile Val Glu Glu Gly  
 245 250 255  
 Asp Met Lys Gly Glu Val Tyr Pro Phe Gly Ile Val Gly Met Ala Asn  
 260 265 270  
 Lys Gly Asp Cys Leu Gln Lys Gly Glu Ser Val Lys Phe Gln Leu Cys  
 275 280 285  
 Val Leu Gly Gln Asn Ala Gln Thr Met Ala Tyr Asn Ile Thr Pro Leu  
 290 295 300  
 Arg Arg Ala Thr Val Glu Cys Val Lys Asp Gln Phe Gly Phe Ile Asn  
 305 310 315 320  
 Tyr Glu Val Gly Asp Ser Lys Lys Leu Phe Phe His Val Lys Glu Val  
 325 330 335  
 Gln Asp Gly Ile Glu Leu Gln Ala Gly Asp Glu Val Glu Phe Ser Val  
 340 345 350  
 Ile Leu Asn Gln Arg Thr Gly Lys Cys Ser Ala Cys Asn Val Trp Arg  
 355 360 365  
 Val Cys Glu Gly Pro Lys Ala Val Ala Ala Pro Arg Pro Asp Arg Leu  
 370 375 380  
 Val Asn Arg Leu Lys Asn Ile Thr Leu Asp Asp Ala Ser Ala Pro Arg  
 385 390 395 400  
 Leu Met Val Leu Arg Gln Pro Arg Gly Pro Asp Asn Ser Met Gly Phe  
 405 410 415  
 Gly Ala Glu Arg Lys Ile Arg Gln Ala Gly Val Ile Asp  
 420 425

&lt;210&gt; 164

&lt;211&gt; 429

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 164

Val	Arg	Met	Phe	Phe	His	Phe	Ser	Glu	Ile	Leu	Asp	Gly	Asn	Gln	Leu
1				5					10					15	
His	Ile	Ala	Asp	Glu	Val	Glu	Phe	Thr	Val	Val	Pro	Asp	Met	Leu	Ser
			20					25					30		
Ala	Gln	Arg	Asn	His	Ala	Ile	Arg	Ile	Lys	Lys	Leu	Pro	Lys	Gly	Thr
		35					40						45		
Val	Ser	Phe	His	Ser	His	Ser	Asp	His	Arg	Phe	Leu	Gly	Thr	Val	Glu
	50					55					60				
Lys	Glu	Ala	Thr	Phe	Ser	Asn	Pro	Lys	Thr	Thr	Ser	Pro	Asn	Lys	Gly
65					70					75					80
Thr	Glu	Lys	Glu	Ala	Glu	Asp	Gly	Ile	Ile	Ala	Tyr	Asp	Asp	Cys	Gly
				85					90					95	
Val	Lys	Leu	Thr	Ile	Ala	Phe	Gln	Ala	Lys	Asp	Val	Glu	Gly	Ser	Thr
		100						105					110		
Ser	Pro	Gln	Ile	Gly	Asp	Lys	Val	Glu	Phe	Ser	Ile	Ser	Asp	Lys	Gln
		115					120					125			
Arg	Pro	Gly	Gln	Gln	Val	Ala	Thr	Cys	Val	Arg	Leu	Leu	Gly	Arg	Asn
		130				135						140			
Ser	Asn	Ser	Lys	Arg	Leu	Leu	Gly	Tyr	Val	Ala	Thr	Leu	Lys	Asp	Asn
145					150					155					160
Phe	Gly	Phe	Ile	Glu	Thr	Ala	Asn	His	Asp	Lys	Glu	Ile	Phe	Phe	His
			165						170					175	
Tyr	Ser	Glu	Phe	Ser	Gly	Asp	Val	Asp	Ser	Leu	Glu	Leu	Gly	Asp	Met
		180						185					190		
Val	Glu	Tyr	Ser	Leu	Ser	Lys	Gly	Lys	Gly	Asn	Lys	Val	Ser	Ala	Glu
		195					200					205			
Lys	Val	Asn	Lys	Thr	His	Ser	Val	Asn	Gly	Ile	Thr	Glu	Glu	Ala	Asp
	210					215					220				
Pro	Thr	Ile	Tyr	Ser	Gly	Lys	Val	Ile	Arg	Pro	Leu	Arg	Ser	Val	Asp
225					230					235					240
Pro	Thr	Gln	Thr	Glu	Tyr	Gln	Gly	Met	Ile	Glu	Ile	Val	Glu	Glu	Gly
			245						250					255	
Asp	Met	Lys	Gly	Glu	Val	Tyr	Pro	Phe	Gly	Ile	Val	Gly	Met	Ala	Asn
		260						265					270		



Lys Gly Asp Cys Leu Gln Lys Gly Glu Ser Val Lys Phe Gln Leu Cys  
           275                                  280                                  285  
 Val Leu Gly Gln Asn Ala Gln Thr Met Ala Tyr Asn Ile Thr Pro Leu  
           290                                  295                                  300  
 Arg Arg Ala Thr Val Glu Cys Val Lys Asp Gln Phe Gly Phe Ile Asn  
 305                                  310                                  315                                  320  
 Tyr Glu Val Gly Asp Ser Lys Lys Leu Phe Phe His Val Lys Glu Val  
                                   325                                  330                                  335  
 Gln Asp Gly Ile Glu Leu Gln Ala Gly Asp Glu Val Glu Phe Ser Val  
                                   340                                  345                                  350  
 Ile Leu Asn Gln Arg Thr Gly Lys Cys Ser Ala Cys Asn Val Trp Arg  
                                   355                                  360                                  365  
 Val Cys Glu Gly Pro Lys Ala Val Ala Ala Pro Arg Pro Asp Arg Leu  
           370                                  375                                  380  
 Val Asn Arg Leu Lys Asn Ile Thr Leu Asp Asp Ala Ser Ala Pro Arg  
 385                                  390                                  395                                  400  
 Leu Met Val Leu Arg Gln Pro Arg Gly Pro Asp Asn Ser Met Gly Phe  
                                   405                                  410                                  415  
 Gly Ala Glu Arg Lys Ile Arg Gln Ala Gly Val Ile Asp  
                                   420                                  425

<210> 165  
 <211> 258  
 <212> PRT  
 <213> Homo sapiens

<400> 165  
 Leu Thr Tyr Thr Pro Glu Asp Val Glu Gly Asn Val Gln Leu Glu Thr  
   1                                  5                                  10                                  15  
 Gly Asp Lys Ile Asn Phe Val Ile Asp Asn Asn Lys His Thr Gly Ala  
           20                                  25                                  30  
 Val Ser Ala Arg Asn Ile Met Leu Leu Lys Lys Lys Gln Ala Arg Cys  
           35                                  40                                  45  
 Gln Gly Val Val Cys Ala Met Lys Glu Ala Phe Gly Phe Ile Glu Arg  
           50                                  55                                  60  
 Gly Asp Val Val Lys Glu Ile Phe Phe His Tyr Ser Glu Phe Lys Gly  
   65                                  70                                  75                                  80  
 Asp Leu Glu Thr Leu Gln Pro Gly Asp Asp Val Glu Phe Thr Ile Lys  
                                   85                                  90                                  95

Asp Arg Asn Gly Lys Glu Val Ala Thr Asp Val Arg Leu Leu Pro Gln  
 100 105 110  
 Gly Thr Val Ile Phe Glu Asp Ile Ser Ile Glu His Phe Glu Gly Thr  
 115 120 125  
 Val Thr Lys Val Ile Pro Lys Val Pro Ser Lys Asn Gln Asn Asp Pro  
 130 135 140  
 Leu Pro Gly Arg Ile Lys Val Asp Phe Val Ile Pro Lys Glu Leu Pro  
 145 150 155 160  
 Phe Gly Asp Lys Asp Thr Lys Ser Lys Val Thr Leu Leu Glu Gly Asp  
 165 170 175  
 His Val Arg Phe Asn Ile Ser Thr Asp Arg Arg Asp Lys Leu Glu Arg  
 180 185 190  
 Ala Thr Asn Ile Glu Val Leu Ser Asn Thr Phe Gln Phe Thr Asn Glu  
 195 200 205  
 Ala Arg Glu Met Gly Val Ile Ala Ala Met Arg Asp Gly Phe Gly Phe  
 210 215 220  
 Ile Lys Cys Val Asp Arg Asp Val Arg Met Phe Phe His Phe Ser Glu  
 225 230 235 240  
 Ile Leu Asp Gly Asn Gln Leu His Ile Ala Asp Glu Val Glu Phe Thr  
 245 250 255  
 Val Val

<210> 166  
 <211> 253  
 <212> PRT  
 <213> Homo sapiens

<400> 166  
 Ile Ala Phe Gln Ala Lys Asp Val Glu Gly Ser Thr Ser Pro Gln Ile  
 1 5 10 15  
 Gly Asp Lys Val Glu Phe Ser Ile Ser Asp Lys Gln Arg Pro Gly Gln  
 20 25 30  
 Gln Val Ala Thr Cys Val Arg Leu Leu Gly Arg Asn Ser Asn Ser Lys  
 35 40 45  
 Arg Leu Leu Gly Tyr Val Ala Thr Leu Lys Asp Asn Phe Gly Phe Ile  
 50 55 60  
 Glu Thr Ala Asn His Asp Lys Glu Ile Phe Phe His Tyr Ser Glu Phe  
 65 70 75 80

Ser Gly Asp Val Asp Ser Leu Glu Leu Gly Asp Met Val Glu Tyr Ser  
                             85                            90                            95  
 Leu Ser Lys Gly Lys Gly Asn Lys Val Ser Ala Glu Lys Val Asn Lys  
                             100                            105                            110  
 Thr His Ser Val Asn Gly Ile Thr Glu Glu Ala Asp Pro Thr Ile Tyr  
                             115                            120                            125  
 Ser Gly Lys Val Ile Arg Pro Leu Arg Ser Val Asp Pro Thr Gln Thr  
                             130                            135                            140  
 Glu Tyr Gln Gly Met Ile Glu Ile Val Glu Glu Gly Asp Met Lys Gly  
                             145                            150                            155                            160  
 Glu Val Tyr Pro Phe Gly Ile Val Gly Met Ala Asn Lys Gly Asp Cys  
                             165                            170                            175  
 Leu Gln Lys Gly Glu Ser Val Lys Phe Gln Leu Cys Val Leu Gly Gln  
                             180                            185                            190  
 Asn Ala Gln Thr Met Ala Tyr Asn Ile Thr Pro Leu Arg Arg Ala Thr  
                             195                            200                            205  
 Val Glu Cys Val Lys Asp Gln Phe Gly Phe Ile Asn Tyr Glu Val Gly  
                             210                            215                            220  
 Asp Ser Lys Lys Leu Phe Phe His Val Lys Glu Val Gln Asp Gly Ile  
                             225                            230                            235                            240  
 Glu Leu Gln Ala Gly Asp Glu Val Glu Phe Ser Val Ile  
                             245                            250

&lt;210&gt; 167

&lt;211&gt; 202

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 167

Met Cys Tyr Gly Lys Cys Ala Arg Cys Ile Gly His Ser Leu Val Gly  
           1                            5                            10                            15  
 Leu Ala Leu Leu Cys Ile Ala Ala Asn Ile Leu Leu Tyr Phe Pro Asn  
                             20                            25                            30  
 Gly Glu Thr Lys Tyr Ala Ser Glu Asn His Leu Ser Arg Phe Val Trp  
                             35                            40                            45  
 Phe Phe Ser Gly Ile Val Gly Gly Gly Leu Leu Met Leu Leu Pro Ala  
           50                            55                            60  
 Phe Val Phe Ile Gly Leu Glu Gln Asp Asp Cys Cys Gly Cys Cys Gly  
           65                            70                            75                            80

His Glu Asn Cys Gly Lys Arg Cys Ala Met Leu Ser Ser Val Leu Ala  
                             85                            90                            95  
 Ala Leu Ile Gly Ile Ala Gly Ser Gly Tyr Cys Val Ile Val Ala Ala  
                             100                            105                            110  
 Leu Gly Leu Ala Glu Gly Pro Leu Cys Leu Asp Ser Leu Gly Gln Trp  
                             115                            120                            125  
 Asn Tyr Thr Phe Ala Ser Thr Glu Gly Gln Tyr Leu Leu Asp Thr Ser  
                             130                            135                            140  
 Thr Trp Ser Glu Cys Thr Glu Pro Lys His Ile Val Glu Trp Asn Val  
                             145                            150                            155                            160  
 Ser Leu Phe Ser Ile Leu Leu Ala Leu Gly Gly Ile Glu Phe Ile Leu  
                             165                            170                            175  
 Cys Leu Ile Gln Val Ile Asn Gly Val Leu Gly Gly Ile Cys Gly Phe  
                             180                            185                            190  
 Cys Cys Ser His Gln Gln Gln Tyr Asp Cys  
                             195                            200

&lt;210&gt; 168

&lt;211&gt; 202

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (115)

&lt;223&gt; Xaa equals stop translation

&lt;400&gt; 168

Met Cys Tyr Gly Lys Cys Ala Arg Cys Ile Gly His Ser Leu Val Gly  
           1                            5                            10                            15  
 Leu Ala Leu Leu Cys Ile Ala Ala Asn Ile Leu Leu Tyr Phe Pro Asn  
                             20                            25                            30  
 Gly Glu Thr Lys Tyr Ala Ser Glu Asn His Leu Ser Arg Phe Val Trp  
                             35                            40                            45  
 Phe Phe Ser Gly Ile Val Gly Gly Gly Leu Leu Met Leu Leu Pro Ala  
           50                            55                            60  
 Phe Val Phe Ile Gly Leu Glu Gln Asp Asp Cys Cys Gly Cys Cys Gly  
           65                            70                            75                            80  
 His Glu Asn Cys Gly Lys Arg Cys Ala Met Leu Ser Ser Val Leu Ala  
                             85                            90                            95  
 Ala Leu Ile Gly Leu Gln Asp Leu Ala Thr Val Ser Leu Trp Gln Pro

	100		105		110										
Leu	Ala	Xaa	Gln	Lys	Asp	His	Tyr	Val	Leu	Ile	Pro	Ser	Ala	Ser	Gly
	115						120					125			
Thr	Thr	Pro	Leu	Pro	Ser	Thr	Glu	Gly	Gln	Tyr	Leu	Leu	Asp	Thr	Ser
	130					135					140				
Thr	Trp	Ser	Glu	Cys	Thr	Glu	Pro	Lys	His	Ile	Val	Glu	Trp	Asn	Val
145					150					155					160
Ser	Leu	Phe	Ser	Ile	Leu	Leu	Ala	Leu	Gly	Gly	Ile	Glu	Phe	Ile	Leu
				165					170					175	
Cys	Leu	Ile	Gln	Val	Ile	Asn	Gly	Val	Leu	Gly	Gly	Ile	Cys	Gly	Phe
			180					185					190		
Cys	Cys	Ser	His	Gln	Gln	Gln	Tyr	Asp	Cys						
	195						200								

&lt;210&gt; 169

&lt;211&gt; 87

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 169

Met	Tyr	Cys	Leu	Gln	Trp	Leu	Leu	Pro	Val	Leu	Leu	Ile	Pro	Lys	Pro
1				5					10					15	
Leu	Asn	Pro	Ala	Leu	Trp	Phe	Ser	His	Ser	Met	Phe	Met	Gly	Phe	Tyr
			20					25					30		
Leu	Leu	Ser	Phe	Leu	Leu	Glu	Arg	Lys	Pro	Cys	Thr	Ile	Cys	Ala	Leu
		35					40					45			
Val	Phe	Leu	Ala	Ala	Leu	Phe	Leu	Ile	Cys	Tyr	Ser	Cys	Trp	Gly	Asn
	50					55					60				
Cys	Phe	Leu	Tyr	His	Cys	Ser	Asp	Ser	Pro	Leu	Pro	Glu	Ser	Ala	His
65					70					75					80
Asp	Pro	Gly	Val	Val	Gly	Thr									
				85											

&lt;210&gt; 170

&lt;211&gt; 87

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 170

Met	Tyr	Cys	Leu	Gln	Trp	Leu	Leu	Pro	Val	Leu	Leu	Ile	Pro	Lys	Pro
1				5					10					15	

Leu Asn Pro Ala Leu Trp Phe Ser His Ser Met Phe Met Gly Phe Tyr  
                   20                  25                  30

Leu Leu Ser Phe Leu Leu Glu Arg Lys Pro Cys Thr Ile Cys Ala Leu  
           35                  40                  45

Val Phe Leu Ala Ala Leu Phe Leu Ile Cys Tyr Ser Cys Trp Gly Asn  
       50                  55                  60

Cys Phe Leu Tyr His Cys Ser Asp Ser Pro Leu Pro Glu Ser Ala His  
       65                  70                  75                  80

Asp Pro Gly Val Val Gly Thr  
                   85

<210> 171

<211> 35

<212> PRT

<213> Homo sapiens

<400> 171

Val Lys Gly Thr Arg Lys Asn Tyr Gln Arg Leu Trp Arg Trp Gly Thr  
       1                  5                  10                  15

Met Leu Leu Gly Met Leu Met Ile Cys Ser Val Ala Gly Asn Leu Trp  
           20                  25                  30

Val Thr Val  
           35

<210> 172

<211> 35

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (20)

<223> Xaa equals stop translation

<400> 172

Leu Lys Ser His Lys Lys Asn Tyr Asn Met Leu Trp Ile Phe Lys Trp  
       1                  5                  10                  15

Gln Val Leu Xaa Leu Phe Phe Val Cys Lys Ile Leu Ser Asn His Trp  
           20                  25                  30

Glu Asn Ile  
           35

<210> 173

<211> 299

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 173

Glu	Gly	Phe	Gly	His	Arg	Pro	His	Lys	Asp	Leu	Trp	Ala	Asn	Lys	Ser	1	5	10	15
Glu	Asn	Glu	Glu	Thr	Ser	Glu	Arg	Ser	Pro	Gln	Ser	Pro	Asp	Val	Gly	20	25	30	
Glu	Ile	Pro	His	Asn	Ser	Gly	Val	Glu	Asp	Ala	Pro	Ile	Ser	Lys	Val	35	40	45	
Gly	Val	Leu	Ala	Ala	Asn	Met	Glu	Ala	Lys	Ala	Ser	Pro	Gln	Arg	Glu	50	55	60	
Lys	Glu	Asp	Lys	Pro	Ala	Glu	Thr	Lys	Lys	Leu	Arg	Ile	Ala	Trp	Pro	65	70	75	80
Pro	Pro	Ala	Glu	Leu	Gly	Ser	Ser	Gly	Ser	Thr	Met	Glu	Glu	Gly	Phe	85	90	95	
Lys	Gly	Ser	Lys	Pro	Lys	Trp	Pro	Pro	Glu	Asp	Glu	Thr	Ser	Lys	Thr	100	105	110	
Glu	Ala	Leu	Gly	Asp	Val	Asp	Leu	Asp	Leu	Lys	Lys	Leu	Arg	Arg	Ser	115	120	125	
Ser	Ser	Leu	Lys	Glu	Arg	Ser	Arg	Pro	Phe	Thr	Val	Ala	Ala	Ser	Phe	130	135	140	
Arg	Thr	Ser	Ser	Leu	Lys	Ser	Pro	Lys	Thr	Leu	Ser	Pro	Leu	Arg	Lys	145	150	155	160
Gly	Trp	Ser	Met	Ser	Glu	Gln	Asn	Glu	Glu	Ser	Gly	Gly	Gly	Ile	Val	165	170	175	
Thr	Ala	Gly	Lys	Gln	Thr	Glu	Asn	Ala	Lys	Ala	Pro	Gly	Lys	Lys	Glu	180	185	190	
Asn	Val	Arg	Lys	Ser	Arg	Trp	Gln	Arg	Glu	Asp	Ser	Ala	Arg	Glu	Glu	195	200	205	
Val	Ala	Arg	Arg	Ser	Lys	Asp	Ile	His	Gly	Val	Glu	Leu	Glu	Ser	Glu	210	215	220	
Asn	Phe	Ile	Glu	Asn	Gly	Ala	Asn	Ile	Tyr	Glu	Gly	Asp	Ser	Gly	Asp	225	230	235	240
Leu	Val	Gln	Gln	Ser	Pro	Leu	Asp	Pro	Lys	Ser	Pro	Ser	Trp	Ser	Gly	245	250	255	
Phe	Val	Asp	Thr	Thr	Pro	Thr	Glu	Glu	Phe	Thr	Thr	Gln	Asn	Gln	Lys	260	265	270	

Ser His Asp Val Gly Phe Arg Glu Gly Glu Val Val Arg Glu Leu Ser  
 275 280 285

Val Glu Glu Gln Ile Lys Arg Asn Arg Tyr Tyr  
 290 295

<210> 174

<211> 299

<212> PRT

<213> Homo sapiens

<400> 174

Glu Gly Phe Gly His Arg Pro His Lys Asp Leu Trp Ala Ser Lys Asn  
 1 5 10 15

Glu Asn Glu Glu Ile Leu Glu Arg Pro Ala Gln Leu Ala Asn Ala Arg  
 20 25 30

Glu Thr Pro His Ser Pro Gly Val Glu Asp Ala Pro Ile Ala Lys Val  
 35 40 45

Gly Val Leu Ala Ala Ser Met Glu Ala Lys Ala Ser Ser Gln Gln Glu  
 50 55 60

Lys Glu Asp Lys Pro Ala Glu Thr Lys Lys Leu Arg Ile Ala Trp Pro  
 65 70 75 80

Pro Pro Thr Glu Leu Gly Ser Ser Gly Ser Ala Leu Glu Glu Gly Ile  
 85 90 95

Lys Met Ser Lys Pro Lys Trp Pro Pro Glu Asp Glu Ile Ser Lys Pro  
 100 105 110

Glu Val Pro Glu Asp Val Asp Leu Asp Leu Lys Lys Leu Arg Arg Ser  
 115 120 125

Ser Ser Leu Lys Glu Arg Ser Arg Pro Phe Thr Val Ala Ala Ser Phe  
 130 135 140

Gln Ser Thr Ser Val Lys Ser Pro Lys Thr Val Ser Pro Pro Ile Arg  
 145 150 155 160

Lys Gly Trp Ser Met Ser Glu Gln Ser Glu Glu Ser Val Gly Gly Arg  
 165 170 175

Val Ala Glu Arg Lys Gln Val Glu Asn Ala Lys Ala Ser Lys Lys Asn  
 180 185 190

Gly Asn Val Gly Lys Thr Thr Trp Gln Asn Lys Glu Ser Lys Gly Glu  
 195 200 205

Thr Gly Lys Arg Ser Lys Glu Gly His Ser Leu Glu Met Glu Asn Glu  
 210 215 220



Asn Leu Val Glu Asn Gly Ala Asp Ser Asp Glu Asp Asp Asn Ser Phe  
225 230 235 240

Leu Lys Gln Gln Ser Pro Gln Glu Pro Lys Ser Leu Asn Trp Ser Ser  
245 250 255

Phe Val Asp Asn Thr Phe Ala Glu Glu Phe Thr Thr Gln Asn Gln Lys  
260 265 270

Ser Gln Asp Val Glu Leu Trp Glu Gly Glu Val Val Lys Glu Leu Ser  
275 280 285

Val Glu Glu Gln Ile Lys Arg Asn Arg Tyr Tyr  
290 295

<210> 175

<211> 47

<212> PRT

<213> Homo sapiens

<400> 175

Tyr Ile Cys Ile Ser Ile Tyr Leu Ser Ile Tyr Leu Ser Ile Ser Ile  
1 5 10 15

Tyr Leu Ser Ile Tyr Thr Tyr Ile His Thr His Thr His Thr His Thr  
20 25 30

Tyr Ile Tyr Ile Tyr Ile Tyr Ile Tyr Ile Tyr Leu Tyr Leu Cys  
35 40 45

<210> 176

<211> 48

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (34)

<223> Xaa equals stop translation

 $\langle 220 \rangle$ 

<221> SITE

<222> (43)

<223> Xaa equals stop translation

<400> 176

Tyr Ile Cys Val Tyr Leu Tyr Val Cys Thr His Pro Cys Ile Tyr Ile  
1 5 10 15

Tyr Leu Ser Val Gln Thr Leu His Met Tyr Thr His Tyr Leu Cys Lys  
20 25 30

Ile Xaa Tyr Met Tyr Asn Ala Tyr Lys Phe Xaa Gln Val Tyr Leu Cys

35

40

45

<210> 177  
 <211> 20  
 <212> PRT  
 <213> Homo sapiens

<400> 177  
 Trp Leu Gly Ala Val Ala His Val Cys Asn Pro Ser Thr Leu Gly Gly  
     1                    5                    10                    15  
 Gln Val Gly Arg  
                     20

<210> 178  
 <211> 20  
 <212> PRT  
 <213> Homo sapiens

<400> 178  
 Trp Leu Gly Val Met Ala Cys Thr Cys Asn Ser Ser Thr Ser Gly Gly  
     1                    5                    10                    15  
 Gln Gly Gly Arg  
                     20

<210> 179  
 <211> 92  
 <212> PRT  
 <213> Homo sapiens

<400> 179  
 Gln Thr Ile Tyr Lys Cys Gln Lys Cys Ala Ser Ile Lys Pro Asp Arg  
     1                    5                    10                    15  
 Ala His His Cys Ser Val Cys Glu Arg Cys Ile Arg Arg Met Asp His  
                     20                    25                    30  
 His Cys Pro Trp Val Asn Asn Cys Val Gly Glu Gly Asn Gln Lys Phe  
             35                    40                    45  
 Phe Val Leu Phe Thr Met Tyr Ile Ala Leu Leu Ser Met His Ala Leu  
     50                    55                    60  
 Tyr Trp Gly Val Trp Gln Phe Val Leu Cys Val Gly Ala Glu Trp Gln  
     65                    70                    75                    80  
 Gln Cys Ser Ser Leu Thr Pro Pro Ala Thr Thr Leu  
                     85                    90

&lt;210&gt; 180

&lt;211&gt; 92

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 180

Glu Val Ile Tyr Lys Cys Pro Lys Cys Cys Cys Ile Lys Pro Glu Arg  
 1 5 10 15

Ala His His Cys Ser Ile Cys Lys Arg Cys Ile Arg Lys Met Asp His  
 20 25 30

His Cys Pro Trp Val Asn Asn Cys Val Gly Glu Lys Asn Gln Arg Phe  
 35 40 45

Phe Val Leu Phe Thr Met Tyr Ile Ala Leu Ser Ser Val His Ala Leu  
 50 55 60

Ile Leu Cys Gly Phe Gln Phe Ile Ser Cys Val Arg Gly Gln Trp Thr  
 65 70 75 80

Glu Cys Ser Asp Phe Ser Pro Pro Ile Thr Val Ile  
 85 90

&lt;210&gt; 181

&lt;211&gt; 72

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 181

Phe Ala Ile Phe Thr Ser Val Met Phe Gly Thr Gln Ile Ser Ser Ile  
 1 5 10 15

Cys Asn Asp Glu Thr Thr Ile Glu Ser Met Arg Ser Arg Asn Ala Asn  
 20 25 30

Leu Asp Glu Asp Glu Arg Ser Arg Asn Asn Ala Trp Lys Asn Leu Gln  
 35 40 45

Leu Val Phe Gly Gly Pro Phe Ser Val Arg Trp Phe Asn Pro Leu Ala  
 50 55 60

Met Pro Phe Val Ala Arg Pro Thr  
 65 70

&lt;210&gt; 182

&lt;211&gt; 68

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; (52)

&lt;223&gt; Xaa equals any of the naturally occurring L-amino acids

&lt;400&gt; 182

Phe	Ser	Leu	Ser	Leu	Ala	Val	Met	Phe	Gly	Thr	Gln	Ile	His	Ser	Ile
1				5					10					15	

Cys	Asn	Asp	Glu	Thr	Glu	Ile	Glu	Arg	Leu	Lys	Ser	Glu	Lys	Pro	Thr
			20					25					30		

Trp	Glu	Arg	Arg	Leu	Arg	Trp	Glu	Gly	Met	Lys	Ser	Val	Phe	Gly	Gly
		35					40						45		

Pro	Pro	Ser	Xaa	Leu	Trp	Met	Asn	Pro	Phe	Val	Gly	Phe	Arg	Phe	Arg
	50					55						60			

Arg	Leu	Pro	Thr
65			

&lt;210&gt; 183

&lt;211&gt; 19

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 183

Pro	Ala	Pro	Gly	Leu	Pro	Gln	Gly	Pro	Cys	Gly	Ala	Asp	Cys	Gln	Pro
1				5				10						15	

Pro Ala Pro

&lt;210&gt; 184

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 184

Pro	Ser	Ile	Gly	Val	Asp	Pro	Lys	Gly	Ser	Cys	Gly	Lys	Glu	Ile	Gln
1				5				10					15		

Ser	Gln	Ser	Pro
			20

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/06830

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/04; C07K 14/00, 16/00; C12N 15/00, 15/63, 15/85, 15/86; C12Q 1/68; G01N 33/53  
US CL : 435/6, 7.1, 7.2, 69.1, 320.1, 325; 530/350, 387.1; 514/2; 536/23.5

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 7.1, 7.2, 69.1, 320.1, 325; 530/350, 387.1; 514/2; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
WEST, CAPLUS, BIOSIS, EMBASE, MEDLINE, GENBANK, SWISSPROT

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ITOH, S. et al. Molecular cloning of 25-hydroxyvitamin D-3 24-hydroxylase (Cyp-24) from mouse kidney: its inducibility by vitamin D-3. Biochim. et Biophys. Acta. 1995, Vol. 1264, pages 26-28, see whole reference.	1-12, 14-16, 21
Y	AKENO, N. et al. Mouse vitamin D-24-hydroxylase: molecular cloning, tissue distribution, and transcriptional regulation by 1alpha,25-Dihydroxyvitamin D3. Endocrinology. 1997, Vol. 138, No. 6, pages 2233-2240, see whole reference.	1-12, 14-16, 21
Y	OHYAMA, Y. et al. Cloning and expression of cDNA encoding 25-hydroxyvitamin D3 24-hydroxylase. FEBS Letters. 1991, Vol. 278, No. 2, pages 195-198, see whole reference.	1-12, 14-16, 21
Y	OHYAMA, Y. et al. Structural characterization of the gene encoding rat 25-hydroxyvitamin D3 24-hydroxylase. 1993, Vol. 32, pages 76-82, see whole reference.	1-12, 14-16, 21



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

20 JUNE 2000

Date of mailing of the international search report

13 JUL 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Joseph F. Murphy

Telephone No. 703-308-0196

# INTERNATIONAL SEARCH REPORT

1 national application No.

PCT/US00/06830

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-12 and 14-16

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/06830

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-12, 14-16 and 21, drawn to an isolated nucleic acid, a recombinant vector, a host cell, an isolated polypeptide and a method of making an isolated polypeptide.

Group II, claim 13, drawn to an antibody.

Group III, claim 17, drawn to a method for preventing, treating or ameliorating a medical condition.

Group IV, claim(s) 18-19, drawn to a method of diagnosing a pathological condition.

Group V, claim 20, drawn to a method for identifying a binding partner.

Group VI, claim 22, drawn to a method of identifying an activity in a biological assay.

Group VII, claim 23, drawn to a binding partner.

The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first recited product, an isolated polynucleotide, a isolated polypeptide, a recombinant vector, a method of preparing the isolated polypeptide. Further pursuant to 37 C.F.R. § (d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

The oligonucleotide sequences set forth in SEQ ID NO: 11-59 and the corresponding amino acid sequences which they encode, set forth in SEQ ID NO: 60-108, respectively.

The claims are deemed to correspond to the species listed above in the following manner:

claim 1: SEQ ID NO: 11-59

claim 11: SEQ ID NO: 60-108

The species listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The special technical feature of each species is its amino acid sequence and/or its encoding nucleic acid sequence. The species are structurally and functionally distinct.

**THIS PAGE BLANK (USPTO)**